

INTERNATIONAL UNION OF PURE AND APPLIED CHEMISTRY

ANALYTICAL CHEMISTRY DIVISION
COMMISSION ON SOLUBILITY DATA

SOLUBILITY DATA SERIES

Volume 49

**ESTERS WITH WATER
PART II: ESTERS 7-C TO 32-C**

SOLUBILITY DATA SERIES

Editor-in-Chief

J. W. LORIMER

*The University of Western Ontario
London, Ontario, Canada*

F. W. Getzen (USA)
*Sub-editor
Liquid/Liquid Systems*

C. L. Young (Australia)
*Sub-editor
Indexes*

EDITORIAL BOARD

M. Th. Cohen-Adad (France)

M. Salomon (USA)

L. Clever (USA)

V. P. Sazanov (Russia)

H. J. M. Grünbauer (Netherlands)

D. G. Shaw (USA)

A. L. Horvath (UK)

R. P. T. Tomkins (USA)

S. Krause (USA)

B. A. Wolf (FRG)

K. N. Marsh (USA)

Managing Editor

P. D. GUJRAL

IUPAC Secretariat, Oxford, UK

INTERNATIONAL UNION OF PURE AND APPLIED CHEMISTRY

IUPAC Secretariat: Bank Court Chambers, 2-3 Pound Way,
Templars Square, Cowley, Oxford OX4 3YF, UK

NOTICE TO READERS

If your library is not already a standing-order customer or subscriber to the Solubility Data Series, may we recommend that you place a standing order or subscription order to receive immediately upon publication all new volumes published in this valuable series. Should you find that these volumes no longer serve your needs, your order can be cancelled at any time without notice.

A complete list of volumes published in the Solubility Data Series will be found on p. 250.

SOLUBILITY DATA SERIES

Volume 49

ESTERS WITH WATER PART II: ESTERS 7-C TO 32-C

Volume Editors

FORREST GETZEN
North Carolina State University
Raleigh, North Carolina, USA

GLENN HEFTER
Murdoch University
Murdoch, Australia

ANDRZEJ MACZYNSKI
Institute of Physical Chemistry
Warsaw, Poland

Contributors (*Evaluators)

GLENN HEFTER*
Murdoch University
Murdoch, Australia

ZOFIA MACZYNSKA
Institute of Physical Chemistry
Warsaw, Poland

ADAM SKRZECZ
Institute of Physical Chemistry
Warsaw, Poland



PERGAMON PRESS

OXFORD · NEW YORK · SEOUL · TOKYO

U.K. Pergamon Press Ltd., Headington Hill Hall,
Oxford OX3 0BW, England

U.S.A. Pergamon Press Inc., 660 White Plains Road,
Tarrytown, New York 10591-5153, USA

KOREA Pergamon Press Korea, KPO Box 315, Seoul 110-603,
Korea

JAPAN Pergamon Press Japan, Tsunashima Building Annex,
3-20-12 Yushima, Bunkyo-ku, Tokyo 113, Japan

Copyright © 1992 International Union of Pure and
Applied Chemistry

All Rights Reserved. No part of this publication may be reproduced, stored in a retrieval system or transmitted in any form or by any means: electronic, electrostatic, magnetic tape, mechanical, photocopying, recording or otherwise, without permission in writing from the copyright holders.

First edition 1992

Library of Congress Cataloging in Publication Data

The Library of Congress has catalogued this serial title as follows:

Solubility data series.—Vol. 1—Oxford; New York: Pergamon, c 1979—v.; 28 cm.

Separately catalogued and classified in LC before no. 18.

ISSN 0191-5622 = Solubility data series.

1. Solubility—Tables—Collected works.

QD543.S6629 541.3'42'05-dc19 85-641351

AACR 2 MARC-S

British Library Cataloguing in Publication Data

A catalogue record for this book is available from the British Library

ISBN 0 08 041908 9

CONTENTS

Foreword		viii
Preface		x
Liquid-Liquid Solubility: Evaluation Philosophy and Methodology		xi
1	7-carbon Esters	1
1.1	2-Propenoic acid, 2-methyl-, 2-cyanoethyl ester	1
1.2	Formic acid cyclohexyl ester	2
1.3	2-Propenoic acid butyl ester	3
1.4	Propanedioic acid diethyl ester	4
1.5	Pentanedioic acid dimethyl ester	5
1.6	Acetic acid 1-methyl-1-butyl ester	6
1.7	Acetic acid 3-methyl-1-butyl ester	9
1.8	Acetic acid pentyl ester	25
1.9	Butanoic acid 1-methylethyl ester	41
1.10	Butanoic acid, 3-methyl-, ethyl ester	47
1.11	Butanoic acid propyl ester	53
1.12	Formic acid hexyl ester	57
1.13	Hexanoic acid methyl ester	58
1.14	Pentanoic acid ethyl ester	59
1.15	Propanoic acid butyl ester	63
1.16	Propanoic acid, 2,2-dimethyl-, ethyl ester	71
1.17	Acetic acid 3-methoxy-1-butyl ester	72
1.18	Propanoic acid, 2-hydroxy-, 1-butyl ester	73
2	8-carbon Esters	74
2.1	Acetic acid phenyl ester	74
2.2	Benzoic acid methyl ester	75
2.3	Benzoic acid, 2-hydroxy-, methyl ester	81
2.4	Acetic acid cyclohexyl ester	85
2.5	Butanedioic acid diethyl ester	91
2.6	2,3-Butanediol diacetate	95
2.7	Hexanedioic acid dimethyl ester	96
2.8	Acetic acid 1,3-dimethylbutyl ester	97
2.9	Acetic acid 2-ethyl-1-butyl ester	98
2.10	Acetic acid hexyl ester	99
2.11	Acetic acid 1-methylpentyl ester	105
2.12	Butanoic acid butyl ester	108
2.13	Formic acid heptyl ester	115
2.14	Heptanoic acid methyl ester	116
2.15	Hexanoic acid ethyl ester	117
2.16	Propanoic acid, 2-methyl-, butyl ester	119
2.17	Propanoic acid 3-methyl-1-butyl ester	120
2.18	Propanoic acid, 2-methyl-, 2-methylpropyl ester	121
2.19	Propanoic acid pentyl ester	122
2.20	Acetic acid 2-butoxyethyl ester	123
3	9-carbon Esters	124
8.1	Acetic acid phenylmethyl ester	124
8.2	Benzoic acid ethyl ester	128
8.3	Benzoic acid, 2-hydroxy-, ethyl ester	133
8.4	1,2,3-Propanetriol triacetate	134
8.5	Propanoic acid cyclohexyl ester	138
8.6	Pentanedioic acid diethyl ester	139
8.7	Acetic acid heptyl ester	140
8.8	Butanoic acid 3-methyl-1-butyl ester	141
8.9	Butanoic acid pentyl ester	142
8.10	Formic acid octyl ester	146
8.11	Heptanoic acid ethyl ester	147
8.12	Pentanoic acid 1-butyl ester	148
8.13	Propanoic acid hexyl ester	149
8.14	Dibutylphosphinic acid methyl ester	150
8.15	Methylphosphonic acid dibutyl ester	152

4	10-carbon Esters	153
4.1	1,2-Benzenedicarboxylic acid dimethyl ester	153
4.2	Hexanedioic acid diethyl ester	156
4.3	Acetic acid 2-ethylhexyl ester	161
4.4	Acetic acid octyl ester	162
4.5	Nonanoic acid methyl ester	163
4.6	Octanoic acid ethyl ester	164
4.7	Propanoic acid, 2-methyl-, hexyl ester	168
4.8	Acetic acid 2-(2-butoxyethoxy) ethyl ester	169
4.9	Dibutylphosphinic acid ethyl ester	173
4.10	Dipropylphosphinic acid butyl ester	175
4.11	Ethylphosphonic acid dibutyl ester	177
5	11-carbon Esters	178
5.1	Heptanedioic acid diethyl ester	178
5.2	Acetic acid isononyl ester	179
5.3	Decanoic acid methyl ester	180
5.4	Nonanoic acid ethyl ester	181
5.5	Dibutylphosphinic acid propyl ester	182
5.6	Propylphosphonic acid dibutyl ester	184
6	12-carbon Esters	185
6.1	1,2-Benzenedicarboxylic acid diethyl ester	185
6.2	Octanedioic acid diethyl ester	187
6.3	Decanoic acid ethyl ester	188
6.4	Dibutylphosphinic acid butyl ester	189
6.5	Butylphosphonic acid dibutyl ester	192
6.6	Phosphoric acid tributyl ester	194
6.7	Phosphoric acid tris(2-methylpropyl) ester	207
7	13-carbon Esters	208
7.1	Benzoic acid, 2-hydroxy-, phenyl ester	208
7.2	Phosphoric acid methyl diphenyl ester	211
7.3	Nonanedioic acid diethyl ester	212
8	14-carbon Esters	213
8.1	1,2-Benzenedicarboxylic acid di-2-propenyl ester	213
8.2	1,2-Benzenedicarboxylic acid bis(1-methylethyl) ester	214
8.3	1,2-Benzenedicarboxylic acid dipropyl ester	215
8.4	Decanedioic acid diethyl ester	216
9	15-carbon Esters	217
9.1	Phosphoric acid dibutyl methylphenyl ester	217
10	16-carbon Esters	218
10.1	1,2-Benzenedicarboxylic acid bis(2-methylpropyl) ester	218
10.2	1,2-Benzenedicarboxylic acid dibutyl ester	219

11	18-carbon Esters	224
11.1	Phosphoric acid butyl bis(methylphenyl) ester	224
11.2	1,2-Benzenedicarboxylic acid dipentyl ester	225
11.3	Phosphoric acid tris(butoxyethyl) ester	226
12	19-carbon Esters	227
12.1	1,2-Benzenedicarboxylic acid butyl phenylmethyl ester	227
13	20-carbon Esters	231
13.1	1,2-Benzenedicarboxylic acid butyl octyl ester	231
14	22-carbon Esters	232
14.1	L-Ascorbic acid 6-hexadecanoate	232
15	24-carbon Esters	233
15.1	1,2-benzenedicarboxylic acid bis(2-ethylhexyl) ester	233
15.2	Phosphoric acid tris(2-ethylhexyl) ester	234
16	26-carbon Esters	235
16.1	1,2-Benzenedicarboxylic acid dinonyl ester	235
17	27-carbon Esters	236
17.1	Benzoic acid, 2-hydroxy-, cinchonan-9-ol, 6'-methoxy, (8 α ,9R)	236
18	32-carbon Esters	237
18.1	1,2-benzenedicarboxylic acid didodecyl ester	237
	System Index	239
	Registry Number Index	244
	Author Index	247
	Solubility Data Series: Published and Forthcoming Volumes	250

FOREWORD

*If the knowledge is
undigested or simply wrong,
more is not better.*

The Solubility Data Series is a project of Commission V.8 (Solubility Data) of the International Union of Pure and Applied Chemistry (IUPAC). The project had its origins in 1973, when the Analytical Chemistry Division of IUPAC set up a Subcommittee on Solubility Data under the chairmanship of the late Prof. A. S. Kertes. When publication of the *Solubility Data Series* began in 1979, the Committee became a full commission of IUPAC, again under the chairmanship of Prof. Kertes, who also became Editor-in-Chief of the Series. The Series has as its goal the preparation of a comprehensive and critical compilation of data on solubilities in all physical systems, including gases, liquids and solids.

The motivation for the Series arose from the realization that, while solubility data are of importance in a wide range of fields in science and technology, the existing data had not been summarized in a form that was at the same time comprehensive and complete. Existing compilations of solubility data indeed existed, but they contained many errors, were in general uncritical, and were seriously out-of-date.

It was also realized that a new series of compilations of data gave educational opportunities, in that careful compilations of existing data could be used to demonstrate what constitutes data of high and lasting quality. As well, if the data were summarized in a sufficiently complete form, any individual could prepare his or her own evaluation, independently of the published evaluation. Thus, a special format was established for each volume, consisting of individual data sheets for each separate publication, and critical evaluations for each separate system, provided sufficient data from different sources were available for comparison. The compilations and, especially, the evaluation were to be prepared by active scientists who were either involved in producing new data, or were interested in using data of high quality. With minor modifications in format, this strategy has continued throughout the Series.

In the standard arrangement of each volume, the Critical Evaluation gives the following information:

(i) A text which discusses the numerical solubility information which has been abstracted from the primary sources in the form of compilation sheets. The text concerns primarily the quality of the data, after consideration of the purity of the materials and their characterization, the experimental method used, the uncertainties in the experimental values, the reproducibility, the agreement with accepted test values, and finally, the fitting of the data to suitable functions, along with statistical tests of the fitted data.

(ii) A set of recommended data, whenever possible, including weighted averages and estimated standard deviations. If applicable, one or more smoothing equations which have been computed or verified by the evaluator are also given.

(iii) A graphical plot of the recommended data, in the form of phase diagrams where appropriate.

The Compilation part consists of data sheets which summarize the experimental data from the primary literature. Here much effort is put into obtaining complete coverage; many good data have appeared in publications from the late nineteenth and early twentieth centuries, or in obscure journals. Data of demonstrably low precision are not compiled, but are mentioned in the Critical Evaluation. Similarly, graphical data, given the uncertainty of accurate conversion to numerical values, are compiled only where no better data are available. The documentation of data of low precision can serve to alert researchers to areas where more work is needed.

A typical data sheet contains the following information:

- (i) list of components: names, formulas, Chemical Abstracts Registry Numbers;
- (ii) primary source of the data;
- (iii) experimental variables;
- (iv) compiler's name;
- (v) experimental values as they appear in the primary source, in modern units with explanations if appropriate;
- (vi) experimental methods used;
- (vii) apparatus and procedure used;
- (viii) source and purity of materials used;
- (ix) estimated error, either from the primary source or estimated by the compiler;
- (x) references relevant to the generation of the data cited in the primary source.

Each volume also contains a general introduction to the particular type of system, such as solubility of gases, of solids in liquids, etc., which contains a discussion of the nomenclature used, the principle of accurate determination of solubilities, and related thermodynamic principles. This general introduction is followed by a specific introduction to the subject matter of the volume itself.

The Series embodies a new approach to the presentation of numerical data, and the details continue to be influenced strongly by the perceived needs of prospective users. The approach used will, it is hoped, encourage attention to the quality of new published work, as authors become more aware that their work will attain permanence only if it meets the standards set out in these volumes. If the Series succeeds in this respect, even partially, the Solubility Data Commission will have justified the labour expended by many scientists throughout the world in its production.

January, 1989

J. W. Lorimer,
London, Canada

PREFACE

This volume, one of a pair in the Solubility Data Series devoted to esters in water, includes solubility data for binary systems containing an ester and water published up through the end of 1988. The first volume contains the esters having two through six carbon atoms with water and the second volume contains esters having seven or more carbon atoms with water. With only rare exceptions, the compiled data and evaluations involve measurements made at constant pressure over rather restricted ranges of temperature. The limited ranges of pressure and temperature involved are to be expected in view of the liquid-liquid condition prescribed for the systems.

For the most part, the data presented show general patterns of behavior expected for the topic systems. While rather extensive data have been reported for many of the systems under consideration, no effort was made to attempt any detailed correlations of behavior or descriptions of trends beyond those presented by the evaluator in the evaluation sections.

The critical evaluations were all prepared by one author and an introductory section has been included to elaborate the philosophy and methodology followed in the evaluations. No attempts were made to extract thermodynamic parameters such as activity coefficients, enthalpies of solution, and the like from the reported data although much of the published work is sufficiently precise for such activities.

Every attempt was made to include all published data for the subject systems through the end of 1988. The editors would be grateful for any information concerning material which was overlooked in the preparation of this work.

The editors would like to express their appreciation to all those who have provided assistance in the volume preparation, especially to fellow members of the IUPAC Commission on Solubility Data and to those who have served as reviewers of this work.

Forrest W. Getzen
North Carolina State University
Raleigh, North Carolina, USA

Andrzej Maczynski
Instytut Chemii Fizycznej
Polskiej Akademii Nauk
Warszawa, POLAND

Glenn T. Hefter
Murdoch University
Murdoch, Western Australia, AUSTRALIA

November 1991.

LIQUID-LIQUID SOLUBILITY: EVALUATION PHILOSOPHY AND METHODOLOGY

G.T. HEFTER

INTRODUCTION

Because of the differing ways in which it is possible to critically evaluate published solubility data it is worthwhile to present a coherent statement on the philosophy and procedures which have been used in this volume. Although this is a personal statement concerning the present volume, many of the ideas have been developed by myself as an evaluator for previous liquid-liquid volumes in the Solubility Data Series [1-3] and are shared, to some degree, by other evaluators.

EVALUATION PHILOSOPHY

The critical evaluation of data is, at least in part, a subjective process. Some physical scientists are uncomfortable with this notion but the history of science shows us it is undeniable. Because some subjective judgements are inevitably involved, it is vital that the evaluator's procedure should be as visible as possible. In this way, if the user does not agree with the evaluator's judgement or arguments or if new data become available it is possible, with a minimum of effort, to derive a new set of preferred or "best" values. For this reason the Critical Evaluations in this and those previous volumes to which I have contributed [1-3] have been, in essence, written like research papers with enough information provided for users to repeat or change the evaluation should they so desire.

The starting point for each Critical Evaluation is that the data compilation is complete, i.e., the data sheets include all the available data. This is an important premise because the addition of extra data may change the evaluation considerably. This is especially true of liquid-liquid systems since very few have been adequately studied over a wide temperature range and disagreements often exist between independent investigations.

In general, a "democracy of observers" is followed. That is, the data of all investigators are given equal weight irrespective of their reputation or location. This is justifiable in scientific terms because experience shows that real "errors" in liquid-liquid solubility data are much greater than individual authors' reported precisions. Thus, the benzene-water system, which is relatively simple and probably the most widely studied of all liquid-liquid systems [2], has solubilities which are known to an accuracy of about $\pm 2\%$ (relative). The same estimated error is obtained regardless of whether a simple average of the available data is taken or whether a complex weighting procedure is used. This suggests that, at the present stage of development of liquid-liquid solubility determinations, equal weighting for all data is the most appropriate course of action. The exceptions to this rule are discussed below.

REJECTION OF DATA

The statement that all data are given equal weight must be qualified: some data are assigned zero weight, i.e., they are rejected. Data rejection is potentially dangerous, especially when few data are available. Nevertheless, since a critical evaluation is required, it has to be considered. The most important reasons for rejecting data are as follows.

1. The data disagree "significantly" from most other studies (especially if these are numerous). Significance is subjective but a criterion of

$$| s_{\text{obs}} - s_{\text{av}} | > 3\sigma_n$$

where s_{obs} is the reported solubility datum being considered for rejection

s_{av} is the arithmetic average (mean) of all other studies at the same temperature and pressure

σ_n is the standard deviation from the mean of these studies

has been generally, but not rigidly, applied [1-3].

It should be noted that this procedure can be invoked only once. That is, having rejected some points in a particular data set it is not statistically valid to recalculate a new value of σ_n and then reject further points. Strictly speaking this means that outlying data should be included in the original calculation of σ_n . However, the amount of available data does not always make this approach realistic and common sense must be applied.

2. Approximate data (one significant figure) are rejected when a reasonable amount of more precise data is available.
3. Data which are grossly inconsistent with chemical systematics or thermodynamic expectations are also rejected.
4. In keeping with the practice adopted in previous liquid-liquid solubility values, data originally presented in graphical form or in volume ratio units are also excluded. In the latter case, data are retained if appropriate density data are readily available to permit conversion to mass units. As density measurements are usually far more reliable than solubility data, the choice of density values is usually not critical.

It is appropriate to add here that data for a particular system coming from the same authors or the same laboratories have generally not been regarded as independent measurements and have therefore been subsumed into a single result (at each temperature).

THE EVALUATION FORMAT

The Critical Evaluations in this volume follow the format evolved in previous liquid-liquid solubility volumes [1-3]. In essence each Evaluation is divided into three parts. In the first part Table 1 lists all the available publications containing quantitative solubility data for the particular system. Broad experimental details such as temperature range, type of solubility measured ((1) in (2), (2) in (1) or mutual) and the method used are also included in Table 1. After mention of any other special features such as critical points, or any special types of investigation, further discussion is usually divided into two parts: the solubility of the organic in the aqueous phase, and that of water in the organic-rich phase. Since these two sections are handled in the same way they will be discussed together.

After listing rejected data, anomalies and caveats, all the available data (usually expressed as g/100g sln) are collected into Tables (usually Tables 2 & 3). To facilitate comparisons, data are interpolated (if necessary) to specific temperatures at 5K intervals near 298K, then 10 or 20K intervals. These temperatures are exact, i.e. 298.15, 303.15 ... K but for convenience are written 298, 303, ... K in the text. The nature of most liquid-liquid solubility data makes graphical interpolation the most appropriate approach [4]. Interpolated values are always designated with an asterisk (*). Very occasionally data are extrapolated but only over a short range (<5K) and only if the solubility curve is not changing sharply.

All the retained values at each temperature are then averaged and the mean and σ_n recorded in the Tables. Few of the systems in this volume have sufficient data to permit meaningful statistical analysis. The value of σ_n is therefore included only as a convenient measure of the spread of the experimental data.

"BEST" VALUES

The word "best", almost always in inverted commas (quotation marks), is used throughout the Critical Evaluations in this and previous liquid-liquid solubility volumes [1-3] in a specific way: to mean "best available estimate". It follows from what has been said above that "best" values are simply the arithmetic average (mean) of all the accepted data, including those obtained by interpolation and excluding rejected data. "Best" values are given in both g/100g sln and mole fraction units (x).

RECOMMENDED VALUES

Solubilities are recommended only if two or more independent studies (including rejected data) agree to within $\pm 5\%$ (relative), i.e.,

$$| 100\sigma_n/s_{av} | \leq 5$$

and there are no other obvious problems. Recommended data are designated by (R) immediately following the numerical value.

For sparingly soluble systems this criterion may be too harsh and

$$| 100\sigma_n/s_{av} | \leq 10 \text{ or } 20$$

may be more appropriate [3]. Very few high molecular weight esters have been sufficiently well characterized to enable even these less restrictive criteria to be applied.

Data which are not "Recommended" are regarded as "Tentative". The likely accuracy of these data varies from system to system and from temperature to temperature but may be inferred from

1. σ_n
2. $100\sigma_n/s_{av}$
3. n (the number of independent measurements).

Occasionally there are definite reasons to suspect the available data are not reliable. Such data are labelled "Doubtful" and the detailed reasons for this classification given. Considerable care should be exercised in the use of such data.

Note that the mole fraction "best" solubilities in each table should be regarded as having the same status ("Recommended", "Tentative" or "Doubtful") and the same (relative) percentage uncertainties (where given) as the corresponding mass % solubilities.

GRAPHICS

In systems for which a reasonable amount of data exists, selected results are plotted graphically over the studied temperature range. Appropriate comments on the plots are made in the figure caption or in the text.

FITTING EQUATIONS

The use of fitting equations to smooth solubility data as a function of temperature is controversial and a topic in its own right which will not be discussed here. As in previous liquid-liquid solubility volumes [1-3], fitting equations are not generally given in the printed version of this volume. There are several reasons for this.

1. There is no single equation form which can be used for all systems.
2. The data for some systems are too scattered or poorly defined for realistic curve fitting.
3. Without special care empirical fitting equations may give misleading results (spurious extrema, etc.) when interpolating and can almost never be reliably used for extrapolation.

For certain systems where reliable data appear evident, fitting equations have been provided. Such equations come directly from the original works or from the efforts of the compilers in the course of their examination of reported data and preparation of smoothed value tables.

Users who wish to derive their own fitting equations are advised to do so with caution. It is suggested that the "best" values be used for this purpose, but it should be noted that this may create anomalies because of the way in which the "best" values are derived (see above). Where fitting equations are given in this volume they should be used only over the stated range.

THERMODYNAMIC CONSISTENCY

Using standard relationships from thermodynamics it can be shown that:

$$\Delta_{\text{sln}}G = -RT \ln a_{\text{sat}}$$

$$\frac{\partial(\Delta_{\text{sln}}G/T)}{\partial T} = - \frac{\Delta_{\text{sln}}H}{RT^2}$$

$$\text{and } \frac{\partial^2(\Delta_{\text{sln}}G/T)}{\partial T^2} = \frac{\partial(\Delta_{\text{sln}}H)}{\partial T} = \Delta_{\text{sln}}C_p$$

where $\Delta_{\text{sln}}G$, $\Delta_{\text{sln}}H$ and $\Delta_{\text{sln}}C_p$ are respectively the Gibbs energy, enthalpy and heat capacity (at constant pressure) changes corresponding to the dissolution of the solute in the solvent and a_{sat} is the activity of the saturated solute in the solvent. If the solute is only sparingly soluble the thermodynamic parameters may be equated with their standard state values, and it should be noted that $\Delta_{\text{sln}}C_p$ may itself be temperature dependent.

Both $\Delta_{\text{sln}}H$ and $\Delta_{\text{sln}}C_p$, at least in principle, can be measured directly by separate calorimetric procedures [5,6] and therefore can serve as an independent check on the temperature dependence of published solubility data. This approach has been used with significant effect in evaluating hydrocarbon solubilities in water [2-3]. However, very few calorimetric data have been reported for ester-water systems and so this approach has not been utilized in the present volume.

REFERENCES

1. G.T. Hefter, in A.F.M. Barton, Ed., *Solubility Data Series*, Vol. 15, *Alcohols with Water* (Pergamon Press, Oxford, 1984).
2. G.T. Hefter, in D.G. Shaw, Ed., *Solubility Data Series*, Vol. 37, *Hydrocarbons with Water and Seawater, Part I: Hydrocarbons C5 to C7* (Pergamon Press, Oxford, 1989).
3. G.T. Hefter, in D.G. Shaw, Ed., *Solubility Data Series*, Vol. 38, *Hydrocarbons with Water and Seawater, Part II: Hydrocarbons C8 to C36* (Pergamon Press, Oxford, 1989).
4. G.T. Hefter, unpublished paper presented at 1st Intl. Symp. *Soly. Phenom.*, London, Canada, August, 1984.
5. S.O. Nilsson and I. Wadso, *J. Chem. Thermodyn.* **1984**, 16, 317.
6. P. Picker, P.A. Leduc, P.R. Philip and J.E. Desnoyers, *J. Chem. Thermodyn.* **7**, 3, 641.

COMPONENTS: (1) 2-Propenoic acid, 2-methyl-, 2-cyanoethyl ester (2-cyanoethyl methacrylate); $C_7H_9NO_2$; [4513-53-5] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Frolov, A.F.; Maiorova, N.M.; Erykov, V.G.; Korshukov, M.A. <i>Zh. Prikl. Khim.</i> <u>1977</u> , 50, 1318-21.
VARIABLES: $T/K = 293$	PREPARED BY: Z. Maczynska
EXPERIMENTAL VALUES: <p>The solubility of 2-methyl-2-propenoic acid 2-cyanoethyl ester in water at 20°C was reported to be 1.7 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 0.0022.</p> <p>The solubility of water in 2-methyl-2-propenoic acid 2-cyanoethyl ester at 20°C was reported to be 2.9 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.19.</p>	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The method was not specified.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified; b.p. 240.0°C, d_4^{20} 1.0400, n_D^{20} 1.4460. (2) Not specified.
ESTIMATED ERROR: Not specified.	
REFERENCES: 	

COMPONENTS: (1) Formic acid cyclohexyl ester (cyclohexyl formate); $C_7H_{12}O_2$; [4351-54-6] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u> , 31, 56-70.																																																						
VARIABLES: $T/K = 273 - 363$	PREPARED BY: Z. Maczynska																																																						
EXPERIMENTAL VALUES:																																																							
Mutual solubility of formic acid cyclohexyl ester and water																																																							
<table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th rowspan="2" style="text-align: left;">t/°C</th> <th colspan="2" style="text-align: center;">g(1)/100g sln</th> <th colspan="2" style="text-align: center;">x_1 (compiler)</th> </tr> <tr> <th style="text-align: center;">(2)-rich phase</th> <th style="text-align: center;">(1)-rich phase</th> <th style="text-align: center;">(2)-rich phase</th> <th style="text-align: center;">(1)-rich phase</th> </tr> </thead> <tbody> <tr> <td>0</td> <td style="text-align: center;">-</td> <td style="text-align: center;">98.90</td> <td style="text-align: center;">-</td> <td style="text-align: center;">0.9267</td> </tr> <tr> <td>9.3</td> <td style="text-align: center;">1.70</td> <td style="text-align: center;">98.58</td> <td style="text-align: center;">0.00242</td> <td style="text-align: center;">0.9070</td> </tr> <tr> <td>19.0</td> <td style="text-align: center;">0.65</td> <td style="text-align: center;">98.44</td> <td style="text-align: center;">0.00092</td> <td style="text-align: center;">0.8987</td> </tr> <tr> <td>29.6</td> <td style="text-align: center;">0.39</td> <td style="text-align: center;">98.63</td> <td style="text-align: center;">0.00055</td> <td style="text-align: center;">0.9101</td> </tr> <tr> <td>39.5</td> <td style="text-align: center;">-</td> <td style="text-align: center;">98.61</td> <td style="text-align: center;">-</td> <td style="text-align: center;">0.9088</td> </tr> <tr> <td>49.9</td> <td style="text-align: center;">-</td> <td style="text-align: center;">98.36</td> <td style="text-align: center;">-</td> <td style="text-align: center;">0.8939</td> </tr> <tr> <td>69.9</td> <td style="text-align: center;">-</td> <td style="text-align: center;">98.10</td> <td style="text-align: center;">-</td> <td style="text-align: center;">0.8789</td> </tr> <tr> <td>80.0</td> <td style="text-align: center;">-</td> <td style="text-align: center;">97.87</td> <td style="text-align: center;">-</td> <td style="text-align: center;">0.8659</td> </tr> <tr> <td>90.3</td> <td style="text-align: center;">-</td> <td style="text-align: center;">97.50</td> <td style="text-align: center;">-</td> <td style="text-align: center;">0.8457</td> </tr> </tbody> </table>		t/°C	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	-	98.90	-	0.9267	9.3	1.70	98.58	0.00242	0.9070	19.0	0.65	98.44	0.00092	0.8987	29.6	0.39	98.63	0.00055	0.9101	39.5	-	98.61	-	0.9088	49.9	-	98.36	-	0.8939	69.9	-	98.10	-	0.8789	80.0	-	97.87	-	0.8659	90.3	-	97.50	-	0.8457
t/°C	g(1)/100g sln		x_1 (compiler)																																																				
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																			
0	-	98.90	-	0.9267																																																			
9.3	1.70	98.58	0.00242	0.9070																																																			
19.0	0.65	98.44	0.00092	0.8987																																																			
29.6	0.39	98.63	0.00055	0.9101																																																			
39.5	-	98.61	-	0.9088																																																			
49.9	-	98.36	-	0.8939																																																			
69.9	-	98.10	-	0.8789																																																			
80.0	-	97.87	-	0.8659																																																			
90.3	-	97.50	-	0.8457																																																			
std. dev. 0.01 0.02																																																							
AUXILIARY INFORMATION																																																							
METHOD/APPARATUS/PROCEDURE: <p>The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.</p>	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial sample; purity 95%; used as received. (2) Not specified.																																																						
	ESTIMATED ERROR: Accuracy of method 0.1 wt% or less, for solubility, see above.																																																						
	REFERENCES:																																																						

COMPONENTS: (1) 2-Propenoic acid butyl ester (butyl acrylate); $C_7H_{12}O_2$; [141-32-2] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Logutov, V.I.; Danov, S.M.; Chubarov, G.A.; Tomashchuk, V.I. <i>Zh. Prikl. Khim.</i> <u>1983</u> , 56(1), 214-6.
VARIABLES: $T/K = 293$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: <p>The solubility of 2-Propenoic acid butyl ester in water at 20°C was reported to be 0.18 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 2.6×10^{-4}.</p> <p>The solubility of water in 2-Propenoic acid butyl ester at 20°C was reported to be 0.65 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.044.</p> <p>The densities of the water-rich phase and the ester rich phase were reported to be d_{20}^{20} 0.9981 and d_{20}^{20} 0.8987 respectively.</p>	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: <p>The analytical method was used. After mixing and separating, the two samples were analysed. The water in organic phase was determined by titration with Karl Fischer reagent. The 2-Propenoic acid butyl ester was measured by the gas-liquid chromatography method with an inert standard or by the bromate titration method. Both procedures gave similar results.</p>	SOURCE AND PURITY OF MATERIALS: (1) Source not specified; used as received; b.p. 147.4°C, d_4^{20} 0.9010, n_D^{20} 1.4200. (2) Twice distilled. ESTIMATED ERROR: Temp. $\pm 0.5^\circ C$. Soly. $\pm(2-3)\%$ (relative) (2) in organic phase, $\pm 3\%$ (relative) (1) in water phase. REFERENCES:

<p>COMPONENTS:</p> <p>(1) Propanedioic acid diethyl ester (diethyl malonate); $C_7H_{12}O_4$; [105-53-3]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Sobotka, H.; Kahn, J.</p> <p><i>J. Am. Chem. Soc.</i> <u>1931</u>, 53, 2935-8.</p>
<p>VARIABLES:</p> <p>$T/K = 293$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of propanedioic acid diethyl ester in water at 20°C was reported to be 2.08 g(1)/100mL(2). The corresponding mass per cent and mole fraction, x_1, values calculated by the compiler are 2.04 g(1)/100g sln and 0.0023.</p> <p>Density of water $d_4^{20} = 0.9982$ (ref 1) was used in the calculation.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The titration method was used. The ester was added dropwise from a micro-burette with a capillary tip to 100, 250, or 500 mL of water in a narrow-mouthed stock bottle with a well-ground glass stopper. The bottle was shaken after each addition of ester. 1-5 mg of Sudan IV dye was put into the water to improve the end-point of the titration. At saturation, one additional drop of ester was sufficient to convert the floating rough indicator particles into dark transparent droplets.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified (Eastman Kodak Laboratories or synthesized); twice-distilled under reduced pressure; d_4^{20} 1.0550, n_D^{20} 1.4144.</p> <p>(2) Distilled.</p> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <p>REFERENCES:</p> <p>1. <i>Selected Values of Properties of Hydrocarbons and Related Compounds</i>, API Research Project 44, Thermodynamics Research Center, Texas A and M University, Texas, <u>1973</u>.</p>

COMPONENTS: (1) Pentanedioic acid dimethyl ester (<i>dimethyl glutarate</i>); $C_7H_{12}O_4$; [1119-40-0] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u> , 31, 56-70.																																																											
VARIABLES: T/K = 273 - 364	PREPARED BY: Z. Maczynska																																																											
EXPERIMENTAL VALUES: Mutual solubility of pentanedioic acid dimethyl ester and water <table border="1" data-bbox="154 521 1141 1032"> <thead> <tr> <th rowspan="2">t/°C</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>7.1</td><td>97.70</td><td>0.0085</td><td>0.8269</td></tr> <tr><td>9.4</td><td>6.3</td><td>97.37</td><td>0.0075</td><td>0.8063</td></tr> <tr><td>20.0</td><td>6.1</td><td>97.72</td><td>0.0072</td><td>0.8282</td></tr> <tr><td>29.6</td><td>5.9</td><td>96.44</td><td>0.0070</td><td>0.7529</td></tr> <tr><td>39.5</td><td>5.8</td><td>95.88</td><td>0.0069</td><td>0.7236</td></tr> <tr><td>49.8</td><td>5.8</td><td>95.36</td><td>0.0069</td><td>0.6980</td></tr> <tr><td>60.3</td><td>6.1</td><td>94.69</td><td>0.0072</td><td>0.6673</td></tr> <tr><td>70.2</td><td>6.4</td><td>92.89</td><td>0.0076</td><td>0.5950</td></tr> <tr><td>80.0</td><td>6.6</td><td>92.86</td><td>0.0079</td><td>0.5939</td></tr> <tr><td>90.5</td><td>8.1</td><td>91.77</td><td>0.0098</td><td>0.5564</td></tr> </tbody> </table> <p>std. dev. 0.10 . 0.03</p>		t/°C	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	7.1	97.70	0.0085	0.8269	9.4	6.3	97.37	0.0075	0.8063	20.0	6.1	97.72	0.0072	0.8282	29.6	5.9	96.44	0.0070	0.7529	39.5	5.8	95.88	0.0069	0.7236	49.8	5.8	95.36	0.0069	0.6980	60.3	6.1	94.69	0.0072	0.6673	70.2	6.4	92.89	0.0076	0.5950	80.0	6.6	92.86	0.0079	0.5939	90.5	8.1	91.77	0.0098	0.5564
t/°C	g(1)/100g sln		x_1 (compiler)																																																									
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																								
0	7.1	97.70	0.0085	0.8269																																																								
9.4	6.3	97.37	0.0075	0.8063																																																								
20.0	6.1	97.72	0.0072	0.8282																																																								
29.6	5.9	96.44	0.0070	0.7529																																																								
39.5	5.8	95.88	0.0069	0.7236																																																								
49.8	5.8	95.36	0.0069	0.6980																																																								
60.3	6.1	94.69	0.0072	0.6673																																																								
70.2	6.4	92.89	0.0076	0.5950																																																								
80.0	6.6	92.86	0.0079	0.5939																																																								
90.5	8.1	91.77	0.0098	0.5564																																																								
AUXILIARY INFORMATION																																																												
METHOD/APPARATUS/PROCEDURE: The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial sample; purity 98%; used as received. (2) Not specified. ESTIMATED ERROR: Accuracy of method 0.1 wt% or less, for solubility, see above. REFERENCES:																																																											

<p>COMPONENTS:</p> <p>(1) Acetic acid 1-methyl-1-butyl ester (1-methylbutyl acetate); C₇H₁₄O₂; [626-38-0]</p> <p>(2) Water; H₂O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia January, 1989</p>												
<p>CRITICAL EVALUATION:</p> <p>Quantitative solubility data for the acetic acid 1-methyl-1-butyl ester (1) - water (2) system have been reported in the publications listed in Table 1.</p> <p style="text-align: center;"><u>TABLE 1: Quantitative Solubility Studies of the Acetic acid 1-methyl-1-butyl ester (1) - Water (2) System</u></p> <table border="1" data-bbox="222 660 1258 829"> <thead> <tr> <th>Reference</th> <th>T/K</th> <th>Solubility</th> <th>Method</th> </tr> </thead> <tbody> <tr> <td>Park and Hopkins (ref 1)</td> <td>298</td> <td>(1) in (2)</td> <td>unspecified</td> </tr> <tr> <td>Doolittle (ref 2)</td> <td>293</td> <td>mutual</td> <td>unspecified</td> </tr> </tbody> </table> <p>The original data in these publications are compiled in the Data Sheets immediately following this Critical Evaluation.</p> <p>As can be seen from Table 1 no data have been reported under comparable conditions and so no Critical Evaluation is possible. Furthermore, the datum of Park and Hopkins (ref 1) is reported only in v/v units and would normally be excluded from consideration. However, assuming a density of 0.86 g/mL for (1) (ref 2) and a density of 1.00 g/mL for the saturated solution of (1) in (2), the solubility reported by Park and Hopkins (ref 1) at 298 K is equivalent to 0.23 g(1)/100 g sln which is in good agreement with Doolittle's value of 0.19 g(1)/100 g sln at 293 K (ref 1). Doolittle (ref 1) also reports a value of 0.88 g(2)/100 g sln for the solubility of water in the ester. The interested user is referred to the relevant Data Sheets for additional information.</p> <p>REFERENCES</p> <ol style="list-style-type: none"> Park, J. G.; Hopkins, M. B. <i>Ind. Eng. Chem.</i> <u>1930</u>, 22, 826-30. Doolittle, A. K. <i>Ind. Eng. Chem.</i> <u>1935</u>, 27, 1169-79. 		Reference	T/K	Solubility	Method	Park and Hopkins (ref 1)	298	(1) in (2)	unspecified	Doolittle (ref 2)	293	mutual	unspecified
Reference	T/K	Solubility	Method										
Park and Hopkins (ref 1)	298	(1) in (2)	unspecified										
Doolittle (ref 2)	293	mutual	unspecified										

<p>COMPONENTS:</p> <p>(1) Acetic acid 1-methyl-1-butyl ester (1-methylbutyl acetate); $C_7H_{14}O_2$; [626-38-0]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Park, J.G.; Hopkins, M.B. <i>Ind. Eng. Chem.</i> <u>1930</u>, 22, 826-30.</p>
<p>VARIABLES:</p> <p>T/K = 298</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of acetic acid 1-methyl-1-butyl ester in water at 25°C was reported to be 0.2 mL(1)/100mL(2).</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The method was not specified.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, commercial samples; used as received; 85-88% of ester, b.p. range 128-134°C.</p> <p>(2) Not specified.</p>
	<p>ESTIMATED ERROR:</p> <p>Not specified.</p>
	<p>REFERENCES:</p>

<p>COMPONENTS:</p> <p>(1) Acetic acid 1-methyl-1-butyl ester (<i>1-methylbutyl acetate</i>); $C_7H_{14}O_2$; [626-38-0]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Doolittle, A.K. <i>Ind. End. Chem.</i> <u>1935</u>, 27, 1169-79.</p>
<p>VARIABLES:</p> <p>$T/K = 293$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of acetic acid 1-methyl-1-butyl ester in water at 20°C was reported to be 0.19 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 2.6×10^{-4}.</p> <p>The solubility of water in acetic acid 1-methyl-1-butyl ester at 20°C was reported to be 0.88 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.060.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The method was not specified.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, commercial product; purity 92%, b.p. range 121-144°C, d_4^{20} 0.862.</p> <p>(2) Not specified.</p> <hr/> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <hr/> <p>REFERENCES:</p>

COMPONENTS:

- (1) Acetic acid 3-methyl-1-butyl ester (*isopentyl acetate*);
C₇H₁₄O₂; [123-92-2]
(2) Water; H₂O; [7732-18-5]

EVALUATOR:

G.T. Hefter, School of Mathematical and Physical Sciences,
Murdoch University, Perth, W.A.,
Australia
January, 1989

CRITICAL EVALUATION:

Quantitative solubility data for the acetic acid 3-methyl-1-butyl ester (1) - water (2) system have been reported in the publications listed in Table 1.

TABLE 1: Quantitative Solubility Studies of the Acetic acid 3-methyl-1-butyl ester (1) - Water (2) System

Reference	T/K	Solubility	Method
Bancroft (ref 1)	293	mutual	titration
Othmer et al. (ref 2)	297	mutual	unspecified
Narasimhan et al. (ref 3)	303	mutual	titration, refractometric
Iguchi and Fuse (ref 4)	298	mutual	titration
Ramanarao et al. (ref 5)	303	mutual	turbidometric
Andreeva et al. (ref 6)	367	mutual	titration
Rao et al. (ref 7)	303	mutual	titration
Skrzecz (ref 8)	288-359	mutual	synthetic, Karl Fischer
Stephenson and Stuart (ref 9)	273-364	mutual	GLC

The original data in these publications are compiled in the Data Sheets immediately following this Critical Evaluation. For convenience, further discussion of this system will be divided into two parts.

1. SOLUBILITY OF ACETIC ACID 3-METHYL-1-BUTYL ESTER (1) IN WATER (2)

All the available data for the solubility of acetic acid 3-methyl-1-butyl ester (1) in water (2) are summarized in Table 2 with the following exceptions.

The value of Bancroft (ref 1) reported in v/v units is excluded from consideration. The data reported by Iguchi and Fuse (ref 4), Ramanarao et al. (ref 5) and Andreeva et al., (ref 6) are substantially higher ($> 3\sigma_n$) than other values and have been rejected. The datum of Othmer et al. (ref 2), which is considerably lower than other studies is also rejected.

(continued next page)

<p>COMPONENTS:</p> <p>(1) Acetic acid 3-methyl-1-butyl ester (<i>isopentyl acetate</i>); C₇H₁₄O₂; [123-92-2]</p> <p>(2) Water; H₂O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia January, 1989</p>
-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

The remaining data, largely due to Skrzecz (ref 8) and Stephenson and Stuart (ref 9) are in excellent agreement, enabling the average "Best" values to be Recommended over a wide temperature range. Selected data are plotted in Figure 1.

TABLE 2: Recommended (R) and Tentative Solubilities of Acetic acid 3-methyl-1-butyl ester (1) in Water (2)

T/K	Solubilities		
	Reported values	"Best" values ($\pm\sigma_n$) ^a	
	g(1)/100g sln	g(1)/100g sln	10 ⁴ x ₁
273	0.34 (ref 9)	0.34	4.7
283	0.260* (ref 9)	0.26	3.6
293	0.218* (ref 8), 0.220* (ref 9)	0.219 ± 0.001 (R)	3.03
298	0.207* (ref 8), 0.208* (ref 9)	0.208 ± 0.001 (R)	2.88
303	0.22 (ref 3), 0.19 (ref 7), 0.198* (ref 8), 0.196* (ref 9)	0.201 ± 0.011 (R)	2.78
313	0.185* (ref 8), 0.182* (ref 9)	0.184 ± 0.002 (R)	2.55
323	0.178* (ref 8), 0.174* (ref 9)	0.176 ± 0.002 (R)	2.44
333	0.178* (ref 8), 0.174* (ref 9)	0.176 ± 0.002 (R)	2.44
343	0.185* (ref 8), 0.179* (ref 9)	0.182 ± 0.003 (R)	2.52
353	0.199* (ref 8), 0.190* (ref 9)	0.195 ± 0.005 (R)	2.70
363	0.219* (ref 8), 0.203 (ref 8)	0.211 ± 0.008 (R)	2.92

^a Obtained by averaging where appropriate; σ_n has no statistical significance. Mole fraction solubilities (x₁) have the same status and (relative) percentage uncertainties as the mass % solubilities.

(continued next page)

COMPONENTS:

- (1) Acetic acid 3-methyl-1-butyl ester (*isopentyl acetate*);
 $C_7H_{14}O_2$; [123-92-2]
 (2) Water; H_2O ; [7732-18-5]

EVALUATOR:

G.T. Hefter, School of Mathematical
 and Physical Sciences,
 Murdoch University, Perth, W.A.,
 Australia
 January, 1989

CRITICAL EVALUATION: (continued)




FIGURE 1. Selected data for the solubility of acetic acid 3-methyl-1-butyl ester (1) in water (2): ref 8 (●); ref 9 (○). Solid line is a least square polynomial fitted to the "Best" values from Table 2.

2. SOLUBILITY OF WATER (2) IN ACETIC ACID 3-METHYL-1-BUTYL ESTER (1)

All the available data for the solubility of water (2) in acetic acid 3-methyl-1-butyl ester (1) are summarized in Table 3 with the following exceptions.

The datum of Bancroft (ref 1) reported in v/v units has been excluded from consideration. The data reported by Othmer et al. (ref 2) and Andreeva et al. (ref 6) are significantly higher than other studies and have been rejected. The data of Iguchi and Fuse (ref 4) and Rao et al. (ref 7) are significantly lower than other studies and are also rejected.

The remaining data, mainly due to Skrzecz (ref 8) and Stephenson and Stuart (ref 9), are in general in good agreement although they differ significantly at $T > 343$ K. Selected data are plotted in Figure 2.

(continued next page)

<p>COMPONENTS:</p> <p>(1) Acetic acid 3-methyl-1-butyl ester (<i>isopentyl acetate</i>); C₇H₁₄O₂; [123-92-2]</p> <p>(2) Water; H₂O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia January, 1989</p>
-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

TABLE 3: Recommended (R) and Tentative Solubilities of Water (2) in Acetic acid 3-methyl-1-butyl ester (1)

T/K	Solubilities		
	Reported values	"Best" values ($\pm\sigma_n$) ^a	
	g(2)/100g sln	g(2)/100g sln	10 ² x ₂
273	0.56 (ref 9)	0.56	3.9
283	0.67* (ref 9)	0.67	4.7
293	0.67* (ref 8), 0.78* (ref 9)	0.73 ± 0.06	5.1
298	0.82* (ref 8), 0.82* (ref 9)	0.82 (R)	5.6
303	0.58 ^b (ref 3), 0.545 ^b (ref 5) 0.83* (ref 8), 0.87* (ref 9)	0.85 ± 0.02 (R)	5.8
313	0.85* (ref 8), 0.96* (ref 9)	0.90 ± 0.06 (R)	6.2
323	0.96* (ref 8), 1.05* (ref 9)	1.00 ± 0.05 (R)	6.8
333	1.11* (ref 8), 1.13* (ref 9)	1.12 ± 0.01 (R)	7.6
343	1.32* (ref 8), 1.21* (ref 9)	1.26 ± 0.06 (R)	8.5
353	1.62* (ref 8), 1.28* (ref 9)	1.5 ± 0.2	10
363	2.03* (ref 8), 1.33* (ref 9)	1.7 ± 0.4	17

^a Obtained by averaging where appropriate; σ_n has no statistical significance. Mole fraction solubilities (x₂) have the same status and (relative) percentage uncertainties as the mass % solubilities.

^b Not included in calculation of "Best" value.

(continued next page)

COMPONENTS:

- (1) Acetic acid 3-methyl-1-butyl ester (isopentyl acetate);
 $C_7H_{14}O_2$; [123-92-2]
- (2) Water; H_2O ; [7732-18-5]

EVALUATOR:

G.T. Hefter, School of Mathematical
 and Physical Sciences,
 Murdoch University, Perth, W.A.,
 Australia
 January, 1989

CRITICAL EVALUATION: (continued)




FIGURE 2. Selected data for the solubility of water (2) in acetic acid 3-methyl-1-butyl ester (1): ref 8 (●); ref 9 (○). Solid line is a least square polynomial fitted to the "Best" values from Table 3.

REFERENCES

1. Bancroft, W. D. *Phys. Rev.* 1895, 3, 114-36.
2. Othmer, D. F.; White, R. E.; Trueger, E. *Ind. Eng. Chem.* 1941, 33, 1240-8; *ibid.* 1513.
3. Narasimhan, K. S.; Reddy, C. C.; Chari, K. S. *J. Chem. Eng. Data* 1962, 7, 457-60.
4. Iguchi, A.; Fuse, K. *Kagaku Kogaku* 1971, 35, 1035-7.
5. Ramanarao, M. V.; Husain, A.; Chari, K. S. *Indian J. Technol.* 1964, 2, 252-4.
6. Andreeva, N. G.; Komarova, L. F.; Garber, Yu. N. *Zh. Prikl. Khim.* 1978, 51, 2031-6.
7. Rao, D. S.; Rao, K. V.; Prasad, A. R.; Chiranjivi, C. *J. Chem. Eng. Data* 1979, 24, 241-4.

(continued next page)

<p>COMPONENTS:</p> <p>(1) Acetic acid 3-methyl-1-butyl ester (isopentyl acetate); C₇H₁₄O₂; [123-92-2]</p> <p>(2) Water; H₂O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia January, 1989</p>
<p>CRITICAL EVALUATION: (continued)</p> <p>8. Skrzecz, A. <i>Pol. J. Chem.</i> <u>1981</u>, 55, 1177-80; see also Skrzecz, A. <i>Thesis, I. Ch. F. PAN, Warszawa, 1979.</i></p> <p>9. Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u>, 31, 56-70.</p> <p>ACKNOWLEDGEMENT</p> <p>The Evaluator thanks Dr. Brian Clare for the graphics.</p>	

COMPONENTS: (1) Acetic acid 3-methyl-1-butyl ester (<i>isopentyl acetate</i>); $C_7H_{14}O_2$; [123-92-2] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Bancroft, W.D. <i>Phys. Rev.</i> <u>1895</u> , 3, 114-36.
VARIABLES: $T/K = 293$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: The solubility of acetic acid 3-methyl-1-butyl ester in water at 20°C was reported to be 0.02 mL(1)/10mL(2). The solubility of water in acetic acid 3-methyl-1-butyl ester at 20°C was reported to be 0.12 mL(2)/10mL(1).	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The titration method was used. 10 mL of solvent in a test tube was titrated with the second component until the solution became cloudy.	SOURCE AND PURITY OF MATERIALS: (1) Laboratory source; dried over $CaCl_2$, distilled. (2) Not specified. ESTIMATED ERROR: Soly. ± 0.01 mL. REFERENCES:

COMPONENTS: (1) Acetic acid 3-methyl-1-butyl ester (<i>isopentyl acetate</i>); $C_7H_{14}O_2$; [123-92-2] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Othmer, D.F.; White, R.E.; Trueger, E. <i>Ind. Eng. Chem.</i> <u>1941</u> , 33, 1240-8, 1513.
VARIABLES: $T/K = 297$	PREPARED BY: Z. Maczynska
EXPERIMENTAL VALUES: <p>The solubility of acetic acid 3-methyl-1-butyl ester in water at 23.5°C was reported to be 0.16 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 2.2×10^{-4}.</p> <p>The solubility of water in acetic acid 3-methyl-1-butyl ester at 23.5°C was reported to be 2.8 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.172.</p>	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: Nothing was specified in the paper.	SOURCE AND PURITY OF MATERIALS: (1) Not specified. (2) Not specified. ESTIMATED ERROR: Temp. $\pm 0.5^\circ C$ (mean of reported range), Soly. about ± 0.04 g(1)/100g sln and ± 2.1 g(2)/100g sln (compiler). REFERENCES:

COMPONENTS: (1) Acetic acid 3-methyl-1-butyl ester (<i>isopentyl acetate</i>); $C_7H_{14}O_2$; [123-92-2] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Narasimhan, K.S.; Reddy, C.C.; Chari, K.S. <i>J. Chem. Eng. Data</i> <u>1962</u> , 7, 457-60.
VARIABLES: $T/K = 303$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: <p>The solubility of acetic acid 3-methyl-1-butyl ester in water at 30°C was reported to be 0.22 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 3.1×10^{-4}.</p> <p>The solubility of water in acetic acid 3-methyl-1-butyl ester at 30°C was reported to be 0.58 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.040.</p>	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: <p>The titration and the analytical methods described by Narasimhan, Reddy and Chari (ref 1) were used. In the titration method, a known weight of mixture was taken in 100mL stoppered conical flask and kept in a thermostat for 1/2 h. which was found to be sufficient for equilibrium. Then, the selected component was added dropwise until a permanent turbidity appeared in the solution.</p> <p>For the analytical procedure, the mixture was stirred for 1 h. in a thermostated equilibrium apparatus. After separation the refractive indexes of both layers were measured and the corresponding compositions were obtained graphically from a previously drawn plot relating the refractive index and composition. The data were reported together with the ternary system acetic acid 3-methyl-1-butyl ester-water-phenol.</p>	SOURCE AND PURITY OF MATERIALS: (1) Source not specified; twice fractionated; purity 100%, b.p. 140.1-142.1°C, d_4^{24} 0.8670, n_D^{20} 1.4005. (2) Distilled. ESTIMATED ERROR: Temp. $\pm 0.1^\circ C$ by (ref 1). Soly. about ± 0.02 g(1)/100g sln and ± 0.18 g(2)/100g sln (compiler). REFERENCES: 1. Narasimhan, K.S.; Reddy, C.C.; Chari, K.S. <i>J. Chem. Eng. Data</i> <u>1962</u> , 7, 340.

<p>COMPONENTS:</p> <p>(1) Acetic acid 3-methyl-1-butyl ester (<i>isopentyl acetate</i>); C₇H₁₄O₂; [123-92-2]</p> <p>(2) Water; H₂O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Iguchi, A.; Fuse, K. Kagaku Kogaku <u>1971</u>, 35(9), 1035-7.</p>
<p>VARIABLES:</p> <p>T/K = 298</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of acetic acid 3-methyl-1-butyl ester in water at 25°C was reported to be 0.43 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 6.0×10^{-4}.</p> <p>The solubility of water in acetic acid 3-methyl-1-butyl ester at 25°C was reported to be 0.32 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.0227.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The titration method was used. The samples were titrated with the second component to obtain turbidity. The method and data were reported together with the ternary system acetic acid 3-methyl-1-butyl ester-water-acetic acid. No further details were reported in the paper.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, commercial grade of high purity, used as received.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm 0.1^\circ\text{C}$. Soly. about ± 0.2 g(1)/100g sln and ± 0.4 g(2)/100g sln (compiler).</p> <p>REFERENCES:</p>

COMPONENTS: (1) Acetic acid 3-methyl-1-butyl ester (<i>isopentyl acetate</i>); $C_7H_{14}O_2$; [123-92-2] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Ramanarao, M.V.; Husain, A.; Chari, K.S. <i>Indian J. Technol.</i> <u>1964</u> , 2, 252-4.
VARIABLES: $T/K = 303$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: <p>The solubility of acetic acid 3-methyl-1-butyl ester in water at 30°C was reported to be 0.2500 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 3.467×10^{-4}.</p> <p>The solubility of water in acetic acid 3-methyl-1-butyl ester at 30°C was reported to be 0.5445 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.03806.</p>	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: <p>The turbidity end point method described by Narasimhan, Reddy and Chari (ref 1) was used. Only the method for the ternary system was described. A known weight of mixture was placed into a 100 mL stoppered conical flask and kept in a thermostat for 1/2 h. which was found to be sufficient for equilibrium. Then, the selected component was added dropwise until a permanent turbidity appeared in the solution.</p>	SOURCE AND PURITY OF MATERIALS: (1) Source not specified; twice distilled; b.p. 142.1°C. (2) Distilled. ESTIMATED ERROR: Temp. $\pm 0.1^\circ C$ (authors). Soly. about ± 0.05 g(1)/100g sln and ± 0.2 g(2)/100g sln (compiler). REFERENCES: 1. Narasimhan, K.S.; Reddy, C.C.; Chari, K.S. <i>J. Chem. Eng. Data</i> <u>1962</u> , 7, 340.

<p>COMPONENTS:</p> <p>(1) Acetic acid 3-methyl-1-butyl ester (<i>isopentyl acetate</i>); C₇H₁₄O₂; [123-92-2]</p> <p>(2) Water; H₂O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Andreeva, N.G.; Komarova, L.F.; Garber, Yu.N.</p> <p>Zh. Prikl. Khim. <u>1978</u>, 51, 2031-6.</p>
<p>VARIABLES:</p> <p>T/K = 367</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of acetic acid 3-methyl-1-butyl ester in water at 94.0°C^a was reported to be $x_1 = 0.011$. The corresponding mass per cent value calculated by the compiler is 7.4 g(1)/100g sln.</p> <p>The solubility of water in acetic acid 3-methyl-1-butyl ester at 94.0°C^a was reported to be $x_2 = 0.179$. The corresponding mass per cent value calculated by the compiler is 2.93 g(2)/100g sln.</p> <p>^a Boiling temperature at atmospheric pressure.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The titration method of Mozhukhin, Tyurikov and Mitropolskaya (ref 1) was used. The data were reported together with the ternary system acetic acid 3-methyl-1-butyl ester-water-acetic acid butyl ester (<i>isopentyl acetate-water-butyl acetate</i>). No further details were reported in the paper.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Not specified.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Soly. about ± 7.2 g(1)/100g sln and ± 0.8 g(2)/100g sln (compiler).</p> <p>REFERENCES:</p> <p>1. Mozhukhin, A.S.; Tyurikov, I.D.; Mitropolskaya, V.A. in <i>Fiz. Khim. Osn. Rektifikatsii</i>, Moskva, <u>1970</u>.</p>

<p>COMPONENTS:</p> <p>(1) Acetic acid 3-methyl-1-butyl ester (<i>isopentyl acetate</i>); C₇H₁₄O₂; [123-92-2]</p> <p>(2) Water; H₂O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Rao, D.S.; Rao, K.V.; Prasad, A.R.; Chiranjivi, C.</p> <p><i>J. Chem. Eng. Data</i> <u>1979</u>, 24, 241-4.</p>
<p>VARIABLES:</p> <p>T/K = 303</p>	<p>PREPARED BY:</p> <p>Z. Maczynska</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of acetic acid 3-methyl-1-butyl ester in water at 30°C was reported to be 0.19 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 2.6×10^{-4}.</p> <p>The solubility of water in acetic acid 3-methyl-1-butyl ester at 30°C was reported to be 0.15 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.011.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The titration method was used. An accurately weighed amount of one component was placed in a conical flask, the second component was titrated into the flask from a burette until the solution became turbid, and the amount titrated was recorded. The conical flask was kept in a constant-temperature bath. The data were reported together with the ternary system acetic acid 3-methyl-1-butyl ester-water-ethanenitrile (<i>isopentyl acetate-water-acetonitrile</i>).</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, reagent grade; used as received.</p> <p>(2) Distilled; free from CO₂.</p> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm 0.1^\circ\text{C}$ (authors). Soly. about ± 0.01 g(1)/100g sln and ± 0.6 g(2)/100g sln (compiler).</p> <p>REFERENCES:</p>

<p>COMPONENTS:</p> <p>(1) Acetic acid 3-methyl-1-butyl ester (<i>isopentyl acetate</i>); $C_7H_{14}O_2$; [123-92-2]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Skrzecz, A. <i>Pol. J. Chem.</i> <u>1981</u>, <i>55</i>, 1177-80.</p> <p>Skrzecz, A. <i>Thesis, Inst. Phys. Chem., Pol. Acad. Sci., Warszawa</i>, <u>1979</u>.</p>																																																	
<p>VARIABLES:</p> <p>$T/K = 288 - 359$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>																																																	
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of acetic acid 3-methyl-1-butyl ester and water</p> <table border="1" data-bbox="253 580 1243 989"> <thead> <tr> <th rowspan="2">T/K</th> <th colspan="2">x_1</th> <th colspan="2">g(1)/100g sln</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr> <td>288.4^a</td> <td>-</td> <td>0.9562</td> <td>-</td> <td>99.370</td> </tr> <tr> <td>296.9</td> <td>0.00029</td> <td>-</td> <td>0.209</td> <td>-</td> </tr> <tr> <td>306.3</td> <td>-</td> <td>0.9470</td> <td>-</td> <td>99.231</td> </tr> <tr> <td>310.1</td> <td>0.00026</td> <td>-</td> <td>0.188</td> <td>-</td> </tr> <tr> <td>320.7</td> <td>0.00025</td> <td>-</td> <td>0.180</td> <td>-</td> </tr> <tr> <td>325.9</td> <td>-</td> <td>0.9304</td> <td>-</td> <td>98.975</td> </tr> <tr> <td>339.3</td> <td>0.00025</td> <td>-</td> <td>0.180</td> <td>-</td> </tr> <tr> <td>344.4</td> <td>-</td> <td>0.9114</td> <td>-</td> <td>98.673</td> </tr> </tbody> </table> <p>^a By the Karl Fischer method. (continued next page)</p>		T/K	x_1		g(1)/100g sln		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	288.4 ^a	-	0.9562	-	99.370	296.9	0.00029	-	0.209	-	306.3	-	0.9470	-	99.231	310.1	0.00026	-	0.188	-	320.7	0.00025	-	0.180	-	325.9	-	0.9304	-	98.975	339.3	0.00025	-	0.180	-	344.4	-	0.9114	-	98.673
T/K	x_1		g(1)/100g sln																																															
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																														
288.4 ^a	-	0.9562	-	99.370																																														
296.9	0.00029	-	0.209	-																																														
306.3	-	0.9470	-	99.231																																														
310.1	0.00026	-	0.188	-																																														
320.7	0.00025	-	0.180	-																																														
325.9	-	0.9304	-	98.975																																														
339.3	0.00025	-	0.180	-																																														
344.4	-	0.9114	-	98.673																																														
<p>AUXILIARY INFORMATION</p>																																																		
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The synthetic method of Alexejew and the analytical method were used. An ampoule with the solution of fixed concentration was placed in a glass tube connected with a thermostat filled completely with distilled water. During the measurements the temperature of the bath was changed continuously and the appearance and disappearance of turbidity within the ampoule was observed visually. For the analytical method, The amount of water in the saturated organic phase was determined using a Karl Fischer titration procedure. The amount of water in the pure ester was taken into account.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Prolabo-Rhone Poulenc, pure grade; distilled; purity 99.97% by glc, 0.08 wt% water by the Karl Fischer method.</p> <p>(2) Distilled.</p> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm(0.2-0.6)^\circ C$. Soly. see above.</p> <p>REFERENCES:</p>																																																	

COMPONENTS: (1) Acetic acid 3-methyl-1-butyl ester (<i>isopentyl acetate</i>); $C_7H_{14}O_2$; [123-92-2] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Skrzecz, A. <i>Pol. J. Chem.</i> <u>1981</u> , 55, 1177-80. Skrzecz, A. <i>Thesis</i> , Inst. Phys. Chem., Pol. Acad. Sci., Warszawa, <u>1979</u> .
--------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

EXPERIMENTAL VALUES: (continued)

Mutual solubility of acetic acid 3-methyl-1-butyl ester and water

T/K	x_1		g(1)/100g sln	
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase
345.5	0.00026	-	0.188	-
358.6	0.00029	-	0.209	-
358.8	-	0.8804	-	98.155

Author's smoothing equations:

$$x_1 = 3.680 \times 10^{-4} - 4.37 \times 10^{-6} (T/K - 273.15) + 3.98 \times 10^{-8} (T/K - 273.15)^2$$

st.dev. = 1.82×10^{-6} (2)-rich phase

$$x_2 = 0.038754 + 2.150 \times 10^{-4} (T/K - 273.15) + 6.82 \times 10^{-6} (T/K - 273.15)^2$$

st. dev. = 6.68×10^{-4} (1)-rich phase

<p>COMPONENTS:</p> <p>(1) Acetic acid 3-methyl-1-butyl ester (isopentyl acetate); C₇H₁₄O₂; [123-92-2]</p> <p>(2) Water; H₂O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u>, 31, 56-70.</p>																																																											
<p>VARIABLES:</p> <p>T/K = 273 - 364</p>	<p>PREPARED BY:</p> <p>Z. Maczynska</p>																																																											
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of acetic acid 3-methyl-1-butyl ester and water</p> <hr/> <table border="1"> <thead> <tr> <th rowspan="2">t/°C</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x₁ (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>0.340</td><td>99.44</td><td>0.000472</td><td>0.9609</td></tr> <tr><td>9.1</td><td>0.265</td><td>99.33</td><td>0.000367</td><td>0.9535</td></tr> <tr><td>19.4</td><td>0.212</td><td>99.24</td><td>0.000294</td><td>0.9475</td></tr> <tr><td>30.3</td><td>0.208</td><td>99.20</td><td>0.000288</td><td>0.9449</td></tr> <tr><td>39.7</td><td>0.184</td><td>99.15</td><td>0.000255</td><td>0.9416</td></tr> <tr><td>50.0</td><td>0.174</td><td>98.89</td><td>0.000241</td><td>0.9250</td></tr> <tr><td>60.1</td><td>0.152</td><td>98.84</td><td>0.000210</td><td>0.9218</td></tr> <tr><td>70.2</td><td>0.203</td><td>98.76</td><td>0.000281</td><td>0.9168</td></tr> <tr><td>80.3</td><td>0.182</td><td>98.80</td><td>0.000252</td><td>0.9193</td></tr> <tr><td>90.7</td><td>0.205</td><td>98.66</td><td>0.000284</td><td>0.9106</td></tr> </tbody> </table> <hr/> <p>std. dev. 0.002 0.01</p>		t/°C	g(1)/100g sln		x ₁ (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	0.340	99.44	0.000472	0.9609	9.1	0.265	99.33	0.000367	0.9535	19.4	0.212	99.24	0.000294	0.9475	30.3	0.208	99.20	0.000288	0.9449	39.7	0.184	99.15	0.000255	0.9416	50.0	0.174	98.89	0.000241	0.9250	60.1	0.152	98.84	0.000210	0.9218	70.2	0.203	98.76	0.000281	0.9168	80.3	0.182	98.80	0.000252	0.9193	90.7	0.205	98.66	0.000284	0.9106
t/°C	g(1)/100g sln		x ₁ (compiler)																																																									
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																								
0	0.340	99.44	0.000472	0.9609																																																								
9.1	0.265	99.33	0.000367	0.9535																																																								
19.4	0.212	99.24	0.000294	0.9475																																																								
30.3	0.208	99.20	0.000288	0.9449																																																								
39.7	0.184	99.15	0.000255	0.9416																																																								
50.0	0.174	98.89	0.000241	0.9250																																																								
60.1	0.152	98.84	0.000210	0.9218																																																								
70.2	0.203	98.76	0.000281	0.9168																																																								
80.3	0.182	98.80	0.000252	0.9193																																																								
90.7	0.205	98.66	0.000284	0.9106																																																								
<p>AUXILIARY INFORMATION</p>																																																												
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, commercial sample; purity 99%; used as received.</p> <p>(2) Not specified.</p> <hr/> <p>ESTIMATED ERROR:</p> <p>Accuracy of method 0.1 wt% or less, for solubility, see above.</p> <hr/> <p>REFERENCES:</p>																																																											

<p>COMPONENTS:</p> <p>(1) Acetic acid pentyl ester (pentyl acetate); $C_7H_{14}O_2$; [628-63-7]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia January, 1989</p>
-------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION:

Quantitative solubility data for the acetic acid pentyl ester (1) - water (2) system have been reported in the publications listed in Table 1.

TABLE 1: Quantitative Solubility Studies of the
Acetic acid pentyl ester (1) - Water (2) System

Reference	T/K	Solubility	Method
Hemptinne (ref 1)	298	(1) in (2)	analytical
Park and Hopkins (ref 2)	298	(1) in (2)	unspecified
Doolittle (ref 3)	293	mutual	unspecified
Rao and Rao (ref 4)	303	mutual	titration
Venkataratnam et al. (ref 5)	303	mutual	titration
Rao and Rao (ref 6)	303	mutual	titration
Krupatkin and Shcherbakova (ref 7)	298	mutual	titration
Skrzecz (ref 8)	286-365	mutual	synthetic, Karl Fischer
Richon and Viillard (ref 9)	298	mutual	calorimetric, refractometric
Stephenson and Stuart (ref 10)	273-364	mutual	GLC

The original data in these publications are compiled in the Data Sheets immediately following this Critical Evaluation. For convenience, further discussion of this system will be divided into two parts.

1. SOLUBILITY OF ACETIC ACID PENTYL ESTER (1) IN WATER (2)

All the available data for the solubility of acetic acid pentyl ester (1) in water (2) are summarized in Table 2 with the following exclusions. The datum of Park and Hopkins (ref 2) reported in v/v units has been excluded from consideration. The datum of Krupatkin and Shcherbakova (ref 7) and the approximate values of Rao et al. (ref 4-6) appear to be high and have been rejected. Selected data are plotted in Figure 1.

(continued next page)

<p>COMPONENTS:</p> <p>(1) Acetic acid pentyl ester (pentyl acetate); $C_7H_{14}O_2$; [628-63-7]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia January, 1989</p>
-------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

TABLE 2: Recommended (R) and Tentative Solubilities
of Acetic acid pentyl ester (1) in Water (2)

T/K	Solubilities		
	Reported values g(1)/100g sln	"Best" values ($\pm\sigma_n$) ^a	
		g(1)/100g sln	10^4x_1
273	0.29 (ref 10)	0.29	4.0
283	0.224* (ref 8), 0.25* (ref 10)	0.24 \pm 0.01 (R)	3.3
293	0.195* (ref 8), 0.22* (ref 10)	0.21 \pm 0.01 (R)	2.9
298	0.162* (ref 1), 0.183* (ref 8), 0.178* ^b (ref 9), 0.19* (ref 10)	0.178 \pm 0.010 (R)	2.46
303	0.172* (ref 8), 0.18* (ref 10)	0.176 \pm 0.004 (R)	2.44
313	0.157* (ref 8), 0.15* (ref 10)	0.154 \pm 0.004 (R)	2.13
323	0.150* (ref 8), 0.10 (ref 10)	0.13 \pm 0.02	1.8
333	0.151* (ref 8), 0.10* (ref 10)	0.13 \pm 0.02	1.8
343	0.161* (ref 8), 0.17* (ref 10)	0.166 \pm 0.005 (R)	2.30
353	0.181* (ref 8), 0.17* (ref 10)	0.176 \pm 0.006 (R)	2.44
363	0.211* (ref 8)	0.21	2.9

^a Obtained by averaging where appropriate; σ_n has no statistical significance. Mole fraction solubilities (x_1) have the same status and (relative) percentage uncertainties as the mass % solubilities.

^b Average of two independent results reported by the original authors.

2. SOLUBILITY OF WATER (2) IN ACETIC ACID PENTYL ESTER (1)

All the available data for the solubility of water (2) in acetic acid pentyl ester (1) are summarized in Table 3 with the following exceptions.

The approximate values of Rao et al. (ref 4-6) are widely scattered and do not add any extra information to the more precise studies (ref 8-10) available and so have been rejected. The datum of Krupatkin and Shcherbakova (ref 7) at 298 K is significantly higher than all other studies (ref 8-10) and has also been rejected. Selected data are plotted in Figure 2.

(continued next page)

COMPONENTS:

- (1) Acetic acid pentyl ester
(pentyl acetate); $C_7H_{14}O_2$;
[628-63-7]
- (2) Water; H_2O ; [7732-18-5]

EVALUATOR:

G.T. Hefter, School of Mathematical
and Physical Sciences,
Murdoch University, Perth, W.A.,
Australia
January, 1989

CRITICAL EVALUATION: (continued)

Although the remaining data (ref 8-10) are in reasonable agreement over a wide temperature range the differences are generally too large to permit many values to be Recommended without further studies.




FIGURE 1. Selected data for the solubility of acetic acid pentyl ester (1) in water (2): ref 8 (●); ref 10 (O). Solid line is a least square polynomial fitted to the "Best" values from Table 2.

(continued next page)

<p>COMPONENTS:</p> <p>(1) Acetic acid pentyl ester (pentyl acetate); C₇H₁₄O₂; [628-63-7]</p> <p>(2) Water; H₂O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia January, 1989</p>
-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

TABLE 3: Recommended (R) and Tentative Solubilities
of Water (2) in Acetic acid pentyl ester (1)

T/K	Solubilities		
	Reported values	"Best" values ($\pm\sigma_n$) ^a	
	g(2)/100g sln	g(2)/100g sln	10 ² x ₂
273	0.79 (ref 10)	0.79	5
283	0.64* (ref 8), 0.81* (ref 10)	0.73 ± 0.09	5.1
293	0.63* (ref 8), 0.87* (ref 10)	0.75 ± 0.12	5.2
298	0.65* (ref 8), 0.91 (ref 9), 0.91* (ref 10)	0.82 ± 0.12	5.6
303	0.67* (ref 8), 0.96* (ref 10)	0.82 ± 0.14	5.6
313	0.75* (ref 8), 1.06* (ref 10)	0.9 ± 0.2	6
323	0.88* (ref 8), 1.16* (ref 10)	1.02 ± 0.14	6.9
333	1.04* (ref 8), 1.26* (ref 10)	1.15 ± 0.11	7.8
343	1.25* (ref 8), 1.34* (ref 10)	1.30 ± 0.05 (R)	8.7
353	1.51* (ref 8), 1.44* (ref 10)	1.48 ± 0.04 (R)	9.8
363	1.80* (ref 8), 1.55* (ref 10)	1.68 ± 0.13	11

^a Obtained by averaging where appropriate; σ_n has no statistical significance. Mole fraction solubilities (x_2) have the same status and (relative) percentage uncertainties as the mass % solubilities.

(continued next page)

COMPONENTS:

- (1) Acetic acid pentyl ester
(*pentyl acetate*); $C_7H_{14}O_2$;
[628-63-7]
- (2) Water; H_2O ; [7732-18-5]

EVALUATOR:

G.T. Hefter, School of Mathematical
and Physical Sciences,
Murdoch University, Perth, W.A.,
Australia
January, 1989

CRITICAL EVALUATION: (continued)




FIGURE 2. Selected data for the solubility of water (2) in acetic acid pentyl ester (1): ref 8 (●); ref 10 (○). Solid line is a least square polynomial fitted to the "Best" values from Table 3.

REFERENCES

1. Hemptinne, A. *Z. Phys. Chem.* 1894, 13, 561-9.
2. Park, J. G.; Hopkins, M. B. *Ind. Eng. Chem.* 1930, 22, 826-30.
3. Doolittle, A. K. *Ind. Eng. Chem.* 1935, 27, 1169-79.
4. Rao, J. R.; Rao, C. V. *J. Appl. Chem.* 1957, 7, 435-9.
5. Venkataratnam, A.; Rao, J. R.; Rao, C. V. *Chem. Eng. Sci.* 1957, 7, 102-10.
6. Rao, J. R.; Rao, C. V. *J. Appl. Chem.* 1959, 9, 69-73.
7. Krupatkin, I. L.; Shcherbakova, T. A. *Zh. Prikl. Khim.* 1971, 44, 307-11.

(continued next page)

COMPONENTS: (1) Acetic acid pentyl ester (<i>pentyl acetate</i>); C ₇ H ₁₄ O ₂ ; [628-63-7] (2) Water; H ₂ O; [7732-18-5]	EVALUATOR: G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia January, 1989
CRITICAL EVALUATION: (continued) 8. Skrzecz, A. <i>Pol. J. Chem.</i> <u>1980</u> , <i>54</i> , 1101-4; see also Skrzecz, A. <i>Thesis, I. Ch. F. PAN, Warszawa, 1979.</i> 9. Richon, D.; Viallard, A. <i>Fluid Phase Equil.</i> <u>1985</u> , <i>21</i> , 279-93. 10. Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u> , <i>31</i> , 56-70. ACKNOWLEDGEMENT The Evaluator thanks Dr. Brian Clare for the graphics.	

COMPONENTS: (1) Acetic acid pentyl ester (pentyl acetate); $C_7H_{14}O_2$; [628-63-7] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Hemptinne, A. <i>Z. Phys. Chem.</i> <u>1894</u> , 13, 561-9.
VARIABLES: $T/K = 298$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: The solubility of acetic acid pentyl ester in water at $25^\circ C$ was reported to be 1.622 g(1)/L sln.	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The analytical method was used. The mixture of water with excess ester was heated for some time in a water bath and the ester phase was filtered. A sample of known volume was then transferred to a smaller flask, heated with the known amount of baryta until complete saponification was obtained and then titrated. No further details were reported in the paper.	SOURCE AND PURITY OF MATERIALS: (1) Not specified. (2) Not specified. ESTIMATED ERROR: Not specified. REFERENCES:

COMPONENTS: (1) Acetic acid pentyl ester (pentyl acetate); $C_7H_{14}O_2$; [628-63-7] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Park, J.G.; Hopkins, M.B. <i>Ind. Eng. Chem.</i> <u>1930</u> , 22, 826-30.
VARIABLES: $T/K = 298$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: The solubility of acetic acid pentyl ester in water at $25^\circ C$ was reported to be 0.2 mL(1)/100mL(2).	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The method was not specified.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial samples; used as received. (2) Not specified.
	ESTIMATED ERROR: Not specified.
	REFERENCES:

COMPONENTS: (1) Acetic acid pentyl ester (pentyl acetate); $C_7H_{14}O_2$; [628-63-7] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Doolittle, A.K. <i>Ind. End. Chem.</i> <u>1935</u> , 27, 1169-79.
VARIABLES: $T/K = 293$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: The solubility of acetic acid pentyl ester in water at 20°C was reported to be 0.17 g(1)/100g sln. The corresponding mole fraction, x_1 , value calculated by the compiler is 2.4×10^{-4} . The solubility of water in acetic acid pentyl ester at 20°C was reported to be 1.15 g(2)/100g sln. The corresponding mole fraction, x_2 , value calculated by the compiler is 0.0776.	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The method was not specified.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial product; purity 84%, mixed isomers; b.p. range 127-155°C, d_4^{20} 0.862. (2) Not specified. ESTIMATED ERROR: Soly. about ± 0.03 g(1)/100g sln and ± 0.5 g(2)/100g sln (compiler). REFERENCES:

<p>COMPONENTS:</p> <p>(1) Acetic acid pentyl ester (<i>pentyl acetate</i>); $C_7H_{14}O_2$; [628-63-7]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Rao, J.R.; Rao, C.V. <i>J. Appl. Chem.</i> <u>1957</u>, 7, 435-9.</p>
<p>VARIABLES:</p> <p>$T/K = 303$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of acetic acid pentyl ester in water at 30°C was reported to be 0.4 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 6×10^{-4}.</p> <p>The solubility of water in acetic acid pentyl ester at 30°C was reported to be 1.7 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.111.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The titration method was used. The data and method were reported together with the ternary system acetic acid pentyl ester-water-methanol.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) May and Baker, reagent grade; 95% was distilled between 120 and 150°C, d^{30}_D 0.8630, n^{30}_D 1.3937.</p> <p>(2) Distilled; free from CO_2.</p> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm 0.02^\circ C$ (authors). Soly. about ± 0.2 g(1)/100g sln and ± 1.0 g(2)/100g sln (compiler).</p> <p>REFERENCES:</p>

COMPONENTS: (1) Acetic acid pentyl ester (pentyl acetate); $C_7H_{14}O_2$; [628-63-7] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Venkataratnam, A.; Rao, J.R.; Rao, C.V. <i>Chem. Eng. Sci.</i> <u>1957</u> , 7, 102-10.
VARIABLES: $T/K = 303$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: <p>The solubility of acetic acid pentyl ester in water at 30°C was reported to be 0.2 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 3×10^{-4}.</p> <p>The solubility of water in acetic acid pentyl ester at 30°C was reported to be 0.4 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.03.</p>	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: <p>The titration method described by Othmer, White and Trueger (ref 1) was used. The data were reported together with the ternary system acetic acid pentyl ester-water-2-propanone (pentyl acetate-water-acetone).</p>	SOURCE AND PURITY OF MATERIALS: (1) May and Baker, analytical grade; used as received; 95% distilled between 120 and 150°C. d^{30} 0.8630, n^{30} 1.3937. (2) Distilled; free of CO_2 . ESTIMATED ERROR: Soly. about ± 0.03 g(1)/100g sln and ± 0.3 g(2)/100g sln (compiler). REFERENCES: 1. Othmer, D.F.; White, R.E.; Trueger, E. <i>Ind. Eng. Chem.</i> <u>1941</u> , 33, 1240.

COMPONENTS: (1) Acetic acid pentyl ester (pentyl acetate); $C_7H_{14}O_2$; [628-63-7] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Rao, J.R.; Rao, C.V. <i>J. Appl. Chem.</i> <u>1959</u> , 9, 69-73.
VARIABLES: $T/K = 303$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: The solubility of acetic acid pentyl ester in water at 30°C was reported to be 0.4 g(1)/100g sln. The corresponding mole fraction, x_1 , value calculated by the compiler is 6×10^{-4} . The solubility of water in acetic acid pentyl ester at 30°C was reported to be 0.9 g(2)/100g sln and 99.2 g(1)/100g sln ^a . The corresponding mole fraction, x_2 , value calculated by the compiler is 0.06. ^a The sum of mass percent values is not equal 100%.	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: Presumably the titration method described by Rao and Rao (ref 1) was used. The data were reported together with the ternary system acetic acid pentyl ester-water-1-propanol.	SOURCE AND PURITY OF MATERIALS: (1) May and Barker, analytical grade; 95% had distilled between 120 and 150°C, d^{30}_D 0.8630, n^{30}_D 1.3937. (2) Distilled. ESTIMATED ERROR: Soly. about ± 0.2 g(1)/100g sln and ± 0.2 g(2)/100g sln (compiler). REFERENCES: 1. Rao, J.R.; Rao, C.V. <i>J. Appl. Chem.</i> <u>1957</u> , 7, 435.

COMPONENTS: (1) Acetic acid pentyl ester (pentyl acetate); $C_7H_{14}O_2$; [628-63-7] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Krupatkin, I.L.; Shcherbakova, T.A. <i>Zh. Prikl. Khim.</i> <u>1971</u> , 44, 307-11.
VARIABLES: $T/K = 298$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: The solubility of acetic acid pentyl ester in water at 25°C was reported to be 0.22 g(1)/100g sln. The corresponding mole fraction, x_1 , value calculated by the compiler is 3.05×10^{-4} . The solubility of water in acetic acid pentyl ester at 25°C was reported to be 1.20 g(2)/100g sln. The corresponding mole fraction, x_2 , value calculated by the compiler is 0.0807.	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The titration method was used. The amount of water in pure ester was taken into account. The data were reported together with the ternary system acetic acid pentyl ester-water-phosphoric acid.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, analytical grade; b.p. range 147-150°C. (2) Twice distilled. ESTIMATED ERROR: Temp. $\pm 0.1^\circ C$ (authors). Soly. about ± 0.03 g(1)/100g sln and ± 0.5 g(2)/100g sln (compiler). REFERENCES:

<p>COMPONENTS:</p> <p>(1) Acetic acid pentyl ester (pentyl acetate); C₇H₁₄O₂; [628-63-7]</p> <p>(2) Water; H₂O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Skrzecz, A. Pol. J. Chem. <u>1980</u>, 54, 1101-4. Skrzecz, A. Thesis, Inst. Phys. Chem., Pol. Acad. Sci., Warszawa, <u>1979</u>.</p>																																																																
<p>VARIABLES:</p> <p>T/K = 286 - 365</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>																																																																
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of acetic acid pentyl ester and water</p> <table border="1" data-bbox="256 560 1254 1104"> <thead> <tr> <th rowspan="2">T/K</th> <th colspan="2">x₁</th> <th colspan="2">g(1)/100g sln</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>285.6</td><td>0.00030</td><td>-</td><td>0.216</td><td>-</td></tr> <tr><td>293.2</td><td>-</td><td>0.9565^a</td><td>-</td><td>99.375</td></tr> <tr><td>299.9</td><td>0.00025</td><td>-</td><td>0.180</td><td>-</td></tr> <tr><td>305.6</td><td>-</td><td>0.9519</td><td>-</td><td>99.306</td></tr> <tr><td>306.1</td><td>0.00023</td><td>-</td><td>0.166</td><td>-</td></tr> <tr><td>328.6</td><td>-</td><td>0.9318</td><td>-</td><td>98.997</td></tr> <tr><td>343.6</td><td>-</td><td>0.9193</td><td>-</td><td>98.800</td></tr> <tr><td>345.9</td><td>0.00023</td><td>-</td><td>0.166</td><td>-</td></tr> <tr><td>353.1</td><td>0.00025</td><td>-</td><td>0.180</td><td>-</td></tr> <tr><td>358.6</td><td>-</td><td>0.8896</td><td>-</td><td>98.312</td></tr> <tr><td>364.5</td><td>0.00030</td><td>-</td><td>0.216</td><td>-</td></tr> </tbody> </table> <p>^a By the Karl Fischer method. Author's smoothing equations: $x_1 = 3.583 \times 10^{-4} - 5.48 \times 10^{-6} (T/K - 273.15) + 5.22 \times 10^{-8} (T/K - 273.15)^2$ st. dev. = 3.39×10^{-6} (2)-rich phase $x_2 = 0.045710 - 3.4441 \times 10^{-4} (T/K - 273.15) + 1.267 \times 10^{-4} (T/K - 273.15)^2$ st. dev. = 3.41×10^{-3} (1)-rich phase</p>		T/K	x ₁		g(1)/100g sln		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	285.6	0.00030	-	0.216	-	293.2	-	0.9565 ^a	-	99.375	299.9	0.00025	-	0.180	-	305.6	-	0.9519	-	99.306	306.1	0.00023	-	0.166	-	328.6	-	0.9318	-	98.997	343.6	-	0.9193	-	98.800	345.9	0.00023	-	0.166	-	353.1	0.00025	-	0.180	-	358.6	-	0.8896	-	98.312	364.5	0.00030	-	0.216	-
T/K	x ₁		g(1)/100g sln																																																														
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																													
285.6	0.00030	-	0.216	-																																																													
293.2	-	0.9565 ^a	-	99.375																																																													
299.9	0.00025	-	0.180	-																																																													
305.6	-	0.9519	-	99.306																																																													
306.1	0.00023	-	0.166	-																																																													
328.6	-	0.9318	-	98.997																																																													
343.6	-	0.9193	-	98.800																																																													
345.9	0.00023	-	0.166	-																																																													
353.1	0.00025	-	0.180	-																																																													
358.6	-	0.8896	-	98.312																																																													
364.5	0.00030	-	0.216	-																																																													
<p>AUXILIARY INFORMATION</p>																																																																	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The synthetic method of Alexejew and the analytical method were used. An ampoule with the solution of fixed concentration was placed into a glass tube connected with a thermostat filled completely with distilled water. The temperature of the bath was changed continuously during the measurements and the appearance and disappearance of turbidity within the ampoule was observed visually. For the analytical method, The amount of water in the saturated organic-phase was determined using a Karl Fischer titration procedure. The amount of water in the pure ester was taken into account.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) POCH Gliwice, pure for analysis grade; distilled; purity 99.92% by glc, 0.27 wt% water by the Karl Fischer method. (2) Distilled.</p> <p>ESTIMATED ERROR:</p> <p>Temp. ±(0.2-0.6)°C. Soly. see above.</p> <p>REFERENCES:</p>																																																																

COMPONENTS: (1) Acetic acid pentyl ester (pentyl acetate); $C_7H_{14}O_2$; [628-63-7] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Richon, D.; Viillard, A. <i>Fluid Phase Equilib.</i> <u>1985</u> , 21, 279-93.																						
VARIABLES: $T/K = 298$	PREPARED BY: A. Skrzecz																						
EXPERIMENTAL VALUES: Mutual solubility of acetic acid pentyl ester and water <table border="1" data-bbox="123 532 1173 747"> <thead> <tr> <th rowspan="2">T/K</th> <th rowspan="2">mol(1)/ 100g(2)</th> <th colspan="2">x_1</th> <th colspan="2">g(1)/100g sln (compiler)</th> </tr> <tr> <th>(2)-rich ph.</th> <th>(1)-rich ph.</th> <th>(2)-rich ph.</th> <th>(1)-rich ph.</th> </tr> </thead> <tbody> <tr> <td>298.1^a</td> <td>-</td> <td>2.33×10^{-4}</td> <td>0.938</td> <td>0.1681^d</td> <td>99.09^e</td> </tr> <tr> <td>298.1^b</td> <td>0.00144</td> <td>2.59×10^{-4c}</td> <td>-</td> <td>0.1871^d</td> <td>-</td> </tr> </tbody> </table> <p data-bbox="123 768 473 868"> ^a Calorimetric method. ^b Refractometric method. ^c Compiler. </p>		T/K	mol(1)/ 100g(2)	x_1		g(1)/100g sln (compiler)		(2)-rich ph.	(1)-rich ph.	(2)-rich ph.	(1)-rich ph.	298.1 ^a	-	2.33×10^{-4}	0.938	0.1681 ^d	99.09 ^e	298.1 ^b	0.00144	2.59×10^{-4c}	-	0.1871 ^d	-
T/K	mol(1)/ 100g(2)			x_1		g(1)/100g sln (compiler)																	
		(2)-rich ph.	(1)-rich ph.	(2)-rich ph.	(1)-rich ph.																		
298.1 ^a	-	2.33×10^{-4}	0.938	0.1681 ^d	99.09 ^e																		
298.1 ^b	0.00144	2.59×10^{-4c}	-	0.1871 ^d	-																		
AUXILIARY INFORMATION																							
METHOD/APPARATUS/PROCEDURE: The calorimetric and differential refractometric methods were used. The calorimetric method was described in the thesis of Richon (ref 1) and by Richon and Viillard (ref 2). For the refractometric measurements, a Phoenix model 1-2000T differential refractometer from Texas Instruments was used, and the solubility was determined from a characteristic calibration curve as described in the thesis of Richon (ref 1).	SOURCE AND PURITY OF MATERIALS: (1) Merck (for analysis); purified by preparation gas chromatographic method; purity >99.5%, water content was negligible. (2) Distilled. ESTIMATED ERROR: Soly. about $\pm 0.01^d$ and $\pm 0.2^e$ g(1)/100g sln (compiler). REFERENCES: 1. Richon, D. <i>Thesis</i> , University de Clermont-Ferrand, <u>1974</u> . 2. Richon, D.; Viillard, A. <i>Can. J. Chem.</i> <u>1976</u> , 54, 2584.																						

COMPONENTS: (1) Acetic acid pentyl ester (pentyl acetate); $C_7H_{14}O_2$; [628-63-7] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u> , 31, 56-70.																																																														
VARIABLES: $T/K = 273 - 364$	PREPARED BY: Z. Maczynska																																																														
EXPERIMENTAL VALUES: Mutual solubility of acetic acid pentyl ester and water <table border="1" data-bbox="259 544 1253 1058"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>0.29</td><td>99.21</td><td>0.00040</td><td>0.9456</td></tr> <tr><td>9.8</td><td>-</td><td>99.19</td><td>-</td><td>0.9443</td></tr> <tr><td>19.7</td><td>0.22</td><td>99.14</td><td>0.00030</td><td>0.9410</td></tr> <tr><td>30.6</td><td>0.16</td><td>98.78</td><td>0.00022</td><td>0.9180</td></tr> <tr><td>39.5</td><td>0.16</td><td>98.91</td><td>0.00022</td><td>0.9262</td></tr> <tr><td>50.0</td><td>0.10</td><td>98.88</td><td>0.00014</td><td>0.9243</td></tr> <tr><td>60.3</td><td>0.10</td><td>98.73</td><td>0.00014</td><td>0.9149</td></tr> <tr><td>70.2</td><td>0.17</td><td>98.63</td><td>0.00023</td><td>0.9088</td></tr> <tr><td>80.1</td><td>0.17</td><td>98.62</td><td>0.00023</td><td>0.9082</td></tr> <tr><td>90.5</td><td>-</td><td>98.46</td><td>-</td><td>0.8984</td></tr> </tbody> </table> <table border="1" data-bbox="259 1058 1253 1118"> <tbody> <tr> <td>std. dev.</td> <td>0.01</td> <td>0.02</td> </tr> </tbody> </table>		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	0.29	99.21	0.00040	0.9456	9.8	-	99.19	-	0.9443	19.7	0.22	99.14	0.00030	0.9410	30.6	0.16	98.78	0.00022	0.9180	39.5	0.16	98.91	0.00022	0.9262	50.0	0.10	98.88	0.00014	0.9243	60.3	0.10	98.73	0.00014	0.9149	70.2	0.17	98.63	0.00023	0.9088	80.1	0.17	98.62	0.00023	0.9082	90.5	-	98.46	-	0.8984	std. dev.	0.01	0.02
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																												
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																											
0	0.29	99.21	0.00040	0.9456																																																											
9.8	-	99.19	-	0.9443																																																											
19.7	0.22	99.14	0.00030	0.9410																																																											
30.6	0.16	98.78	0.00022	0.9180																																																											
39.5	0.16	98.91	0.00022	0.9262																																																											
50.0	0.10	98.88	0.00014	0.9243																																																											
60.3	0.10	98.73	0.00014	0.9149																																																											
70.2	0.17	98.63	0.00023	0.9088																																																											
80.1	0.17	98.62	0.00023	0.9082																																																											
90.5	-	98.46	-	0.8984																																																											
std. dev.	0.01	0.02																																																													
AUXILIARY INFORMATION																																																															
METHOD/AppARATUS/PROCEDURE: <p>The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.</p>	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial sample; purity 99%; used as received. (2) Not specified. ESTIMATED ERROR: Accuracy of method 0.1 wt% or less, for solubility, see above. REFERENCES:																																																														

COMPONENTS:	EVALUATOR:
(1) Butanoic acid 1-methylethyl ester (<i>isopropyl butyrate</i>); C ₇ H ₁₄ O ₂ ; [638-11-9]	G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia
(2) Water; H ₂ O; [7732-18-5]	January, 1989

CRITICAL EVALUATION:

Quantitative solubility data for the butanoic acid 1-methylethyl ester (1) - water (2) system have been reported in the publications listed in Table 1.

TABLE 1: Quantitative Solubility Studies of the Butanoic acid 1-methylethyl ester (1) - Water (2) System

Reference	T/K	Solubility	Method
Bomshtein et al. (ref 1)	283-363	mutual	titration
Stephenson and Stuart (ref 2)	273-363	mutual	GLC

The original data in these publications are compiled in the Data Sheets immediately following this Critical Evaluation. For convenience, further discussion of this system will be divided into two parts.

1. SOLUBILITY OF BUTANOIC ACID 1-METHYLETHYL ESTER (1) IN WATER (2)

All the available data (ref 1,2) for the solubility of butanoic acid 1-methylethyl ester (1) in water (2) are summarized in Table 2 and plotted in Figure 1. The data are in serious disagreement differing by up to an order of magnitude. Interestingly, the data of Bomshtein et al. (ref 1) are lower than those reported by Stephenson and Stuart (ref 2) whereas for most other systems studied by the same authors the reverse is true. In the absence of any other studies it is not possible to prefer one data set over the other and no "Best" values are proposed, however, the data of Stephenson and Stuart (ref 2) are usually reliable. Clearly this system warrants further investigation.

TABLE 2: Reported Solubilities of Butanoic acid 1-methylethyl ester (1) in Water (2)

T/K	Reported solubilities	
	g(1)/100g sln	10 ³ x ₁ ^a
273	3.12 (ref 2)	4.4
283	0.20 (ref 1), 2.91* (ref 2)	4.1
293	0.22 (ref 1), 2.74* (ref 2)	3.8
298	0.24* (ref 1), 2.67* (ref 2)	3.8

(continued next page)

<p>COMPONENTS:</p> <p>(1) Butanoic acid 1-methylethyl ester (<i>isopropyl butyrate</i>); $C_7H_{14}O_2$; [638-11-9]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia</p> <p>January, 1989</p>
-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

T/K	Reported solubilities	
	g(1)/100g sln	$10^3 x_1^a$
303	0.25 (ref 1), 2.61* (ref 2)	3.7
313	0.30 (ref 1), 2.53* (ref 2)	3.6
323	0.40 (ref 1), 2.49* (ref 2)	3.5
333	0.50 (ref 1), 2.56* (ref 2)	3.7
343	0.65 (ref 1), 2.66* (ref 2)	3.8
353	0.88 (ref 1), 2.82* (ref 2)	4.0
363	1.40 (ref 1), 3.00* (ref 2)	4.3

^a Values of ref 2, see text.




FIGURE 1. Data for the solubility of butanoic acid 1-methylethyl ester (1) in water (2): ref 1 (●); ref 2 (○). Solid lines are least squares polynomial fits and are included only for illustrative purposes.

(continued next page)

<p>COMPONENTS:</p> <p>(1) Butanoic acid 1-methylethyl ester (<i>isopropyl butyrate</i>); C₇H₁₄O₂; [638-11-9]</p> <p>(2) Water; H₂O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia January, 1989</p>
-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

2. SOLUBILITY OF WATER (2) IN BUTANOIC ACID 1-METHYLETHYL ESTER (1)

All the available data for the solubility of water (2) in butanoic acid 1-methylethyl ester (1) are summarized in Table 3 and plotted in Figure 2.

Although the two studies show a rather different dependence on temperature the results, unlike the aqueous phase solubilities, are in reasonable agreement and the average values may be regarded as Tentative pending further studies.

TABLE 3: Tentative Solubilities
of Water (2) in Butanoic acid 1-methylethyl ester (1)

T/K	Solubilities		
	Reported values g(2)/100g sln	"Best" values ($\pm\sigma_n$) ^a	
		g(2)/100g sln	x ₂
273	1.71 (ref 2)	1.7	0.11
283	0.90 (ref 1), 1.81* (ref 2)	1.4 ± 0.5	0.09
293	1.20 (ref 1), 1.92* (ref 2)	1.6 ± 0.4	0.11
298	1.35* (ref 1), 2.00* (ref 2)	1.7 ± 0.3	0.11
303	1.50 (ref 1), 2.06* (ref 2)	1.8 ± 0.3	0.12
313	1.80 (ref 1), 2.20* (ref 2)	2.0 ± 0.2	0.13
323	2.10 (ref 1), 2.37* (ref 2)	2.2 ± 0.2	0.14
333	2.30 (ref 1), 2.57* (ref 2)	2.4 ± 0.2	0.15
343	2.60 (ref 1), 2.78* (ref 2)	2.7 ± 0.1	0.17
353	3.40 (ref 1), 3.00* (ref 2)	3.2 ± 0.2	0.19
363	5.50 (ref 1), 3.20* (ref 2)	4.4 ± 1.3	0.25

^a Obtained by averaging where appropriate; σ_n has no statistical significance. Mole fraction solubilities (x₂) have the same status and (relative) percentage uncertainties as the mass % solubilities.

(continued next page)

<p>COMPONENTS:</p> <p>(1) Butanoic acid 1-methylethyl ester (<i>isopropyl butyrate</i>); C₇H₁₄O₂; [638-11-9]</p> <p>(2) Water; H₂O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia January, 1989</p>
-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)




FIGURE 2. Data for the solubility of water (2) in butanoic acid 1-methylethyl ester (1): ref 1 (●); ref 2 (○). Solid lines are least squares polynomial fits and are included only for illustrative purposes.

REFERENCES

1. Bomshtein, A. L.; Trofimov, A. N.; Serafimov, L. A. *Zh. Prikl. Khim.* **1978**, *51*, 1280-2.
2. Stephenson, R.; Stuart, J. *J. Chem. Eng. Data* **1986**, *31*, 56-70.

ACKNOWLEDGEMENT

The Evaluator thanks Dr. Brian Clare for the graphics.

<p>COMPONENTS:</p> <p>(1) Butanoic acid 1-methylethyl ester (<i>isopropyl butyrate</i>); $C_7H_{14}O_2$; [638-11-9]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Bomshtein, A.L.; Trofimov, A.N.; Serafimov, L.A.</p> <p><i>Zh. Prikl. Khim.</i> 1978, 51, 1280-2.</p>																																																						
<p>VARIABLES:</p> <p>$T/K = 283 - 363$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>																																																						
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of butanoic acid 1-methylethyl ester and water</p> <table border="1" data-bbox="145 539 1128 1013"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">x_1</th> <th colspan="2">g(1)/100g sln (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>10</td><td>0.00028</td><td>0.9384</td><td>0.20</td><td>99.10</td></tr> <tr><td>20</td><td>0.00031</td><td>0.9193</td><td>0.22</td><td>98.80</td></tr> <tr><td>30</td><td>0.00035</td><td>0.9009</td><td>0.25</td><td>98.50</td></tr> <tr><td>40</td><td>0.00042</td><td>0.8831</td><td>0.30</td><td>98.20</td></tr> <tr><td>50</td><td>0.00056</td><td>0.8659</td><td>0.40</td><td>97.90</td></tr> <tr><td>60</td><td>0.00069</td><td>0.8547</td><td>0.50</td><td>97.70</td></tr> <tr><td>70</td><td>0.00091</td><td>0.8384</td><td>0.65</td><td>97.40</td></tr> <tr><td>80</td><td>0.00123</td><td>0.7973</td><td>0.88</td><td>96.60</td></tr> <tr><td>90</td><td>0.00196</td><td>0.7041</td><td>1.40</td><td>94.50</td></tr> </tbody> </table>		$t/^\circ C$	x_1		g(1)/100g sln (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	10	0.00028	0.9384	0.20	99.10	20	0.00031	0.9193	0.22	98.80	30	0.00035	0.9009	0.25	98.50	40	0.00042	0.8831	0.30	98.20	50	0.00056	0.8659	0.40	97.90	60	0.00069	0.8547	0.50	97.70	70	0.00091	0.8384	0.65	97.40	80	0.00123	0.7973	0.88	96.60	90	0.00196	0.7041	1.40	94.50
$t/^\circ C$	x_1		g(1)/100g sln (compiler)																																																				
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																			
10	0.00028	0.9384	0.20	99.10																																																			
20	0.00031	0.9193	0.22	98.80																																																			
30	0.00035	0.9009	0.25	98.50																																																			
40	0.00042	0.8831	0.30	98.20																																																			
50	0.00056	0.8659	0.40	97.90																																																			
60	0.00069	0.8547	0.50	97.70																																																			
70	0.00091	0.8384	0.65	97.40																																																			
80	0.00123	0.7973	0.88	96.60																																																			
90	0.00196	0.7041	1.40	94.50																																																			
<p>AUXILIARY INFORMATION</p>																																																							
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The titration method was used at constant temperature. No further details were reported in the paper.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified; distilled; without impurities by glc, b.p. $125.0^\circ C$, n_D^{20} 1.394.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <p>REFERENCES:</p>																																																						

<p>COMPONENTS:</p> <p>(1) Butanoic acid 1-methylethyl ester (isopropyl butyrate); $C_7H_{14}O_2$; [638-11-9]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u>, <i>31</i>, 56-70.</p>																																																											
<p>VARIABLES:</p> <p>$T/K = 273 - 363$</p>	<p>PREPARED BY:</p> <p>Z. Maczynska</p>																																																											
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of butanoic acid 1-methylethyl ester and water</p> <table border="1" data-bbox="273 546 1257 1050"> <thead> <tr> <th rowspan="2">t/°C</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>3.12</td><td>98.29</td><td>0.00444</td><td>0.8883</td></tr> <tr><td>9.7</td><td>3.08</td><td>98.13</td><td>0.00438</td><td>0.8789</td></tr> <tr><td>19.5</td><td>2.65</td><td>98.18</td><td>0.00375</td><td>0.8819</td></tr> <tr><td>30.8</td><td>2.75</td><td>97.90</td><td>0.00390</td><td>0.8658</td></tr> <tr><td>39.5</td><td>2.57</td><td>97.91</td><td>0.00364</td><td>0.8663</td></tr> <tr><td>49.7</td><td>2.49</td><td>97.70</td><td>0.00352</td><td>0.8546</td></tr> <tr><td>60.2</td><td>2.49</td><td>97.39</td><td>0.00352</td><td>0.8377</td></tr> <tr><td>70.1</td><td>2.70</td><td>97.18</td><td>0.00382</td><td>0.8266</td></tr> <tr><td>80.0</td><td>2.84</td><td>96.93</td><td>0.00403</td><td>0.8137</td></tr> <tr><td>90.2</td><td>2.98</td><td>96.85</td><td>0.00423</td><td>0.8097</td></tr> </tbody> </table> <p>std. dev. 0.02 0.02</p>		t/°C	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	3.12	98.29	0.00444	0.8883	9.7	3.08	98.13	0.00438	0.8789	19.5	2.65	98.18	0.00375	0.8819	30.8	2.75	97.90	0.00390	0.8658	39.5	2.57	97.91	0.00364	0.8663	49.7	2.49	97.70	0.00352	0.8546	60.2	2.49	97.39	0.00352	0.8377	70.1	2.70	97.18	0.00382	0.8266	80.0	2.84	96.93	0.00403	0.8137	90.2	2.98	96.85	0.00423	0.8097
t/°C	g(1)/100g sln		x_1 (compiler)																																																									
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																								
0	3.12	98.29	0.00444	0.8883																																																								
9.7	3.08	98.13	0.00438	0.8789																																																								
19.5	2.65	98.18	0.00375	0.8819																																																								
30.8	2.75	97.90	0.00390	0.8658																																																								
39.5	2.57	97.91	0.00364	0.8663																																																								
49.7	2.49	97.70	0.00352	0.8546																																																								
60.2	2.49	97.39	0.00352	0.8377																																																								
70.1	2.70	97.18	0.00382	0.8266																																																								
80.0	2.84	96.93	0.00403	0.8137																																																								
90.2	2.98	96.85	0.00423	0.8097																																																								
<p>AUXILIARY INFORMATION</p>																																																												
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, commercial sample; purity 98%; used as received.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Accuracy of method 0.1 wt% or less, for solubility, see above.</p> <p>REFERENCES:</p>																																																											

COMPONENTS: (1) Butanoic acid, 3-methyl-, ethyl ester (ethyl isovalerate); $C_7H_{14}O_2$; [108-64-5] (2) Water; H_2O ; [7732-18-5]	EVALUATOR: G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia January, 1989
-----------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION:

Quantitative solubility data for the 3-methylbutanoic acid ethyl ester (1) - water (2) system have been reported in the publications listed in Table 1.

TABLE 1: Quantitative Solubility Studies of the 3-Methylbutanoic acid ethyl ester (1) - Water (2) System

Reference	T/K	Solubility	Method
Bancroft (ref 1)	293	mutual	titration
Chang and Moulton (ref 2)	298	mutual	titration
Stephenson and Stuart (ref 3)	273-364	mutual	GLC

The original data in these publications are compiled in the Data Sheets immediately following this Critical Evaluation. For convenience, further discussion of this system will be divided into two parts.

1. SOLUBILITY OF 3-METHYLBUTANOIC ACID ETHYL ESTER (1) IN WATER (2)

All the available data for the solubility of 3-methylbutanoic acid ethyl ester (1) in water (2) are summarized in Table 2 except for the datum of Bancroft which is reported in v/v units and hence excluded from this Evaluation.

At 298 K, the only temperature where comparison is possible, the datum of Chang and Moulton (ref 2), is in quite good agreement with the interpolated value of Stephenson and Stuart (ref 3), lending confidence to the latter's values at other temperatures. Nevertheless, pending further studies, the data of Stephenson and Stuart must be regarded as Tentative although it may be noted that these authors' data are usually reliable.

TABLE 2: Tentative Solubilities of 3-Methylbutanoic acid ethyl ester (1) in Water (2)

T/K	Solubilities		
	Reported values	"Best" values ^a	
	g(1)/100g sln	g(1)/100g sln	10^4x_1
273	0.34 (ref 3)	0.3	4
283	0.24 (ref 3)	0.2	3
293	0.21* (ref 3)	0.2	3
298	0.16 (ref 2), 0.20* (ref 3)	0.20	2.8

(continued next page)

COMPONENTS: (1) Butanoic acid, 3-methyl-, ethyl ester (ethyl isovalerate); $C_7H_{14}O_2$; [108-64-5] (2) Water; H_2O ; [7732-18-5]	EVALUATOR: G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia January, 1989
-----------------------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

T/K	Solubilities		
	Reported values	"Best" values ^a	
	g(1)/100g sln	g(1)/100g sln	10^4x_1
303	0.19* (ref 3)	0.2	3
313	0.17* (ref 3)	0.2	3
323	0.16* (ref 3)	0.2	3
333	0.16* (ref 3)	0.2	3
343	0.16* (ref 3)	0.2	3
353	0.16* (ref 3)	0.2	3
363	0.16* (ref 3)	0.2	3

^a Rounded values of ref 3.

2. SOLUBILITY OF WATER (2) IN 3-METHYLBUTANOIC ACID ETHYL ESTER (1)

The situation with regard to the solubility of water (2) in 3-methylbutanoic acid ethyl ester (1) is identical to that of the H_2O -rich phase and the comments under Part 1. of this Evaluation are equally applicable and should be consulted in conjunction with Table 3.

TABLE 3: Tentative Solubilities of Water (2) in 3-Methylbutanoic acid ethyl ester (1)

T/K	Solubilities		
	Reported values	"Best" values ^a	
	g(2)/100g sln	g(2)/100g sln	10^2x_2
273	0.41 (ref 3)	0.4	3
283	0.48* (ref 3)	0.5	3
293	0.52* (ref 3)	0.5	3
298	0.47 (ref 2), 0.61* (ref 3)	0.54 ^b	3.8
303	0.69* (ref 3)	0.7	5
313	0.71* (ref 3)	0.7	5
323	0.79* (ref 3)	0.8	6

(continued next page)

COMPONENTS: (1) Butanoic acid, 3-methyl-, ethyl ester (ethyl isovalerate); C ₇ H ₁₄ O ₂ ; [108-64-5] (2) Water; H ₂ O; [7732-18-5]	EVALUATOR: G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia January, 1989
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

T/K	Solubilities		
	Reported values	"Best" values ^a	
	g(2)/100g sln	g(2)/100g sln	10 ² x ₂
333	0.83* (ref 3)	0.8	6
343	0.85* (ref 3)	0.9	6
353	0.94* (ref 3)	0.9	6
363	1.11* (ref 3)	1.1	8

^a Rounded values of ref 3.

^b Average value.

REFERENCES

1. Bancroft, W. D. *Phys. Rev.* 1895, 3, 114-36.
2. Chang, Y. C.; Moulton, R. W. *Ind. Eng. Chem.* 1953, 45, 2350-61.
3. Stephenson, R.; Stuart, J. *J. Chem. Eng. Data* 1986, 31, 56-70.

<p>COMPONENTS:</p> <p>(1) Butanoic acid, 3-methyl-, ethyl ester (ethyl isovalerate); $C_7H_{14}O_2$; [108-64-5]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Bancroft, W.D. <i>Phys. Rev.</i> <u>1895</u>, 3, 114-36.</p>
<p>VARIABLES:</p> <p>$T/K = 293$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of 3-methylbutanoic acid ethyl ester in water at 20°C was reported to be 0.02 mL(1)/10mL(2).</p> <p>The solubility of water in 3-methylbutanoic acid ethyl ester at 20°C was reported to be 0.04 mL(2)/10mL(1).</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The titration method was used. 10 mL of solvent in a test tube was titrated with the second component until the solution became cloudy.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Laboratory source; dried over $CaCl_2$, distilled. (2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Soly. ± 0.01 mL.</p> <p>REFERENCES:</p>

<p>COMPONENTS:</p> <p>(1) Butanoic acid, 3-methyl-, ethyl ester (ethyl isovalerate); $C_7H_{14}O_2$; [108-64-5]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Chang, Yi-C.; Moulton, R.W. <i>Ind. Eng. Chem.</i> <u>1953</u>, 45, 2350-61.</p>
<p>VARIABLES:</p> <p>$T/K = 298$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of 3-methylbutanoic acid ethyl ester in water at 25°C was reported to be 0.16 g(1)/100g(2). The corresponding mass per cent and mole fraction, x_1, values calculated by the compiler are 0.16 g(1)/100g sln and 2.2×10^{-4}.</p> <p>The solubility of water in 3-methylbutanoic acid ethyl ester at 25°C was reported to be 0.47 g(2)/100g(1). The corresponding mass per cent and mole fraction, x_2, values calculated by the compiler are 0.47 g(2)/100g sln and 0.033.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The titration method was used. The 100g portion of solvent in a thermostated container was titrated with the second component from a 1 mL pipette calibrated to 0.01 mL. An average of two closest values (definitely cloud and definitely clear) was presented. The amount of solute (about 0.2 mL) was very difficult to judge and it was accurate only to 0.05 mL. The data were reported together with the ternary system 3-methylbutanoic acid ethyl ester-water-benzene.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified; washed with dilute KOH and H_2O, dehydrated with anhydron, fractionally distilled; purity 99.8% by a saponification test, b.p. range 133-134°C at 755 mm Hg, n_D^{20} 1.3968.</p> <p>(2) Distilled (distillation from $KMnO_4$ caused no change in the refractive index).</p> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm 0.1^\circ C$. Soly. $\pm 25\%$ (probably relative error).</p> <p>REFERENCES:</p>

<p>COMPONENTS:</p> <p>(1) Butanoic acid, 3-methyl-, ethyl ester (ethyl isovalerate); $C_7H_{14}O_2$; [108-64-5]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u>, <i>31</i>, 56-70.</p>																																																											
<p>VARIABLES:</p> <p>$T/K = 273 - 364$</p>	<p>PREPARED BY:</p> <p>Z. Maczynska</p>																																																											
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of 3-methylbutanoic acid ethyl ester and water</p> <table border="1" data-bbox="249 592 1251 1096"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>0.34</td><td>99.588</td><td>0.00047</td><td>0.97097</td></tr> <tr><td>9.0</td><td>0.24</td><td>99.527</td><td>0.00033</td><td>0.96679</td></tr> <tr><td>19.4</td><td>0.22</td><td>99.442</td><td>0.00030</td><td>0.96103</td></tr> <tr><td>30.6</td><td>0.20</td><td>99.355</td><td>0.00028</td><td>0.95519</td></tr> <tr><td>39.8</td><td>0.17</td><td>99.293</td><td>0.00023</td><td>0.95106</td></tr> <tr><td>49.5</td><td>0.16</td><td>99.211</td><td>0.00022</td><td>0.94565</td></tr> <tr><td>60.1</td><td>0.15</td><td>99.173</td><td>0.00021</td><td>0.94316</td></tr> <tr><td>70.0</td><td>0.16</td><td>99.155</td><td>0.00022</td><td>0.94199</td></tr> <tr><td>80.3</td><td>0.16</td><td>99.057</td><td>0.00022</td><td>0.93563</td></tr> <tr><td>90.5</td><td>0.16</td><td>98.877</td><td>0.00022</td><td>0.92415</td></tr> </tbody> </table> <p>std. dev. 0.01 0.004</p>		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	0.34	99.588	0.00047	0.97097	9.0	0.24	99.527	0.00033	0.96679	19.4	0.22	99.442	0.00030	0.96103	30.6	0.20	99.355	0.00028	0.95519	39.8	0.17	99.293	0.00023	0.95106	49.5	0.16	99.211	0.00022	0.94565	60.1	0.15	99.173	0.00021	0.94316	70.0	0.16	99.155	0.00022	0.94199	80.3	0.16	99.057	0.00022	0.93563	90.5	0.16	98.877	0.00022	0.92415
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																									
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																								
0	0.34	99.588	0.00047	0.97097																																																								
9.0	0.24	99.527	0.00033	0.96679																																																								
19.4	0.22	99.442	0.00030	0.96103																																																								
30.6	0.20	99.355	0.00028	0.95519																																																								
39.8	0.17	99.293	0.00023	0.95106																																																								
49.5	0.16	99.211	0.00022	0.94565																																																								
60.1	0.15	99.173	0.00021	0.94316																																																								
70.0	0.16	99.155	0.00022	0.94199																																																								
80.3	0.16	99.057	0.00022	0.93563																																																								
90.5	0.16	98.877	0.00022	0.92415																																																								
<p>AUXILIARY INFORMATION</p>																																																												
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, commercial sample; purity 99%; used as received.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Accuracy of method 0.1 wt% or less, for solubility, see above.</p> <p>REFERENCES:</p>																																																											

COMPONENTS: (1) Butanoic acid propyl ester (propyl butyrate); $C_7H_{14}O_2$; [105-66-8] (2) Water; H_2O ; [7732-18-5]	EVALUATOR: G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia January, 1989
----------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION:

Quantitative solubility data for the butanoic acid propyl ester (1) - water (2) system have been reported in the publications listed in Table 1.

TABLE 1: Quantitative Solubility Studies of the Butanoic acid propyl ester (1) - Water (2) System

Reference	T/K	Solubility	Method
Fuehner (ref 1)	290	(1) in (2)	titration
Stephenson and Stuart (ref 2)	273-364	mutual	GLC

The original data in these publications are compiled in the Data Sheets immediately following this Critical Evaluation. For convenience, further discussion of this system will be divided into two parts.

1. SOLUBILITY OF BUTANOIC ACID PROPYL ESTER (1) IN WATER (2)

All the available data are summarized in Table 2. At 290 K, the only temperature where comparison is possible, the values of Fuehner (ref 1) and Stephenson and Stuart (ref 2) are in good agreement giving confidence in the remaining data of ref 2. However, in the absence of independent studies the data of Stephenson and Stuart must be regarded as Tentative although it may be noted that the solubilities of these authors are usually reliable.

TABLE 2: Tentative Solubilities of Butanoic acid propyl ester (1) in Water (2)

T/K	Solubilities		
	Reported values	"Best" values ^a	
	g(1)/100g sln	g(1)/100g sln	10^4x_1
273	0.30 (ref 2)	0.3	4
283	0.22* (ref 2)	0.2	3
290	0.162 (ref 1), 0.19* (ref 2)	0.18	2.5
293	0.19* (ref 2)	0.2	3
298	0.18* (ref 2)	0.2	3
303	0.17* (ref 2)	0.2	3

(continued next page)

COMPONENTS: (1) Butanoic acid propyl ester (propyl butyrate); $C_7H_{14}O_2$; [105-66-8] (2) Water; H_2O ; [7732-18-5]	EVALUATOR: G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia January, 1989
----------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

T/K	Solubilities		
	Reported values	"Best" values ^a	
	g(1)/100g sln	g(1)/100g sln	10^4x_1
313	0.16* (ref 2)	0.2	3
323	0.16* (ref 2)	0.2	3
333	0.16* (ref 2)	0.2	3
343	0.16* (ref 2)	0.2	3
353	0.16* (ref 2)	0.2	3
363	0.16* (ref 2)	0.2	3

^a Rounded values of ref 2 except at 290 K.

2. SOLUBILITY OF WATER (2) IN BUTANOIC ACID PROPYL ESTER (1)

Only the data of Stephenson and Stuart (ref 2) are available for the solubility of water (2) in propyl butyrate (1) and so no Critical Evaluation is possible. The interested user is referred to the relevant Data Sheet for the experimental solubilities but it may be noted that the data of Stephenson and Stuart (ref 2) are usually reliable.

REFERENCES

1. Fuehner, H. *Ber. Dtsch. Chem. Ges.* 1924, 57, 510-5.
2. Stephenson, R.; Stuart, J. *J. Chem. Eng. Data* 1986, 31, 56-70.

COMPONENTS: (1) Butanoic acid propyl ester (propyl butyrate); $C_7H_{14}O_2$; [105-66-8] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Fuehner, H. Ber. Dtsch. Chem. Ges. <u>1924</u> , 57, 510-5.
VARIABLES: $T/K = 290$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: The solubility of butanoic acid propyl ester in water at $17^\circ C$ was reported to be 0.195 vol%, 0.162 g(1)/100g sln and 0.0124 mol(1)/L sln. The corresponding mole fraction, x_1 , value calculated by the compiler is 2.24×10^{-4} .	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The titration method was used. The ester was added from pipette to the flask with a constant amount of water (50, 100 or 1000 mL) so long as, on shaking, the mixture remained transparent.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial product. (2) Not specified. ESTIMATED ERROR: Not specified. REFERENCES:

<p>COMPONENTS:</p> <p>(1) Butanoic acid propyl ester (propyl butyrate); $C_7H_{14}O_2$; [105-66-8]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u>, <i>31</i>, 56-70.</p>																																																																
<p>VARIABLES:</p> <p>$T/K = 273 - 364$</p>	<p>PREPARED BY:</p> <p>Z. Maczynska</p>																																																																
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of butanoic acid propyl ester and water</p> <table border="1" data-bbox="259 534 1253 1038"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr> <td>0</td> <td>0.30</td> <td>99.36</td> <td>0.00042</td> <td>0.9555</td> </tr> <tr> <td>9.6</td> <td>0.23</td> <td>-</td> <td>0.00032</td> <td>-</td> </tr> <tr> <td>20.1</td> <td>0.19</td> <td>99.37</td> <td>0.00026</td> <td>0.9562</td> </tr> <tr> <td>29.6</td> <td>0.15</td> <td>99.36</td> <td>0.00021</td> <td>0.9555</td> </tr> <tr> <td>39.5</td> <td>0.14</td> <td>99.34</td> <td>0.00019</td> <td>0.9542</td> </tr> <tr> <td>49.8</td> <td>0.17</td> <td>99.16</td> <td>0.00023</td> <td>0.9423</td> </tr> <tr> <td>60.1</td> <td>0.21</td> <td>98.52</td> <td>0.00029</td> <td>0.9021</td> </tr> <tr> <td>70.1</td> <td>0.15</td> <td>-</td> <td>0.00021</td> <td>-</td> </tr> <tr> <td>80.1</td> <td>0.14</td> <td>99.01</td> <td>0.00019</td> <td>0.9326</td> </tr> <tr> <td>90.5</td> <td>0.17</td> <td>98.19</td> <td>0.00023</td> <td>0.8824</td> </tr> <tr> <td>std. dev.</td> <td>0.01</td> <td>0.01</td> <td></td> <td></td> </tr> </tbody> </table>		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	0.30	99.36	0.00042	0.9555	9.6	0.23	-	0.00032	-	20.1	0.19	99.37	0.00026	0.9562	29.6	0.15	99.36	0.00021	0.9555	39.5	0.14	99.34	0.00019	0.9542	49.8	0.17	99.16	0.00023	0.9423	60.1	0.21	98.52	0.00029	0.9021	70.1	0.15	-	0.00021	-	80.1	0.14	99.01	0.00019	0.9326	90.5	0.17	98.19	0.00023	0.8824	std. dev.	0.01	0.01		
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																														
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																													
0	0.30	99.36	0.00042	0.9555																																																													
9.6	0.23	-	0.00032	-																																																													
20.1	0.19	99.37	0.00026	0.9562																																																													
29.6	0.15	99.36	0.00021	0.9555																																																													
39.5	0.14	99.34	0.00019	0.9542																																																													
49.8	0.17	99.16	0.00023	0.9423																																																													
60.1	0.21	98.52	0.00029	0.9021																																																													
70.1	0.15	-	0.00021	-																																																													
80.1	0.14	99.01	0.00019	0.9326																																																													
90.5	0.17	98.19	0.00023	0.8824																																																													
std. dev.	0.01	0.01																																																															
<p>AUXILIARY INFORMATION</p>																																																																	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, commercial sample; purity 99%; used as received.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Accuracy of method 0.1 wt% or less, for solubility, see above.</p> <p>REFERENCES:</p>																																																																

COMPONENTS: (1) Formic acid hexyl ester (hexyl formate); C ₇ H ₁₄ O ₂ ; [629-33-4] (2) Water; H ₂ O; [7732-18-5]	ORIGINAL MEASUREMENTS: Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u> , 31, 56-70.																																																																
VARIABLES: T/K = 273 - 364	PREPARED BY: Z. Maczynska																																																																
EXPERIMENTAL VALUES: Mutual solubility of formic acid hexyl ester and water <table border="1" data-bbox="133 534 1127 1038"> <thead> <tr> <th rowspan="2">t/°C</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x₁ (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr> <td>0</td> <td>-</td> <td>99.474</td> <td>-</td> <td>0.96319</td> </tr> <tr> <td>8.7</td> <td>0.17</td> <td>99.475</td> <td>0.00024</td> <td>0.96326</td> </tr> <tr> <td>19.7</td> <td>0.15</td> <td>99.492</td> <td>0.00021</td> <td>0.96441</td> </tr> <tr> <td>29.7</td> <td>0.13</td> <td>99.401</td> <td>0.00018</td> <td>0.95826</td> </tr> <tr> <td>39.6</td> <td>0.13</td> <td>99.340</td> <td>0.00018</td> <td>0.95418</td> </tr> <tr> <td>49.8</td> <td>0.11</td> <td>99.346</td> <td>0.00015</td> <td>0.95458</td> </tr> <tr> <td>60.2</td> <td>0.11</td> <td>99.281</td> <td>0.00015</td> <td>0.95027</td> </tr> <tr> <td>70.3</td> <td>0.12</td> <td>-</td> <td>0.00017</td> <td>-</td> </tr> <tr> <td>80.2</td> <td>0.13</td> <td>99.153</td> <td>0.00018</td> <td>0.94185</td> </tr> <tr> <td>90.6</td> <td>0.14</td> <td>99.201</td> <td>0.00019</td> <td>0.94499</td> </tr> <tr> <td>std. dev.</td> <td>0.01</td> <td>0.003</td> <td></td> <td></td> </tr> </tbody> </table>		t/°C	g(1)/100g sln		x ₁ (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	-	99.474	-	0.96319	8.7	0.17	99.475	0.00024	0.96326	19.7	0.15	99.492	0.00021	0.96441	29.7	0.13	99.401	0.00018	0.95826	39.6	0.13	99.340	0.00018	0.95418	49.8	0.11	99.346	0.00015	0.95458	60.2	0.11	99.281	0.00015	0.95027	70.3	0.12	-	0.00017	-	80.2	0.13	99.153	0.00018	0.94185	90.6	0.14	99.201	0.00019	0.94499	std. dev.	0.01	0.003		
t/°C	g(1)/100g sln		x ₁ (compiler)																																																														
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																													
0	-	99.474	-	0.96319																																																													
8.7	0.17	99.475	0.00024	0.96326																																																													
19.7	0.15	99.492	0.00021	0.96441																																																													
29.7	0.13	99.401	0.00018	0.95826																																																													
39.6	0.13	99.340	0.00018	0.95418																																																													
49.8	0.11	99.346	0.00015	0.95458																																																													
60.2	0.11	99.281	0.00015	0.95027																																																													
70.3	0.12	-	0.00017	-																																																													
80.2	0.13	99.153	0.00018	0.94185																																																													
90.6	0.14	99.201	0.00019	0.94499																																																													
std. dev.	0.01	0.003																																																															
AUXILIARY INFORMATION																																																																	
METHOD/APPARATUS/PROCEDURE: The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial sample; purity 96%; used as received. (2) Not specified. ESTIMATED ERROR: Accuracy of method 0.1 wt% or less, for solubility, see above. REFERENCES:																																																																

<p>COMPONENTS:</p> <p>(1) Hexanoic acid methyl ester (methyl caproate); $C_7H_{14}O_2$; [106-70-7]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u>, <i>31</i>, 56-70.</p>																																																											
<p>VARIABLES:</p> <p>$T/K = 273 - 364$</p>	<p>PREPARED BY:</p> <p>Z. Maczynska</p>																																																											
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of hexanoic acid methyl ester and water</p> <table border="1" data-bbox="251 544 1246 1048"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>0.241</td><td>99.62</td><td>0.000334</td><td>0.9732</td></tr> <tr><td>9.3</td><td>0.177</td><td>99.54</td><td>0.000245</td><td>0.9677</td></tr> <tr><td>20.0</td><td>0.166</td><td>99.45</td><td>0.000230</td><td>0.9616</td></tr> <tr><td>29.7</td><td>0.167</td><td>99.40</td><td>0.000231</td><td>0.9582</td></tr> <tr><td>39.5</td><td>0.129</td><td>99.32</td><td>0.000179</td><td>0.9528</td></tr> <tr><td>50.0</td><td>0.137</td><td>99.25</td><td>0.000190</td><td>0.9482</td></tr> <tr><td>60.0</td><td>0.132</td><td>99.19</td><td>0.000183</td><td>0.9442</td></tr> <tr><td>70.2</td><td>0.128</td><td>99.20</td><td>0.000177</td><td>0.9449</td></tr> <tr><td>80.2</td><td>0.138</td><td>99.17</td><td>0.000191</td><td>0.9429</td></tr> <tr><td>90.5</td><td>0.147</td><td>99.11</td><td>0.000204</td><td>0.9390</td></tr> </tbody> </table> <p>std. dev. 0.001 0.01</p>		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	0.241	99.62	0.000334	0.9732	9.3	0.177	99.54	0.000245	0.9677	20.0	0.166	99.45	0.000230	0.9616	29.7	0.167	99.40	0.000231	0.9582	39.5	0.129	99.32	0.000179	0.9528	50.0	0.137	99.25	0.000190	0.9482	60.0	0.132	99.19	0.000183	0.9442	70.2	0.128	99.20	0.000177	0.9449	80.2	0.138	99.17	0.000191	0.9429	90.5	0.147	99.11	0.000204	0.9390
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																									
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																								
0	0.241	99.62	0.000334	0.9732																																																								
9.3	0.177	99.54	0.000245	0.9677																																																								
20.0	0.166	99.45	0.000230	0.9616																																																								
29.7	0.167	99.40	0.000231	0.9582																																																								
39.5	0.129	99.32	0.000179	0.9528																																																								
50.0	0.137	99.25	0.000190	0.9482																																																								
60.0	0.132	99.19	0.000183	0.9442																																																								
70.2	0.128	99.20	0.000177	0.9449																																																								
80.2	0.138	99.17	0.000191	0.9429																																																								
90.5	0.147	99.11	0.000204	0.9390																																																								
<p>AUXILIARY INFORMATION</p>																																																												
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, commercial sample; purity 99%; used as received.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Accuracy of method 0.1 wt% or less, for solubility, see above.</p> <p>REFERENCES:</p>																																																											

<p>COMPONENTS:</p> <p>(1) Pentanoic acid ethyl ester (ethyl valerate); $C_7H_{14}O_2$; [539-82-2]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia January, 1989</p>
---------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION:

Quantitative solubility data for the pentanoic acid ethyl ester (1) - water (2) system have been reported in the publications listed in Table 1.

TABLE 1: Quantitative Solubility Studies of the
Pentanoic acid ethyl ester (1) - Water (2) System

Reference	T/K	Solubility	Method
Hemptinne (ref 1)	298	(1) in (2)	analytical
Bomshtein et al. (ref 2)	283-353	mutual	titration

The original data in these publications are compiled in the Data Sheets immediately following this Critical Evaluation. For convenience, further discussion of this system will be divided into two parts.

1. SOLUBILITY OF PENTANOIC ACID ETHYL ESTER (1) IN WATER (2)

For the solubility of pentanoic acid ethyl ester (1) in water (2) all the available data are summarized in Table 2. At 298 K, the only temperature where comparison is possible, the datum of Hemptinne (ref 1) is in good agreement with the interpolated value of Bomshtein et al. (ref 2). However, it should be noted that the solubilities reported by Bomshtein et al. often differ significantly from the "Best" values in well-characterized systems. For this reason the data in Table 2 should be regarded as very Tentative.

(continued next page)

<p>COMPONENTS:</p> <p>(1) Pentanoic acid ethyl ester (ethyl valerate); $C_7H_{14}O_2$; [539-82-2]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia January, 1989</p>
---------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

TABLE 2: Tentative Solubilities
of Pentanoic acid ethyl ester (1) in Water (2)

T/K	Solubilities		
	Reported values	"Best" values ^a	
	g(1)/100g sln	g(1)/100g sln	10^4x_1
283	0.24 (ref 2)	0.2	3
293	0.25 (ref 2)	0.25	3.5
298	0.237 (ref 1), 0.255* (ref 2)	0.25	3.5
303	0.26 (ref 2)	0.3	4
313	0.27 (ref 2)	0.3	4
323	0.32 (ref 2)	0.3	4
333	0.37 (ref 2)	0.4	6
343	0.55 (ref 2)	0.6	8
353	0.78 (ref 2)	0.8	11

^a Rounded values of ref (2), but see text.

2. SOLUBILITY OF WATER (2) IN PENTANOIC ACID ETHYL ESTER (1)

As the only data available for the solubility of water (2) in pentanoic acid ethyl ester (1) are those of Bomshtein *et al.* (ref 2) no Critical Evaluation is possible. The interested user is referred to the relevant Data Sheet but the comments made under part 1. of this Evaluation above should be noted.

REFERENCES

- Hemptinne, A. *Z. Phys. Chem.* 1894, 13, 561-9.
- Bomshtein, A. L.; Trofimov, A. N.; Serafimov, L. A. *Zh. Prikl. Khim.* 1978, 51, 1280-2.

<p>COMPONENTS:</p> <p>(1) Pentanoic acid ethyl ester (ethyl valerate); $C_7H_{14}O_2$; [539-82-2]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Hemptinne, A. <i>Z. Phys. Chem.</i> <u>1894</u>, 13, 561-9.</p>
<p>VARIABLES:</p> <p>$T/K = 298$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of pentanoic acid ethyl ester in water at $25^\circ C$ was reported to be 2.366 g(1)/L sln.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method was used. The mixture of water with excess ester was heated for some time in a water bath and the ester phase was filtered. A sample of known volume was then transferred to a smaller flask, heated with the known amount of baryta until complete saponification was obtained and then titrated. No further details were reported in the paper.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Not specified. (2) Not specified.</p>
	<p>ESTIMATED ERROR:</p> <p>Not specified.</p>
	<p>REFERENCES:</p>

COMPONENTS: (1) Pentanoic acid ethyl ester (ethyl valerate); $C_7H_{14}O_2$; [539-82-2] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Bomshtein, A.L.; Trofimov, A.N.; Serafimov, L.A. <i>Zh. Prikl. Khim.</i> <u>1978</u> , 51, 1280-2.																																																	
VARIABLES: $T/K = 283 - 353$	PREPARED BY: A. Skrzecz																																																	
EXPERIMENTAL VALUES: Mutual solubility of pentanoic acid ethyl ester and water <table border="1" data-bbox="249 530 1243 963"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">x_1</th> <th colspan="2">g(1)/100g sln (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>10</td><td>0.00033</td><td>0.9928</td><td>0.24</td><td>99.90</td></tr> <tr><td>20</td><td>0.00034</td><td>0.9650</td><td>0.25</td><td>99.50</td></tr> <tr><td>30</td><td>0.00036</td><td>0.9257</td><td>0.26</td><td>98.90</td></tr> <tr><td>40</td><td>0.00038</td><td>0.9009</td><td>0.27</td><td>98.50</td></tr> <tr><td>50</td><td>0.00044</td><td>0.8715</td><td>0.32</td><td>98.00</td></tr> <tr><td>60</td><td>0.00051</td><td>0.8492</td><td>0.37</td><td>97.60</td></tr> <tr><td>70</td><td>0.00077</td><td>0.8226</td><td>0.55</td><td>97.10</td></tr> <tr><td>80</td><td>0.00109</td><td>0.7828</td><td>0.78</td><td>96.30</td></tr> </tbody> </table>		$t/^\circ C$	x_1		g(1)/100g sln (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	10	0.00033	0.9928	0.24	99.90	20	0.00034	0.9650	0.25	99.50	30	0.00036	0.9257	0.26	98.90	40	0.00038	0.9009	0.27	98.50	50	0.00044	0.8715	0.32	98.00	60	0.00051	0.8492	0.37	97.60	70	0.00077	0.8226	0.55	97.10	80	0.00109	0.7828	0.78	96.30
$t/^\circ C$	x_1		g(1)/100g sln (compiler)																																															
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																														
10	0.00033	0.9928	0.24	99.90																																														
20	0.00034	0.9650	0.25	99.50																																														
30	0.00036	0.9257	0.26	98.90																																														
40	0.00038	0.9009	0.27	98.50																																														
50	0.00044	0.8715	0.32	98.00																																														
60	0.00051	0.8492	0.37	97.60																																														
70	0.00077	0.8226	0.55	97.10																																														
80	0.00109	0.7828	0.78	96.30																																														
AUXILIARY INFORMATION																																																		
METHOD/APPARATUS/PROCEDURE: The titration method was used at constant temperature. No further details were reported in the paper.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified; distilled; without impurities by glc, b.p. $145.2^\circ C$, n_D^{20} 1.400. (2) Not specified. ESTIMATED ERROR: Not specified. REFERENCES:																																																	

<p>COMPONENTS:</p> <p>(1) Propanoic acid butyl ester (butyl propionate); $C_7H_{14}O_2$; [590-01-2]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia January, 1989</p>
-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION:

Quantitative solubility data for the propanoic acid butyl ester (1) - water (2) system have been reported in the publications listed in Table 1.

TABLE 1: Quantitative Solubility Studies of the Propanoic acid butyl ester (1) - Water (2) System

Reference	T/K	Solubility	Method
Bridgman (ref 1)	280-303	(2) in (1)	synthetic
Park and Hopkins (ref 2)	298	(1) in (2)	unspecified
Doolittle (ref 3)	293	mutual	unspecified
Bomshtein et al. (ref 4)	368	mutual	titration
Stephenson and Stuart (ref 5)	273-364	mutual	GLC

The original data in these publications are compiled in the Data Sheets immediately following this Critical Evaluation. For convenience, further discussion of this system will be divided into two parts.

1. SOLUBILITY OF PROPANOIC ACID BUTYL ESTER (1) IN WATER (2)

Apart from the single temperature values of Doolittle (ref 3) and Bomshtein et al. (ref 4), only the data of Stephenson and Stuart (ref 5) are available for the solubilities of propanoic acid butyl ester (1) in water (2) over a range of temperatures. The datum of Doolittle (ref 3) is in reasonable agreement with that of Stephenson and Stuart (ref 5) but Bomshtein et al.'s (ref 4) value is an order of magnitude higher. In the absence of confirmatory studies it is not possible to assess the quality of these data although it may be noted that the data of Stephenson and Stuart (ref 5) are generally reliable and those of Bomshtein et al. (ref 4) are often high.

(continued next page)

<p>COMPONENTS:</p> <p>(1) Propanoic acid butyl ester (butyl propionate); C₇H₁₄O₂; [590-01-2]</p> <p>(2) Water; H₂O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia January, 1989</p>
-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

TABLE 2: Reported Solubilities
of Propanoic acid butyl ester (1) in Water (2)

T/K	Reported Solubilities	
	g(1)/100g sln	10 ⁴ x ₁ ^a
273	0.298 (ref 5)	4.1
283	0.228* (ref 5)	3.2
293	0.15 (ref 3), 0.193* (ref 5)	2.7
298	0.180* (ref 5)	2.5
303	0.169* (ref 5)	2.3
313	0.155* (ref 5)	2.1
323	0.146* (ref 5)	2.0
333	0.143* (ref 5)	2.0
343	0.141* (ref 5)	1.9
353	0.142* (ref 5)	2.0
363	1.4 ^b (ref 4), 0.142* (ref 5)	2.0

^a Data of Stephenson and Stuart (ref 5).

^b 368 K (boiling point of mixture).

2. SOLUBILITY OF WATER (2) IN PROPANOIC ACID BUTYL ESTER (1)

All the available data for the solubility of water (2) in propanoic acid butyl ester (1) are summarized in Table 3. The situation for the organic-rich phase is better than for the aqueous phase, with the values of Bridgman (ref 1) being in excellent agreement with those of Stephenson and Stuart (ref 5) from 283-303 K. At other temperatures only the data of Stephenson and Stuart are available and in the absence of confirmatory studies must be regarded as Tentative.

(continued next page)

<p>COMPONENTS:</p> <p>(1) Propanoic acid butyl ester (butyl propionate); C₇H₁₄O₂; [590-01-2]</p> <p>(2) Water; H₂O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia January, 1989</p>
-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	--------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

TABLE 3: Recommended (R) and Tentative Solubilities
of Water (2) in Propanoic acid butyl ester (1)

T/K	Solubilities		
	Reported values	"Best" values ($\pm\sigma_n$) ^a	
	g(2)/100g sln	g(2)/100g sln	10 ² x ₂
273	0.48 (ref 5)	0.5	4
283	0.470* (ref 1), 0.53* (ref 5)	0.50 ± 0.03 (R)	3.5
293	0.546* (ref 1), 0.80 ^b (ref 3), 0.62* (ref 5)	0.58 ± 0.04 (R)	4.0
298	0.606* (ref 1), 0.68* (ref 5)	0.64 ± 0.04 (R)	4.5
303	0.686 (ref 1), 0.73* (ref 5)	0.71 ± 0.02 (R)	4.9
313	0.82* (ref 5)	0.8	6
323	0.86* (ref 5)	0.9	6
333	0.89* (ref 5)	0.9	6
343	0.92* (ref 5)	0.9	6
353	0.94* (ref 5)	0.9	6
363	3.76 ^c (ref 4), 1.00* (ref 5)	1.0	7

^a Obtained by averaging where appropriate; σ_n has no statistical significance. Mole fraction solubilities (x₂) have the same status and (relative) percentage uncertainties as the mass % solubilities.

^b Not included in calculation of "Best" value, see text.

^c 368 K; not included in calculation of "Best" value, see text.

REFERENCES

1. Bridgman, J. A. *Ind. Eng. Chem.* 1928, 20, 184-7.
2. Park, J. G.; Hopkins, M. B. *Ind. Eng. Chem.* 1930, 22, 826-30.
3. Doolittle, A. K. *Ind. Eng. Chem.* 1935, 27, 1169-79.
4. Bomshtein, A. L.; Trofimov, A. N.; Serafimov, L. A. *Zh. Prikl. Khim.* 1984, 57, 18-23.
5. Stephenson, R.; Stuart, J. *J. Chem. Eng. Data* 1986, 31, 56-70.

<p>COMPONENTS:</p> <p>(1) Propanoic acid butyl ester (<i>butyl propionate</i>); $C_7H_{14}O_2$; [590-01-2]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Bridgman, J.A. <i>Ind. End. Chem.</i> <u>1928</u>, 20, 184-7.</p>																				
<p>VARIABLES:</p> <p>$T/K = 280 - 303$</p>	<p>PREPARED BY:</p> <p>Z. Maczynska</p>																				
<p>EXPERIMENTAL VALUES:</p> <p style="text-align: center;">Solubility of water in propanoic acid butyl ester</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: center;">$t/^\circ C$</th> <th style="text-align: center;">$g(2)/100g(1)$</th> <th style="text-align: center;">$g(2)/100g$ solution (compiler)</th> <th style="text-align: center;">x_2 (compiler)</th> </tr> </thead> <tbody> <tr> <td style="text-align: center;">7.0</td> <td style="text-align: center;">0.452</td> <td style="text-align: center;">0.450</td> <td style="text-align: center;">0.0316</td> </tr> <tr> <td style="text-align: center;">16.0</td> <td style="text-align: center;">0.513</td> <td style="text-align: center;">0.510</td> <td style="text-align: center;">0.0357</td> </tr> <tr> <td style="text-align: center;">22.5</td> <td style="text-align: center;">0.576</td> <td style="text-align: center;">0.573</td> <td style="text-align: center;">0.0400</td> </tr> <tr> <td style="text-align: center;">30.0</td> <td style="text-align: center;">0.691</td> <td style="text-align: center;">0.686</td> <td style="text-align: center;">0.0475</td> </tr> </tbody> </table>		$t/^\circ C$	$g(2)/100g(1)$	$g(2)/100g$ solution (compiler)	x_2 (compiler)	7.0	0.452	0.450	0.0316	16.0	0.513	0.510	0.0357	22.5	0.576	0.573	0.0400	30.0	0.691	0.686	0.0475
$t/^\circ C$	$g(2)/100g(1)$	$g(2)/100g$ solution (compiler)	x_2 (compiler)																		
7.0	0.452	0.450	0.0316																		
16.0	0.513	0.510	0.0357																		
22.5	0.576	0.573	0.0400																		
30.0	0.691	0.686	0.0475																		
<p>AUXILIARY INFORMATION</p>																					
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The synthetic method similar to that described by Groschuff (ref 1) was used. A glass tube of about 100 mL capacity with a stopcock was filled with weighed (1) and (2). By shaking the tube after it had been warmed somewhat, the water was dissolved completely in the ester, after which the tube was placed in a 2-liter beaker filled with water and equipped with an agitator. By alternate slow cooling and warming, it was possible to determine, within about $0.5^\circ C$, the temperature at which water was precipitated from the solution, as shown by the clouding of the liquid which would clear again when the temperature was slightly increased.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified; distilled; b.p. range $144-146^\circ C$; boiled for some time in open flask before used.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm 0.25^\circ C$.</p> <p>REFERENCES:</p> <p>1. Groschuff, E. <i>Z. Elektrochem.</i> <u>1911</u>, 17, 348.</p>																				

COMPONENTS: (1) Propanoic acid butyl ester (<i>butyl propionate</i>); $C_7H_{14}O_2$; [590-01-2] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Park, J.G.; Hopkins, M.B. <i>Ind. Eng. Chem.</i> <u>1930</u> , 22, 826-30.
VARIABLES: $T/K = 298$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: The solubility of propanoic acid butyl ester in water at 25°C was reported to be 0.2 mL(1)/100mL(2).	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The method was not specified.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial samples; used as received. (2) Not specified.
	ESTIMATED ERROR: Not specified.
	REFERENCES:

<p>COMPONENTS:</p> <p>(1) Propanoic acid butyl ester (butyl propionate); $C_7H_{14}O_2$; [590-01-2]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Doolittle, A.K. <i>Ind. End. Chem.</i> <u>1935</u>, 27, 1169-79.</p>
<p>VARIABLES:</p> <p>$T/K = 293$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of propanoic acid butyl ester in water at 20°C was reported to be 0.15 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 2.1×10^{-4}.</p> <p>The solubility of water in propanoic acid butyl ester at 20°C was reported to be 0.80 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.055.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The method was not specified.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, commercial product; purity 99%, b.p. range 124-171°C, d_4^{20} 0.874.</p> <p>(2) Not specified.</p> <hr/> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <hr/> <p>REFERENCES:</p>

COMPONENTS: (1) Propanoic acid butyl ester (butyl propionate); $C_7H_{14}O_2$; [590-01-2] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Bomshtein, A.L.; Trofimov, A.N.; Serafimov, L.A. Zh. Prikl. Khim. <u>1984</u> , 57 18-23.
VARIABLES: $T/K = 368$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: <p>The solubility of propanoic acid butyl ester in water at $94.8^\circ C^a$ was reported to be $x_1 = 0.002$. The corresponding mass per cent value calculated by the compiler is 1.4 g(1)/100g sln.</p> <p>The solubility of water in propanoic acid butyl ester at $94.8^\circ C^a$ was reported to be $x_2 = 0.220$. The corresponding mass per cent value calculated by the compiler is 3.76 g(2)/100g sln.</p> <p>^a Boiling temperature at 101.32 kPa.</p>	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: <p>The titration method was used. The data and method were reported together with the ternary system propanoic acid butyl ester-water-acetic acid. No further details were reported in the paper.</p>	SOURCE AND PURITY OF MATERIALS: (1) Source not specified; distilled; without impurities by glc, b.p. $145.3^\circ C$, n_D^{20} 1.4011. (2) Not specified.
ESTIMATED ERROR: Not specified.	
REFERENCES:	

COMPONENTS: (1) Propanoic acid butyl ester (butyl propionate); $C_7H_{14}O_2$; [590-01-2] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u> , 31, 56-70.																																																																
VARIABLES: $T/K = 273 - 364$	PREPARED BY: Z. Maczynska																																																																
EXPERIMENTAL VALUES: Mutual solubility of propanoic acid butyl ester and water <hr/> <table border="1" data-bbox="249 532 1247 1040"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>0.298</td><td>99.52</td><td>0.000413</td><td>0.9663</td></tr> <tr><td>9.4</td><td>0.230</td><td>99.47</td><td>0.000319</td><td>0.9629</td></tr> <tr><td>19.8</td><td>0.197</td><td>99.38</td><td>0.000273</td><td>0.9568</td></tr> <tr><td>30.6</td><td>0.167</td><td>99.14</td><td>0.000231</td><td>0.9410</td></tr> <tr><td>40.2</td><td>0.144</td><td>99.15</td><td>0.000199</td><td>0.9417</td></tr> <tr><td>50.0</td><td>0.146</td><td>99.15</td><td>0.000202</td><td>0.9417</td></tr> <tr><td>59.9</td><td>0.142</td><td>99.11</td><td>0.000197</td><td>0.9390</td></tr> <tr><td>70.1</td><td>0.140</td><td>99.08</td><td>0.000194</td><td>0.9371</td></tr> <tr><td>80.2</td><td>0.147</td><td>99.07</td><td>0.000204</td><td>0.9365</td></tr> <tr><td>90.4</td><td>0.142</td><td>99.00</td><td>0.000197</td><td>0.9320</td></tr> <tr> <td>std. dev.</td> <td>0.003</td> <td>0.01</td> <td></td> <td></td> </tr> </tbody> </table>		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	0.298	99.52	0.000413	0.9663	9.4	0.230	99.47	0.000319	0.9629	19.8	0.197	99.38	0.000273	0.9568	30.6	0.167	99.14	0.000231	0.9410	40.2	0.144	99.15	0.000199	0.9417	50.0	0.146	99.15	0.000202	0.9417	59.9	0.142	99.11	0.000197	0.9390	70.1	0.140	99.08	0.000194	0.9371	80.2	0.147	99.07	0.000204	0.9365	90.4	0.142	99.00	0.000197	0.9320	std. dev.	0.003	0.01		
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																														
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																													
0	0.298	99.52	0.000413	0.9663																																																													
9.4	0.230	99.47	0.000319	0.9629																																																													
19.8	0.197	99.38	0.000273	0.9568																																																													
30.6	0.167	99.14	0.000231	0.9410																																																													
40.2	0.144	99.15	0.000199	0.9417																																																													
50.0	0.146	99.15	0.000202	0.9417																																																													
59.9	0.142	99.11	0.000197	0.9390																																																													
70.1	0.140	99.08	0.000194	0.9371																																																													
80.2	0.147	99.07	0.000204	0.9365																																																													
90.4	0.142	99.00	0.000197	0.9320																																																													
std. dev.	0.003	0.01																																																															
AUXILIARY INFORMATION																																																																	
METHOD/APPARATUS/PROCEDURE: <p>The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.</p>	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial sample; purity 99%; used as received. (2) Not specified. <hr/> ESTIMATED ERROR: Accuracy of method 0.1.wt% or less, for solubility, see above. <hr/> REFERENCES:																																																																

COMPONENTS: (1) Propanoic acid, 2,2-dimethyl-, ethyl ester (ethyl trimethylacetate); C ₇ H ₁₄ O ₂ ; [3938-95-2] (2) Water; H ₂ O; [7732-18-5]	ORIGINAL MEASUREMENTS: Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u> , <i>31</i> , 56-70.																																																											
VARIABLES: T/K = 273 - 364	PREPARED BY: Z. Maczynska																																																											
EXPERIMENTAL VALUES: Mutual solubility of 2,2-dimethylpropanoic acid ethyl ester and water <hr/> <table border="1" data-bbox="133 592 1120 1093"> <thead> <tr> <th rowspan="2">t/°C</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x₁ (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>0.304</td><td>99.65</td><td>0.000422</td><td>0.9752</td></tr> <tr><td>9.5</td><td>0.298</td><td>99.57</td><td>0.000413</td><td>0.9697</td></tr> <tr><td>20.0</td><td>0.183</td><td>99.49</td><td>0.000253</td><td>0.9643</td></tr> <tr><td>30.7</td><td>0.168</td><td>99.39</td><td>0.000233</td><td>0.9575</td></tr> <tr><td>40.1</td><td>0.135</td><td>99.33</td><td>0.000187</td><td>0.9535</td></tr> <tr><td>50.0</td><td>0.152</td><td>99.29</td><td>0.000210</td><td>0.9508</td></tr> <tr><td>60.2</td><td>0.126</td><td>99.24</td><td>0.000174</td><td>0.9475</td></tr> <tr><td>70.3</td><td>0.120</td><td>99.22</td><td>0.000166</td><td>0.9462</td></tr> <tr><td>80.4</td><td>0.124</td><td>99.18</td><td>0.000172</td><td>0.9436</td></tr> <tr><td>90.4</td><td>0.112</td><td>99.19</td><td>0.000155</td><td>0.9443</td></tr> </tbody> </table> <hr/> std. dev. 0.004 0.01		t/°C	g(1)/100g sln		x ₁ (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	0.304	99.65	0.000422	0.9752	9.5	0.298	99.57	0.000413	0.9697	20.0	0.183	99.49	0.000253	0.9643	30.7	0.168	99.39	0.000233	0.9575	40.1	0.135	99.33	0.000187	0.9535	50.0	0.152	99.29	0.000210	0.9508	60.2	0.126	99.24	0.000174	0.9475	70.3	0.120	99.22	0.000166	0.9462	80.4	0.124	99.18	0.000172	0.9436	90.4	0.112	99.19	0.000155	0.9443
t/°C	g(1)/100g sln		x ₁ (compiler)																																																									
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																								
0	0.304	99.65	0.000422	0.9752																																																								
9.5	0.298	99.57	0.000413	0.9697																																																								
20.0	0.183	99.49	0.000253	0.9643																																																								
30.7	0.168	99.39	0.000233	0.9575																																																								
40.1	0.135	99.33	0.000187	0.9535																																																								
50.0	0.152	99.29	0.000210	0.9508																																																								
60.2	0.126	99.24	0.000174	0.9475																																																								
70.3	0.120	99.22	0.000166	0.9462																																																								
80.4	0.124	99.18	0.000172	0.9436																																																								
90.4	0.112	99.19	0.000155	0.9443																																																								
AUXILIARY INFORMATION																																																												
METHOD/APPARATUS/PROCEDURE: The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial sample; purity 98%; used as received. (2) Not specified.																																																											
ESTIMATED ERROR: Accuracy of method 0.1 wt% or less, for solubility, see above.																																																												
REFERENCES:																																																												

COMPONENTS: (1) Acetic acid 3-methoxy-1-butyl ester (3-methoxybutyl acetate); C ₇ H ₁₄ O ₃ ; [4435-53-4] (2) Water; H ₂ O; [7732-18-5]	ORIGINAL MEASUREMENTS: Doolittle, A.K. <i>Ind. End. Chem.</i> <u>1935</u> , 27, 1169-79.
VARIABLES: <i>T/K</i> = 293	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: <p>The solubility of acetic acid 3-methoxy-1-butyl ester in water at 20°C was reported to be 6.46 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 0.00844.</p> <p>The solubility of water in acetic acid 3-methoxy-1-butyl ester at 20°C was reported to be 3.72 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.239.</p>	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The method was not specified.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial product; purity 99%, b.p. range 135-173°C, d_4^{20} 0.956. (2) Not specified.
ESTIMATED ERROR: Not specified.	
REFERENCES:	

COMPONENTS: (1) Propanoic acid, 2-hydroxy-, 1-butyl ester (<i>butyl lactate</i>); $C_7H_{14}O_3$; [138-22-7] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Doolittle, A.K. <i>Ind. End. Chem.</i> <u>1935</u> , 27, 1169-79.
VARIABLES: $T/K = 293$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: The solubility of 2-hydroxypropanoic acid 1-butyl ester in water at 20°C was reported to be 4.0 g(1)/100g sln. The corresponding mole fraction, x_1 , value calculated by the compiler is 0.0051. The solubility of water in 2-hydroxypropanoic acid 1-butyl ester at 20°C was reported to be 14.5 g(2)/100g sln. The corresponding mole fraction, x_2 , value calculated by the compiler is 0.579.	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The method was not specified.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial product; purity 97%, b.p. range 145-230°C, d_4^{20} 0.980. (2) Not specified.
	ESTIMATED ERROR: Not specified.
	REFERENCES:

<p>COMPONENTS:</p> <p>(1) Acetic acid phenyl ester (<i>phenyl acetate</i>); $C_8H_8O_2$; [122-79-2]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Richon, D.; Viillard, A. <i>Fluid Phase Equilib.</i> <u>1985</u>, <i>21</i>, 279-93.</p>
<p>VARIABLES:</p> <p>$T/K = 298$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of acetic acid phenyl ester in water at 298.1 K was reported to be 0.00416 mol(1)/100 g(2). The corresponding mass per cent and mole fraction, x_1, values calculated by the compiler are 0.558 g(1)/100g sln and 7.42×10^{-4}.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The refractometric method was used. The Phoenix model 1-2000T differential refractometer from Texas Instruments was used and the solubility was determined from a characteristic calibration curve as described in the thesis of Richon (ref 1).</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Merck (for synthesis); purified by preparation gas chromatographic method; purity >99.5%, water content was negligible.</p> <p>(2) Distilled.</p> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <p>REFERENCES:</p> <p>1. Richon, D. <i>Thesis</i>, University de Clermont-Ferrand, <u>1974</u>.</p>

<p>COMPONENTS:</p> <p>(1) Benzoic acid methyl ester (methyl benzoate); $C_8H_8O_2$; [93-58-3]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988</p>
-----------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION:

Quantitative solubility data for the benzoic acid methyl ester (1) - water (2) system have been reported in the publications listed in Table 1.

TABLE 1: Quantitative Solubility Studies of the Benzoic acid methyl ester (1) - Water (2) System

Reference	T/K	Solubility	Method
Gilbert and Lauer (ref 1)	288-308	mutual	titration
Richon and Viillard (ref 2)	298	(1) in (2)	refractometric
Stephenson and Stuart (ref 3)	273-364	mutual	GLC

The original data in these publications are compiled in the Data Sheets immediately following this Critical Evaluation. For convenience, further discussion of this system will be divided into two parts.

1. SOLUBILITY OF BENZOIC ACID METHYL ESTER (1) IN WATER (2)

All the available data for the solubility of benzoic acid methyl ester (1) in water (2) are summarized in Table 2 except that the approximate values of Gilbert and Lauer (ref 1) have been rejected because they are in general much higher than the other studies (ref 2,3).

At 298 K, the only temperature where comparison is possible, the data of Richon and Viillard (ref 2) and Stephenson and Stuart (ref 3) are in good agreement enabling the average value to be Recommended. At other temperatures only the data of Stephenson and Stuart (ref 3) are available and, pending further studies, they must be regarded as Tentative although it may be noted that the solubilities of these authors are usually reliable.

(continued next page)

<p>COMPONENTS:</p> <p>(1) Benzoic acid methyl ester (methyl benzoate); $C_8H_8O_2$; [93-58-3]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988</p>
-----------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

TABLE 2: Recommended (R) and Tentative Solubilities
of Benzoic acid methyl ester (1) in Water (2)

T/K	Solubilities		
	Reported values	"Best" values ^a	
	g(1)/100g sln	g(1)/100g sln	$10^4 x_1$
273	0.221 (ref 3)	0.22	2.9
283	0.220* (ref 3)	0.22	2.9
293	0.223* (ref 3)	0.22	2.9
298	0.202 (ref 2), 0.227* (ref 3)	0.215 ± 0.013^b (R)	2.85^b
303	0.233* (ref 3)	0.23	3.0
313	0.245* (ref 3)	0.25	3.3
323	0.264* (ref 3)	0.26	3.4
333	0.287* (ref 3)	0.29	3.8
343	0.315* (ref 3)	0.32	4.3
353	0.347* (ref 3)	0.35	4.7
363	0.405* (ref 3)	0.41	5.4

^a Rounded values of ref 3 except at 298 K.

^b Average value; x_1 has the same status and (relative) percentage uncertainty as the mass % solubility.

2. SOLUBILITY OF WATER (2) IN BENZOIC ACID METHYL ESTER (1)

All the data for the solubility of water (2) in benzoic acid methyl ester (1) are summarized in Table 3. The two data sets (ref 1,3) are in fair agreement but in the absence of other independent studies the reported solubilities must be regarded as Tentative since the aqueous phase data of Gilbert and Lauer (ref 1) were rejected (see above). However, it may be noted that the data of Stephenson and Stuart (ref 3) are usually reliable.

(continued next page)

<p>COMPONENTS:</p> <p>(1) Benzoic acid methyl ester (methyl benzoate); $C_8H_8O_2$; [93-58-3]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988</p>
-----------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

TABLE 3: Tentative Solubilities
of Water (2) in Benzoic acid methyl ester (1)

T/K	Solubilities		
	Reported values	"Best" values ^a	
	g(2)/100g sln	g(2)/100g sln	10^2x_2
273	0.46 (ref 3)	0.5	4
283	0.54* (ref 3)	0.5	4
293	0.4* (ref 1), 0.62* (ref 3)	0.6	5
298	0.6* (ref 1), 0.67* (ref 3)	0.6	5
303	0.9* (ref 1), 0.73* (ref 3)	0.7	5
313	0.87* (ref 3)	0.9	6
323	1.02 (ref 3)	1.0	7
333	1.18* (ref 3)	1.2	9
343	1.37* (ref 3)	1.4	10
353	1.60* (ref 3)	1.6	11
363	1.96* (ref 3)	2.0	14

^a Rounded values of ref 3, see text.

REFERENCES

1. Gilbert, E. C.; Lauer, B. E. *J. Phys. Chem.* 1927, *31*, 1050-2.
2. Richon, D.; Viillard, A. *Fluid Phase Equil.* 1985, *21*, 279-93.
3. Stephenson, R.; Stuart, J. *J. Chem. Eng. Data* 1986, *31*, 56-70.

<p>COMPONENTS:</p> <p>(1) Benzoic acid methyl ester (methyl benzoate); $C_8H_8O_2$; [93-58-3]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Gilbert, E.C.; Lauer, B.E. <i>J. Phys. Chem.</i> <u>1927</u>, 31, 1050-2.</p>																								
<p>VARIABLES:</p> <p>$T/K = 288 - 308$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>																								
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of benzoic acid methyl ester and water</p> <table border="1" data-bbox="239 520 1237 778"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr> <td>15</td> <td>0.1</td> <td>99.8</td> <td>0.00013</td> <td>0.985</td> </tr> <tr> <td>25</td> <td>0.4</td> <td>99.4</td> <td>0.00053</td> <td>0.956</td> </tr> <tr> <td>35</td> <td>0.5</td> <td>98.8</td> <td>0.00066</td> <td>0.916</td> </tr> </tbody> </table>		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	15	0.1	99.8	0.00013	0.985	25	0.4	99.4	0.00053	0.956	35	0.5	98.8	0.00066	0.916
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																						
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																					
15	0.1	99.8	0.00013	0.985																					
25	0.4	99.4	0.00053	0.956																					
35	0.5	98.8	0.00066	0.916																					
<p>AUXILIARY INFORMATION</p>																									
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The titration method was used as described by Walton and Jenkins (ref 1). A measured quantity of one component in a large test tube placed in a transparent constant temperature bath was titrated from a calibrated burette. The mixture was stirred rapidly with a motor stirrer and the equilibrium end point taken at the instant the solution became clear (with a lamp on the opposite side). The end points were sharp and easily duplicated. The temperature of the burette was recorded and from the density of the material the weight was computed. The solutions were carefully protected from moisture. The mutual solubility was determined from the densities of the respective saturated solutions on the assumption that mixing laws were valid in the range of low concentrations involved. Several runs were made at each temperature.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Eastman Kodak Company, chemically pure grade; redistilled under reduced pressure, a fraction boiling $88-91^\circ C$ at 17mm Hg being collected.</p> <p>(2) Distilled.</p> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <p>REFERENCES:</p> <p>1. Walton, J.H.; Jenkins, J.D. <i>J. Am. Chem. Soc.</i> <u>1923</u>, 45, 2555.</p>																								

COMPONENTS: (1) Benzoic acid methyl ester (methyl benzoate); $C_8H_8O_2$; [93-58-3] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Richon, D.; Viillard, A. <i>Fluid Phase Equilib.</i> <u>1985</u> , 21, 279-93.
VARIABLES: $T/K = 298$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: The solubility of benzoic acid methyl ester in water at 298.1 K was reported to be 0.00149 mol(1)/100 g(2). The corresponding mass per cent and mole fraction, x_1 , values calculated by the compiler are 0.202 g(1)/100g sln and 2.68×10^{-4} .	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The refractometric method was used. The Phoenix model 1-2000T differential refractometer from Texas Instruments was used and the solubility was determined from a characteristic calibration curve as described in the thesis of Richon (ref 1).	SOURCE AND PURITY OF MATERIALS: (1) Source not specified; purified by preparation gas chromatographic method; purity >99.5%, water content was negligible. (2) Distilled. ESTIMATED ERROR: Not specified. REFERENCES: 1. Richon, D. <i>Thesis</i> , University de Clermont-Ferrand, <u>1974</u> .

<p>COMPONENTS:</p> <p>(1) Benzoic acid methyl ester (methyl benzoate); $C_8H_8O_2$; [93-58-3]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u>, 31, 56-70.</p>																																																																
<p>VARIABLES:</p> <p>$T/K = 273 - 364$</p>	<p>PREPARED BY:</p> <p>Z. Maczynska</p>																																																																
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of benzoic acid methyl ester and water</p> <table border="1" data-bbox="243 526 1243 1036"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>0.221</td><td>99.54</td><td>0.000293</td><td>0.9662</td></tr> <tr><td>9.7</td><td>0.221</td><td>99.46</td><td>0.000293</td><td>0.9606</td></tr> <tr><td>20.1</td><td>0.213</td><td>99.35</td><td>0.000282</td><td>0.9529</td></tr> <tr><td>29.6</td><td>0.282</td><td>99.27</td><td>0.000374</td><td>0.9473</td></tr> <tr><td>40.2</td><td>0.247</td><td>99.16</td><td>0.000327</td><td>0.9398</td></tr> <tr><td>49.8</td><td>0.258</td><td>99.02</td><td>0.000342</td><td>0.9304</td></tr> <tr><td>60.1</td><td>0.286</td><td>98.76</td><td>0.000379</td><td>0.9133</td></tr> <tr><td>70.2</td><td>0.325</td><td>98.62</td><td>0.000431</td><td>0.9043</td></tr> <tr><td>80.3</td><td>0.358</td><td>98.44</td><td>0.000475</td><td>0.8930</td></tr> <tr><td>90.5</td><td>0.408</td><td>97.96</td><td>0.000542</td><td>0.8640</td></tr> <tr> <td>std. dev.</td> <td>0.002</td> <td>0.02</td> <td></td> <td></td> </tr> </tbody> </table>		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	0.221	99.54	0.000293	0.9662	9.7	0.221	99.46	0.000293	0.9606	20.1	0.213	99.35	0.000282	0.9529	29.6	0.282	99.27	0.000374	0.9473	40.2	0.247	99.16	0.000327	0.9398	49.8	0.258	99.02	0.000342	0.9304	60.1	0.286	98.76	0.000379	0.9133	70.2	0.325	98.62	0.000431	0.9043	80.3	0.358	98.44	0.000475	0.8930	90.5	0.408	97.96	0.000542	0.8640	std. dev.	0.002	0.02		
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																														
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																													
0	0.221	99.54	0.000293	0.9662																																																													
9.7	0.221	99.46	0.000293	0.9606																																																													
20.1	0.213	99.35	0.000282	0.9529																																																													
29.6	0.282	99.27	0.000374	0.9473																																																													
40.2	0.247	99.16	0.000327	0.9398																																																													
49.8	0.258	99.02	0.000342	0.9304																																																													
60.1	0.286	98.76	0.000379	0.9133																																																													
70.2	0.325	98.62	0.000431	0.9043																																																													
80.3	0.358	98.44	0.000475	0.8930																																																													
90.5	0.408	97.96	0.000542	0.8640																																																													
std. dev.	0.002	0.02																																																															
<p>AUXILIARY INFORMATION</p>																																																																	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, commercial sample; purity 99%; used as received.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Accuracy of method 0.1 wt% or less, for solubility, see above.</p> <p>REFERENCES:</p>																																																																

<p>COMPONENTS:</p> <p>(1) Benzoic acid, 2-hydroxy-, methyl ester (methyl salicylate); $C_8H_8O_3$; [119-36-8]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia March, 1990</p>
-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION:

Quantitative solubility data for the 2-hydroxybenzoic acid methyl ester (1) - water (2) system have been reported in the publications listed in Table 1.

TABLE 1: Quantitative Solubility Studies of the 2-Hydroxybenzoic acid methyl ester (1) - Water (2) System

Reference	T/K	Solubility	Method
Seidell (ref 1)	288-303	(1) in (2)	titration
Stephenson and Stuart (ref 2)	273-364	mutual	GLC

The original data in these publications are compiled in the Data Sheets immediately following this Critical Evaluation. For convenience, further discussion of this system will be divided into two parts.

1. SOLUBILITY OF 2-HYDROXYBENZOIC ACID METHYL ESTER (1) IN WATER (2)

All the available quantitative solubility data for the solubility of 2-hydroxybenzoic acid methyl ester (1) in water (2) are summarized in Table 2.

At 298 K the data of Seidell (ref 1) and Stephenson and Stuart (ref 2) are in excellent agreement. Furthermore Seidell reported the solubility of (1) in (2) was constant over the range 288-303 K which is consistent, within the limits of his likely experimental errors, with the more recent study of Stephenson and Stuart. This gives confidence in the reliability of the data of Stephenson and Stuart at other temperatures although, in the absence of confirmatory studies, their solubilities at temperatures other than 298 K must be regarded as Tentative.

(continued next page)

<p>COMPONENTS:</p> <p>(1) Benzoic acid, 2-hydroxy-, methyl ester (methyl salicylate); C₈H₈O₃; [119-36-8]</p> <p>(2) Water; H₂O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia March, 1990</p>
-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

TABLE 2: Recommended (R) and Tentative Solubilities of 2-Hydroxybenzoic acid methyl ester (1) in Water (2)

T/K	Solubilities		
	Reported values	"Best" values ^a	
	g(1)/100g sln	g(1)/100g sln	10 ⁴ x ₁
273	0.12* (ref 2)	0.12	1.4
283	0.12* (ref 2)	0.12	1.4
293	0.12* (ref 2)	0.12	1.4
298	0.12 (ref 1), 0.12* (ref 2)	0.12 (R)	1.4
303	0.13 (ref 2)	0.13	1.5
313	0.15 (ref 2)	0.15	1.8
323	0.17* (ref 2)	0.17	2.0
333	0.20* (ref 2)	0.20	2.4
343	0.22* (ref 2)	0.22	2.6
353	0.25* (ref 2)	0.25	2.9
363	0.27* (ref 2)	0.27	3.2

^a Values of ref 2, see text.

2. SOLUBILITY OF WATER (2) IN 2-HYDROXYBENZOIC ACID METHYL ESTER (1)

The only data available for the solubility of water (2) in 2-hydroxybenzoic acid methyl ester (1) are those of Stephenson and Stuart (ref 2) and so no Critical Evaluation is possible. The interested user is referred to the relevant Data Sheet for the original solubility values but it may be noted that the data of these authors are usually reliable.

REFERENCES

- Seidell, A. *Hygienic Lab. Bull.* 1910, No. 67, 98pp. (U.S. Govt. Printing Office, Washington, DC).
- Stephenson, R.; Stuart, J. *J. Chem. Eng. Data* 1986, 31, 56-70.

<p>COMPONENTS:</p> <p>(1) Benzoic acid, 2-hydroxy-, methyl ester (methyl salicylate); $C_8H_8O_3$; [119-36-8]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Seidell, A. <i>Hygienic Lab. Bull.</i> <u>1910</u>, No. 67, 98 pp (US Govt. Printing Office, Washington, DC).</p>
<p>VARIABLES:</p> <p>$T/K = 288 - 303$</p>	<p>PREPARED BY:</p> <p>G.T. Hefter</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of 2-hydroxybenzoic acid methyl ester in water at 25°C was reported to be 0.12 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 1.4×10^{-4}.</p> <p>The solubility of 2-hydroxybenzoic acid methyl ester at other temperatures (15-30°C) was reported to be constant with an approximate value of $0.1 \text{ cm}^3(1)/100\text{cm}^3(2)$.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The titration method was used. Portions (50 cm^3) of (2) were placed in an Erlenmeyer flask and titrated with (1) to opalescence. The flask and its contents were then cooled to the lowest temperature desired and allowed to warm slowly. The temperature was continuously recorded with a thermometer and the last drop or two of (1) was added as the desired temperature was reached.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not stated; purified by fractional distillation; b.p. 220-221°C, $d_{25}^{25} 1.182 \text{ g/cm}^3$.</p> <p>(2) Distilled (no details given).</p> <p>ESTIMATED ERROR:</p> <p>Temp. not stated. Soly. not stated.</p> <p>REFERENCES:</p>

<p>COMPONENTS:</p> <p>(1) Benzoic acid, 2-hydroxy-, methyl ester (methyl salicylate); $C_8H_8O_3$; [119-36-8]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u>, <i>31</i>, 56-70.</p>																																																																
<p>VARIABLES:</p> <p>$T/K = 273 - 364$</p>	<p>PREPARED BY:</p> <p>Z. Maczynska</p>																																																																
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of 2-hydroxybenzoic acid methyl ester and water</p> <table border="1" data-bbox="239 580 1243 1090"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>0.116</td><td>99.84</td><td>0.000137</td><td>0.9866</td></tr> <tr><td>9.5</td><td>0.131</td><td>99.83</td><td>0.000155</td><td>0.9858</td></tr> <tr><td>19.9</td><td>0.120</td><td>99.74</td><td>0.000142</td><td>0.9784</td></tr> <tr><td>29.7</td><td>0.124</td><td>99.70</td><td>0.000147</td><td>0.9752</td></tr> <tr><td>39.5</td><td>0.145</td><td>99.66</td><td>0.000172</td><td>0.9720</td></tr> <tr><td>50.0</td><td>0.175</td><td>99.59</td><td>0.000207</td><td>0.9664</td></tr> <tr><td>60.1</td><td>-</td><td>99.52</td><td>-</td><td>0.9608</td></tr> <tr><td>70.2</td><td>0.221</td><td>99.40</td><td>0.000262</td><td>0.9515</td></tr> <tr><td>80.1</td><td>0.270</td><td>99.25</td><td>0.000320</td><td>0.9400</td></tr> <tr><td>90.5</td><td>0.270</td><td>99.20</td><td>0.000320</td><td>0.9362</td></tr> <tr> <td>std. dev.</td> <td>0.004</td> <td>0.01</td> <td></td> <td></td> </tr> </tbody> </table>		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	0.116	99.84	0.000137	0.9866	9.5	0.131	99.83	0.000155	0.9858	19.9	0.120	99.74	0.000142	0.9784	29.7	0.124	99.70	0.000147	0.9752	39.5	0.145	99.66	0.000172	0.9720	50.0	0.175	99.59	0.000207	0.9664	60.1	-	99.52	-	0.9608	70.2	0.221	99.40	0.000262	0.9515	80.1	0.270	99.25	0.000320	0.9400	90.5	0.270	99.20	0.000320	0.9362	std. dev.	0.004	0.01		
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																														
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																													
0	0.116	99.84	0.000137	0.9866																																																													
9.5	0.131	99.83	0.000155	0.9858																																																													
19.9	0.120	99.74	0.000142	0.9784																																																													
29.7	0.124	99.70	0.000147	0.9752																																																													
39.5	0.145	99.66	0.000172	0.9720																																																													
50.0	0.175	99.59	0.000207	0.9664																																																													
60.1	-	99.52	-	0.9608																																																													
70.2	0.221	99.40	0.000262	0.9515																																																													
80.1	0.270	99.25	0.000320	0.9400																																																													
90.5	0.270	99.20	0.000320	0.9362																																																													
std. dev.	0.004	0.01																																																															
<p>AUXILIARY INFORMATION</p>																																																																	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, commercial sample; purity 99%; used as received.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Accuracy of method 0.1 wt% or less, for solubility, see above.</p> <p>REFERENCES:</p>																																																																

<p>COMPONENTS:</p> <p>(1) Acetic acid cyclohexyl ester (cyclohexyl acetate); $C_8H_{14}O_2$; [622-45-7]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988</p>
---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION:

Quantitative solubility data for the acetic acid cyclohexyl ester (1) - water (2) system have been reported in the publications listed in Table 1.

TABLE 1: Quantitative Solubility Studies of the
Acetic acid cyclohexyl ester (1) - Water (2) System

Reference	T/K	Solubility	Method
Doolittle (ref 1)	293	mutual	unspecified
Othmer et al. (ref 2)	297	mutual	unspecified
Stephenson and Stuart (ref 3)	273-363	mutual	GLC

The original data in these publications are compiled in the Data Sheets immediately following this Critical Evaluation. For convenience, further discussion of this system will be divided into two parts.

1. SOLUBILITY OF ACETIC ACID CYCLOHEXYL ESTER (1) IN WATER (2)

All the available data for the solubility of acetic acid cyclohexyl ester (1) in water (2) are summarized in Table 2 with the exception of the datum of Doolittle (ref 1) which is substantially lower than other data (ref 2,3) and has therefore been rejected.

At 296.7 K, the only temperature where comparison is possible, the data of Othmer et al. (ref 2) and Stephenson and Stuart (ref 3) are in good agreement and the average value is Recommended. At other temperatures only the data of Stephenson and Stuart (ref 3) are available and, pending further studies, must be regarded as Tentative, although it may be noted that solubilities reported by these authors are usually reliable.

(continued next page)

<p>COMPONENTS:</p> <p>(1) Acetic acid cyclohexyl ester (cyclohexyl acetate); $C_8H_{14}O_2$; [622-45-7]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988</p>
---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

TABLE 2: Recommended (R) and Tentative Solubilities
of Acetic acid cyclohexyl ester (1) in Water (2)

T/K	Solubilities		
	Reported values	"Best" values ^a	
	g(1)/100g sln	g(1)/100g sln	10^4x_1
273	0.393 (ref 3)	0.39	5.0
283	0.325* (ref 3)	0.33	4.2
293	0.292* (ref 3)	0.29	3.7
296.7	0.29 (ref 2), 0.282* (ref 3)	0.286 ± 0.006 (R) ^b	3.63^b
298	0.277* (ref 3)	0.28	3.6
303	0.264* (ref 3)	0.26	3.3
313	0.242* (ref 3)	0.24	3.0
323	0.225* (ref 3)	0.23	2.9
333	0.217* (ref 3)	0.22	2.8
343	0.224* (ref 3)	0.22	2.8
353	0.247* (ref 3)	0.25	3.2
363	0.290* (ref 3)	0.29	3.7

^a Rounded values of ref 3 except at 296.7 K.

^b Average value; x_1 has the same status and (relative) percentage uncertainty as the mass % solubility.

2. SOLUBILITY OF WATER (2) IN ACETIC ACID CYCLOHEXYL ESTER (1)

All the available solubility data for water (2) in acetic acid cyclohexyl ester (1) are summarized in Table 3. Serious disagreement exists between the values reported (at slightly different temperatures) by Doolittle (ref 1) and Othmer et al. (ref 2) on the one hand and Stephenson and Stuart (ref 3) on the other. In the absence of confirmatory studies it is not possible to distinguish between these data and thus no Critical Evaluation is possible, although it may be noted that the data of Stephenson and Stuart (ref 3) are usually reliable.

(continued next page)

<p>COMPONENTS:</p> <p>(1) Acetic acid cyclohexyl ester (cyclohexyl acetate); C₈H₁₄O₂; [622-45-7]</p> <p>(2) Water; H₂O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988</p>
---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

TABLE 3: Reported Solubilities
of Water (2) in Acetic acid cyclohexyl ester (1)

T/K	Solubilities	
	Reported values g(2)/100g sln	"Best" values ^a 10 ² x ₂ ^a
273	0.67 (ref 3)	5.1
283	0.65* (ref 3)	5.0
293	1.42 (ref 1), 0.70* (ref 3)	5.3
298	1.45 ^b (ref 2), 0.74* (ref 3)	5.6
303	0.80* (ref 3)	6.0
313	0.91* (ref 3)	6.8
323	1.00* (ref 3)	7.4
333	1.05* (ref 3)	7.7
343	1.07* (ref 3)	7.9
353	1.15* (ref 3)	8.4
363	1.36* (ref 3)	9.8

^a Rounded values of ref 3, see text.

^b 296.7 K.

REFERENCES

1. Doolittle, A. K. *Ind. Eng. Chem.* 1935, *27*, 1169-79.
2. Othmer, D. F.; White, R. E.; Trueger, E. *Ind. Eng. Chem.* 1941, *33*, 1240-8; *ibid.* 1513.
3. Stephenson, R.; Stuart, J. *J. Chem. Eng. Data* 1986, *31*, 56-70.

<p>COMPONENTS:</p> <p>(1) Acetic acid cyclohexyl ester (cyclohexyl acetate); $C_8H_{14}O_2$; [622-45-7]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Doolittle, A.K. <i>Ind. End. Chem.</i> <u>1935</u>, 27, 1169-79.</p>
<p>VARIABLES:</p> <p>$T/K = 293$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of acetic acid cyclohexyl ester in water at 20°C was reported to be 0.16 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 2.0×10^{-4}.</p> <p>The solubility of water in acetic acid cyclohexyl ester at 20°C was reported to be 1.42 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.102.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The method was not specified.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, commercial product; purity 79%, b.p. range 165-193°C, d_4^{20} 0.963.</p> <p>(2) Not specified.</p> <hr/> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <hr/> <p>REFERENCES:</p>

COMPONENTS: (1) Acetic acid cyclohexyl ester (cyclohexyl acetate); $C_8H_{14}O_2$; [622-45-7] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Othmer, D.F.; White, R.E.; Trueger, E. <i>Ind. Eng. Chem.</i> <u>1941</u> , 33, 1240-8, 1513.
VARIABLES: $T/K = 297$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: <p>The solubility of acetic acid cyclohexyl ester in water at 23.5°C was reported to be 0.29 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 3.7×10^{-4}.</p> <p>The solubility of water in acetic acid cyclohexyl ester at 23.5°C was reported to be 1.45 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.104.</p>	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: Nothing was specified in the paper.	SOURCE AND PURITY OF MATERIALS: (1) Not specified. (2) Not specified. ESTIMATED ERROR: Temp. $\pm 0.5^\circ C$ (mean of reported range). REFERENCES:

<p>COMPONENTS:</p> <p>(1) Acetic acid cyclohexyl ester (cyclohexyl acetate); $C_8H_{14}O_2$; [622-45-7]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u>, 31, 56-70.</p>																																																																
<p>VARIABLES:</p> <p>$T/K = 273 - 363$</p>	<p>PREPARED BY:</p> <p>Z. Maczynska</p>																																																																
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of acetic acid cyclohexyl ester and water</p> <table border="1" data-bbox="244 531 1239 1052"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>0.393</td><td>99.33</td><td>0.000499</td><td>0.9494</td></tr> <tr><td>9.4</td><td>0.317</td><td>99.35</td><td>0.000403</td><td>0.9509</td></tr> <tr><td>19.8</td><td>0.304</td><td>99.29</td><td>0.000386</td><td>0.9466</td></tr> <tr><td>29.7</td><td>0.260</td><td>99.21</td><td>0.000330</td><td>0.9408</td></tr> <tr><td>39.6</td><td>0.251</td><td>99.10</td><td>0.000319</td><td>0.9331</td></tr> <tr><td>49.8</td><td>0.229</td><td>99.01</td><td>0.000291</td><td>0.9268</td></tr> <tr><td>60.1</td><td>0.218</td><td>98.95</td><td>0.000277</td><td>0.9227</td></tr> <tr><td>70.0</td><td>0.224</td><td>98.93</td><td>0.000284</td><td>0.9213</td></tr> <tr><td>80.1</td><td>0.257</td><td>98.85</td><td>0.000326</td><td>0.9159</td></tr> <tr><td>90.3</td><td>0.286</td><td>98.64</td><td>0.000363</td><td>0.9018</td></tr> <tr> <td>std. dev.</td> <td>0.002</td> <td>0.01</td> <td></td> <td></td> </tr> </tbody> </table>		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	0.393	99.33	0.000499	0.9494	9.4	0.317	99.35	0.000403	0.9509	19.8	0.304	99.29	0.000386	0.9466	29.7	0.260	99.21	0.000330	0.9408	39.6	0.251	99.10	0.000319	0.9331	49.8	0.229	99.01	0.000291	0.9268	60.1	0.218	98.95	0.000277	0.9227	70.0	0.224	98.93	0.000284	0.9213	80.1	0.257	98.85	0.000326	0.9159	90.3	0.286	98.64	0.000363	0.9018	std. dev.	0.002	0.01		
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																														
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																													
0	0.393	99.33	0.000499	0.9494																																																													
9.4	0.317	99.35	0.000403	0.9509																																																													
19.8	0.304	99.29	0.000386	0.9466																																																													
29.7	0.260	99.21	0.000330	0.9408																																																													
39.6	0.251	99.10	0.000319	0.9331																																																													
49.8	0.229	99.01	0.000291	0.9268																																																													
60.1	0.218	98.95	0.000277	0.9227																																																													
70.0	0.224	98.93	0.000284	0.9213																																																													
80.1	0.257	98.85	0.000326	0.9159																																																													
90.3	0.286	98.64	0.000363	0.9018																																																													
std. dev.	0.002	0.01																																																															
<p>AUXILIARY INFORMATION</p>																																																																	
<p>METHOD/Apparatus/PROCEDURE:</p> <p>The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, commercial sample; purity 99%; used as received.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Accuracy of method 0.1 wt% or less, for solubility, see above.</p> <p>REFERENCES:</p>																																																																

<p>COMPONENTS:</p> <p>(1) Butanedioic acid diethyl ester (diethyl succinate); $C_8H_{14}O_4$; [123-25-1]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988</p>
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION:

Quantitative solubility data for the butanedioic acid diethyl ester (1) - water (2) system have been reported in the publications listed in Table 1.

TABLE 1: Quantitative Solubility Studies of the
Butanedioic acid diethyl ester (1) - Water (2) System

Reference	T/K	Solubility	Method
Sobotka and Kahn (ref 1)	293	(1) in (2)	titration
Stephenson and Stuart (ref 2)	274-364	mutual	GLC

The original data in these publications are compiled in the Data Sheets immediately following this Critical Evaluation. For convenience, further discussion of this system will be divided into two parts.

1. SOLUBILITY OF BUTANEDIOIC ACID DIETHYL ESTER (1) IN WATER (2)

All the available data for the solubility of butanedioic acid diethyl ester (1) in water (2) are summarized in Table 2. At 293 K, the only temperature where comparison is possible, the data of Sobotka and Kahn (ref 1) and Stephenson and Stuart (ref 2) are in good agreement and the average value is Recommended. At other temperatures only the data of Stephenson and Stuart (ref 2) are available and must therefore be regarded as Tentative although it may be noted that the solubilities reported by these authors are generally reliable.

(continued next page)

COMPONENTS: (1) Butanedioic acid diethyl ester (diethyl succinate); $C_8H_{14}O_4$; [123-25-1] (2) Water; H_2O ; [7732-18-5]	EVALUATOR: G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988
----------------------------------------------------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

TABLE 2: Recommended (R) and Tentative Solubilities
of Butanedioic acid diethyl ester (1) in Water (2)

T/K	Solubilities		
	Reported values	"Best" values ^a	
	g(1)/100g sln	g(1)/100g sln	10^3x_1
273	2.86 (ref 2)	2.9	3.0
283	2.50* (ref 2)	2.5	2.6
293	1.89 (ref 1), 2.17* (ref 2)	2.03 ± 0.14 (R) ^b	2.14^b
298	2.15* (ref 2)	2.2	2.3
303	1.95* (ref 2)	2.0	2.1
313	1.81* (ref 2)	1.8	1.9
323	1.75* (ref 2)	1.8	1.9
333	1.75* (ref 2)	1.8	1.9
343	1.78* (ref 2)	1.8	1.9
353	1.86* (ref 2)	1.9	2.0
363	2.05* (ref 2)	2.1	2.2

^a Rounded values of ref 2 except at 293 K.

^b Average value; x_1 has the same status and (relative) percentage uncertainty as the mass % solubility.

2. SOLUBILITY OF WATER (2) IN BUTANEDIOIC ACID DIETHYL ESTER (1)

The only data available for the solubility of water (2) in butanedioic acid diethyl ester (1) are those of Stephenson and Stuart (ref 2) and so no Critical Evaluation is possible. The interested user is referred to the relevant Data Sheet for the experimental solubilities but it may be noted that the data of these authors are usually reliable.

REFERENCES

- Sobotka, H.; Kahn, J. *J. Am. Chem. Soc.* **1931**, *53*, 2935-8.
- Stephenson, R.; Stuart, J. *J. Chem. Eng. Data* **1986**, *31*, 56-70.

COMPONENTS: (1) Butanedioic acid diethyl ester (diethyl succinate); $C_8H_{14}O_4$; [123-25-1] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Sobotka, H.; Kahn, J. <i>J. Am. Chem. Soc.</i> <u>1931</u> , 53, 2935-8.
VARIABLES: $T/K = 293$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: The solubility of butanedioic acid diethyl ester in water at 20°C was reported to be 1.92 g(1)/100mL(2). The corresponding mass per cent and mole fraction, x_1 , values calculated by the compiler are 1.89 g(1)/100g sln and 0.00199. Density of water $d_4^{20} = 0.9982$ (ref 1) was used in the calculation.	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The titration method was used. The ester was added dropwise from a micro-burette with a capillary tip to 100, 250, or 500 mL of water in a narrow-mouthed stock bottle with a well-ground glass stopper. The bottle was shaken after each addition of ester. 1-5 mg of Sudan IV dye was put into the water to improve the end-point of the titration. At saturation, one additional drop of ester was sufficient to convert the floating rough indicator particles into dark transparent droplets.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified (Eastman Kodak Laboratories or synthesized); twice-distilled under reduced pressure; d_4^{20} 1.0402, n_D^{20} 1.4200. (2) Distilled. ESTIMATED ERROR: Not specified. REFERENCES: 1. <i>Selected Values of Properties of Hydrocarbons and Related Compounds</i> , API Research Project 44, Thermodynamics Research Center, Texas A and M University, Texas, <u>1973</u> .

<p>COMPONENTS:</p> <p>(1) Butanedioic acid diethyl ester (diethyl succinate); $C_8H_{14}O_4$; [123-25-1]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u>, 31, 56-70.</p>																																																											
<p>VARIABLES:</p> <p>$T/K = 273 - 364$</p>	<p>PREPARED BY:</p> <p>Z. Maczynska</p>																																																											
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of butanedioic acid diethyl ester and water</p> <table border="1" data-bbox="235 520 1233 1030"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>2.86</td><td>98.72</td><td>0.00303</td><td>0.8886</td></tr> <tr><td>10.0</td><td>2.67</td><td>98.49</td><td>0.00283</td><td>0.8709</td></tr> <tr><td>20.0</td><td>2.13</td><td>98.22</td><td>0.00225</td><td>0.8509</td></tr> <tr><td>29.7</td><td>1.99</td><td>97.92</td><td>0.00209</td><td>0.8296</td></tr> <tr><td>39.8</td><td>1.79</td><td>97.58</td><td>0.00188</td><td>0.8066</td></tr> <tr><td>50.0</td><td>1.79</td><td>97.13</td><td>0.00188</td><td>0.7778</td></tr> <tr><td>60.3</td><td>1.74</td><td>96.76</td><td>0.00183</td><td>0.7554</td></tr> <tr><td>70.0</td><td>1.79</td><td>96.40</td><td>0.00188</td><td>0.7347</td></tr> <tr><td>80.1</td><td>1.84</td><td>96.03</td><td>0.00193</td><td>0.7144</td></tr> <tr><td>90.7</td><td>2.09</td><td>95.77</td><td>0.00220</td><td>0.7007</td></tr> </tbody> </table> <p>std. dev. 0.05 0.07</p>		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	2.86	98.72	0.00303	0.8886	10.0	2.67	98.49	0.00283	0.8709	20.0	2.13	98.22	0.00225	0.8509	29.7	1.99	97.92	0.00209	0.8296	39.8	1.79	97.58	0.00188	0.8066	50.0	1.79	97.13	0.00188	0.7778	60.3	1.74	96.76	0.00183	0.7554	70.0	1.79	96.40	0.00188	0.7347	80.1	1.84	96.03	0.00193	0.7144	90.7	2.09	95.77	0.00220	0.7007
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																									
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																								
0	2.86	98.72	0.00303	0.8886																																																								
10.0	2.67	98.49	0.00283	0.8709																																																								
20.0	2.13	98.22	0.00225	0.8509																																																								
29.7	1.99	97.92	0.00209	0.8296																																																								
39.8	1.79	97.58	0.00188	0.8066																																																								
50.0	1.79	97.13	0.00188	0.7778																																																								
60.3	1.74	96.76	0.00183	0.7554																																																								
70.0	1.79	96.40	0.00188	0.7347																																																								
80.1	1.84	96.03	0.00193	0.7144																																																								
90.7	2.09	95.77	0.00220	0.7007																																																								
<p>AUXILIARY INFORMATION</p>																																																												
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, commercial sample; purity 99%; used as received.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Accuracy of method 0.1 wt% or less, for solubility, see above.</p> <p>REFERENCES:</p>																																																											

<p>COMPONENTS:</p> <p>(1) 2,3-Butanediol diacetate (2,3-butylene diacetate); $C_8H_{14}O_4$; [1114-92-7]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Othmer, D.F.; Bergen, W.S.; Shlechter, N.; Bruins, P.F.</p> <p><i>Ind. Eng. Chem.</i> <u>1945</u>, 37, 890-4.</p>																								
<p>VARIABLES:</p> <p>$T/K = 299 - 348$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>																								
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of 2,3-Butanediol diacetate and water</p> <table border="1" data-bbox="137 512 1131 762"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr> <td>26</td> <td>3.62</td> <td>96.3</td> <td>0.00387</td> <td>0.729</td> </tr> <tr> <td>50</td> <td>2.87</td> <td>96.6</td> <td>0.00305</td> <td>0.746</td> </tr> <tr> <td>75</td> <td>3.7</td> <td>95.9</td> <td>0.00396</td> <td>0.708</td> </tr> </tbody> </table>		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	26	3.62	96.3	0.00387	0.729	50	2.87	96.6	0.00305	0.746	75	3.7	95.9	0.00396	0.708
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																						
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																					
26	3.62	96.3	0.00387	0.729																					
50	2.87	96.6	0.00305	0.746																					
75	3.7	95.9	0.00396	0.708																					
<p>AUXILIARY INFORMATION</p>																									
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>Presumably the titration method reported by Othmer, White and Trueger (ref 1) was used. The data were reported together with the ternary system 2,3-Butanediol diacetate-water-2,3-butanediol (2,3-butylene diacetate-water-butylene glycol).</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Synthesized from meso-2,3-butylene glycol; b.p. range $1^\circ C$.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <p>REFERENCES:</p> <p>1. Othmer, D.F.; White, R.E.; Trueger, E. <i>Ind. Eng. Chem.</i> <u>1941</u>, 33, 1240.</p>																								

<p>COMPONENTS:</p> <p>(1) Hexanedioic acid dimethyl ester (<i>dimethyl adipate</i>); $C_8H_{14}O_4$; [627-93-0]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u>, 31, 56-70.</p>																																																											
<p>VARIABLES:</p> <p>$T/K = 282 - 364$</p>	<p>PREPARED BY:</p> <p>Z. Maczynska</p>																																																											
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of hexanedioic acid dimethyl ester and water</p> <table border="1" data-bbox="246 532 1240 1056"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr> <td>9.2</td> <td>3.81</td> <td>98.06</td> <td>0.00408</td> <td>0.8394</td> </tr> <tr> <td>19.6</td> <td>3.07</td> <td>97.70</td> <td>0.00326</td> <td>0.8146</td> </tr> <tr> <td>30.9</td> <td>2.85</td> <td>97.33</td> <td>0.00302</td> <td>0.7903</td> </tr> <tr> <td>39.5</td> <td>2.77</td> <td>96.93</td> <td>0.00294</td> <td>0.7655</td> </tr> <tr> <td>49.8</td> <td>2.70</td> <td>96.34</td> <td>0.00286</td> <td>0.7313</td> </tr> <tr> <td>60.0</td> <td>2.73</td> <td>95.90</td> <td>0.00289</td> <td>0.7075</td> </tr> <tr> <td>70.5</td> <td>2.87</td> <td>95.27</td> <td>0.00305</td> <td>0.6756</td> </tr> <tr> <td>80.2</td> <td>3.50</td> <td>94.59</td> <td>0.00374</td> <td>0.6439</td> </tr> <tr> <td>90.5</td> <td>3.23</td> <td>94.28</td> <td>0.00344</td> <td>0.6302</td> </tr> <tr> <td>std. dev.</td> <td>0.06</td> <td>0.06</td> <td></td> <td></td> </tr> </tbody> </table>		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	9.2	3.81	98.06	0.00408	0.8394	19.6	3.07	97.70	0.00326	0.8146	30.9	2.85	97.33	0.00302	0.7903	39.5	2.77	96.93	0.00294	0.7655	49.8	2.70	96.34	0.00286	0.7313	60.0	2.73	95.90	0.00289	0.7075	70.5	2.87	95.27	0.00305	0.6756	80.2	3.50	94.59	0.00374	0.6439	90.5	3.23	94.28	0.00344	0.6302	std. dev.	0.06	0.06		
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																									
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																								
9.2	3.81	98.06	0.00408	0.8394																																																								
19.6	3.07	97.70	0.00326	0.8146																																																								
30.9	2.85	97.33	0.00302	0.7903																																																								
39.5	2.77	96.93	0.00294	0.7655																																																								
49.8	2.70	96.34	0.00286	0.7313																																																								
60.0	2.73	95.90	0.00289	0.7075																																																								
70.5	2.87	95.27	0.00305	0.6756																																																								
80.2	3.50	94.59	0.00374	0.6439																																																								
90.5	3.23	94.28	0.00344	0.6302																																																								
std. dev.	0.06	0.06																																																										
<p>AUXILIARY INFORMATION</p>																																																												
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, commercial sample; purity 98%; used as received.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Accuracy of method 0.1 wt% or less, for solubility, see above.</p> <p>REFERENCES:</p>																																																											

COMPONENTS: (1) Acetic acid 1,3-dimethylbutyl ester (1,3-dimethylbutyl acetate); $C_8H_{16}O_2$; [108-84-9] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Doolittle, A.K. <i>Ind. End. Chem.</i> <u>1935</u> , 27, 1169-79.
VARIABLES: $T/K = 293$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: <p>The solubility of acetic acid 1,3-dimethylbutyl ester in water at 20°C was reported to be 0.08 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 1.0×10^{-4}.</p> <p>The solubility of water in acetic acid 1,3-dimethylbutyl ester at 20°C was reported to be 0.89 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.067.</p>	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The method was not specified.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial product; purity 95%, b.p. range 140-147°C, d_4^{20} 0.857, n_D^{20} 1.4008. (2) Not specified.
ESTIMATED ERROR: Not specified.	
REFERENCES:	

<p>COMPONENTS:</p> <p>(1) Acetic acid 2-ethyl-1-butyl ester (2-ethylbutyl acetate); $C_8H_{16}O_2$; [10031-87-5]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Doolittle, A.K.</p> <p><i>Ind. End. Chem.</i> <u>1935</u>, 27, 1169-79.</p>
<p>VARIABLES:</p> <p>$T/K = 293$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of acetic acid 2-ethyl-1-butyl ester in water at 20°C was reported to be 0.06 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 7×10^{-5}.</p> <p>The solubility of water in acetic acid 2-ethyl-1-butyl ester at 20°C was reported to be 0.75 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.057.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The method was not specified.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, commercial product; purity 91%, b.p. range 157-164°C, d_4^{20} 0.876, n_D^{20} 1.4103.</p> <p>(2) Not specified.</p> <hr/> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <hr/> <p>REFERENCES:</p>

<p>COMPONENTS:</p> <p>(1) Acetic acid hexyl ester (hexyl acetate); $C_8H_{16}O_2$; [142-92-7]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988</p>
-----------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION:

Quantitative solubility data for the acetic acid hexyl ester (1) - water (2) system have been reported in the publications listed in Table 1.

TABLE 1: Quantitative Solubility Studies of the
Acetic acid hexyl ester (1) - Water (2) System

Reference	T/K	Solubility	Method
Massaldi and King (ref 1)	298	(1) in (2)	GLC
Alvarez and Neila (ref 2)	298	mutual	titration
Stephenson and Stuart (ref 3)	273-363	mutual	GLC

The original data in these publications are compiled in the Data Sheets immediately following this Critical Evaluation. For convenience, further discussion of this system will be divided into two parts.

1. SOLUBILITY OF ACETIC ACID HEXYL ESTER (1) IN WATER (2)

All the available data for the solubility of acetic acid hexyl ester (1) in water (2) are summarized in Table 2 with the exception of the approximate value of Alvarez and Neila (ref 2) which is much higher than the other studies (ref 1,3) and is therefore rejected. At 298 K, the only temperature where comparison is possible, the data of Massaldi and King (ref 1) and Stephenson and Stuart (ref 3) are in excellent agreement and the average value is Recommended. At other temperatures only the values of Stephenson and Stuart (ref 3) are available. Although their results for this system are rather scattered, the solubilities reported by these authors are generally reliable and pending further studies can be considered as Tentative.

(continued next page)

<p>COMPONENTS:</p> <p>(1) Acetic acid hexyl ester (<i>hexyl acetate</i>); $C_8H_{16}O_2$; [142-92-7]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988</p>
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

TABLE 2: Recommended (R) and Tentative Solubilities
of Acetic acid hexyl ester (1) in Water (2)

T/K	Solubilities		
	Reported values	"Best" values ^a	
	g(1)/100g sln	g(1)/100g sln	$10^5 x_1$
273	0.084 (ref 3)	0.08	10
283	0.066 (ref 3)	0.07	9
293	0.056* (ref 3)	0.06	7
298	0.051 (ref 1), 0.053 (ref 3)	0.052 ± 0.001 (R) ^b	6.5
303	0.052* (ref 3)	0.05	6
313	0.051* (ref 3)	0.05	6
323	0.051* (ref 3)	0.05	6
333	0.052* (ref 3)	0.05	6
343	0.056* (ref 3)	0.06	7
353	0.060* (ref 3)	0.06	7
363	0.066* (ref 3)	0.07	9

^a Rounded values of ref 3 except at 298 K.

^b Average value; x_1 has same status and (relative) percentage uncertainty as the mass % solubility.

2. SOLUBILITY OF WATER (2) IN ACETIC ACID HEXYL ESTER (1)

All the available data for the solubility of water (2) in acetic acid hexyl ester (1) are listed in Table 3. At 298 K, the only temperature where comparison is possible, the approximate value of Alvarez and Neila (ref 2) is not in very good agreement with the interpolated datum of Stephenson and Stuart (ref 3). Consequently, in the absence of other independent studies, all the data must be regarded as very Tentative. However, it may be noted that the solubilities reported by Stephenson and Stuart (ref 3) are generally reliable although the present values are rather scattered.

(continued next page)

<p>COMPONENTS:</p> <p>(1) Acetic acid hexyl ester (hexyl acetate); $C_8H_{16}O_2$; [142-92-7]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988</p>
-----------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

TABLE 3: Tentative Solubilities
of Water (2) in Acetic acid hexyl ester (1)

T/K	Solubilities		
	Reported values	"Best" values ^a	
	g(2)/100g sln	g(2)/100g sln	10^2x_2
273	0.494 (ref 3)	0.5	4
283	0.54* (ref 3)	0.5	4
293	0.58* (ref 3)	0.6	5
298	1.1 ^b (ref 2), 0.61* (ref 3)	0.6	5
303	0.63* (ref 3)	0.6	5
313	0.66* (ref 3)	0.7	5
323	0.69* (ref 3)	0.7	5
333	0.72* (ref 3)	0.7	5
343	0.74* (ref 3)	0.7	5
353	0.76* (ref 3)	0.8	6
363	0.78* (ref 3)	0.8	6

^a Rounded values of ref 3, see text.

^b Not included in "Best" value.

REFERENCES

1. Massaldi, H. A.; King, C. J. *J. Chem. Eng. Data* 1973, 18, 393-7.
2. Alvarez, J. R.; Neila, J. J. *An. Quim.* 1978, 74, 326-32.
3. Stephenson, R.; Stuart, J. *J. Chem. Eng. Data* 1986, 31, 56-70.

<p>COMPONENTS:</p> <p>(1) Acetic acid hexyl ester (hexyl acetate); $C_8H_{16}O_2$; [142-92-7]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Massaldi, H.A.; King, C.J. <i>J. Chem. Eng. Data</i> <u>1973</u>, 18, 393-7.</p>
<p>VARIABLES:</p> <p>$T/K = 298$</p>	<p>PREPARED BY:</p> <p>Z. Maczynska</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of acetic acid hexyl ester in water at 25°C was reported to be 0.00354 mol(1)/L sln.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method was used. A technique based on head-space analysis using gas-liquid chromatography was developed to determine solubilities of sparingly soluble organics. Saturated solutions need not be prepared in advance; thus, phase separation problems were avoided. Furthermore, the analysis of liquid samples was not required. This method is versatile enough to give determinations provided that the vapor pressure of the pure liquid substance is known. The gas chromatograph was a Varian Aero-graph Model 1740 with a flame ionization detector.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Eastman Kodak Co., reagent grade; purity not specified; used as received.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm 0.05^\circ C$.</p> <p>REFERENCES:</p>

COMPONENTS: (1) Acetic acid hexyl ester (hexyl acetate); $C_8H_{16}O_2$; [142-92-7] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Alvarez, J.R.; Neila, J.J. An. Quim. <u>1978</u> , 74, 326-32.
VARIABLES: $T/K = 298$	PREPARED BY: Z. Maczynska
EXPERIMENTAL VALUES: <p>The solubility of acetic acid hexyl ester in water at 25°C was reported to be 0.1 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 0.00013.</p> <p>The solubility of water in acetic acid hexyl ester at 25°C was reported to be 1.1 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.082.</p>	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: Presumably the titration method was used. The data were reported together with the ternary system acetic acid hexyl ester-water-phenol. No further details were reported in the paper.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial product; used as received; $d_4^{21.3}$ 0.8681, n_D^{20} 1.4093. (2) Twice distilled over $KMnO_4$.
	ESTIMATED ERROR: Not specified.
	REFERENCES:

COMPONENTS: (1) Acetic acid hexyl ester (hexyl acetate); $C_8H_{16}O_2$; [142-92-7] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u> , <i>31</i> , 56-70.																																																																
VARIABLES: $T/K = 273 - 363$	PREPARED BY: Z. Maczynska																																																																
EXPERIMENTAL VALUES: Mutual solubility of acetic acid hexyl ester and water <table border="1" data-bbox="253 520 1251 1030"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">$g(1)/100g\ sln$</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>0.084</td><td>99.506</td><td>0.000105</td><td>0.96177</td></tr> <tr><td>9.5</td><td>0.070</td><td>99.504</td><td>0.000087</td><td>0.96162</td></tr> <tr><td>19.7</td><td>0.043</td><td>99.434</td><td>0.000054</td><td>0.95641</td></tr> <tr><td>29.6</td><td>0.054</td><td>99.270</td><td>0.000067</td><td>0.94441</td></tr> <tr><td>40.0</td><td>0.056</td><td>99.254</td><td>0.000070</td><td>0.94325</td></tr> <tr><td>50.0</td><td>0.047</td><td>99.367</td><td>0.000059</td><td>0.95148</td></tr> <tr><td>60.7</td><td>0.051</td><td>99.305</td><td>0.000064</td><td>0.94695</td></tr> <tr><td>70.0</td><td>0.057</td><td>99.251</td><td>0.000071</td><td>0.94303</td></tr> <tr><td>80.0</td><td>0.063</td><td>99.220</td><td>0.000079</td><td>0.94079</td></tr> <tr><td>90.2</td><td>0.066</td><td>99.215</td><td>0.000082</td><td>0.94044</td></tr> <tr> <td>std. dev.</td> <td>0.003</td> <td>0.004</td> <td></td> <td></td> </tr> </tbody> </table>		$t/^\circ C$	$g(1)/100g\ sln$		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	0.084	99.506	0.000105	0.96177	9.5	0.070	99.504	0.000087	0.96162	19.7	0.043	99.434	0.000054	0.95641	29.6	0.054	99.270	0.000067	0.94441	40.0	0.056	99.254	0.000070	0.94325	50.0	0.047	99.367	0.000059	0.95148	60.7	0.051	99.305	0.000064	0.94695	70.0	0.057	99.251	0.000071	0.94303	80.0	0.063	99.220	0.000079	0.94079	90.2	0.066	99.215	0.000082	0.94044	std. dev.	0.003	0.004		
$t/^\circ C$	$g(1)/100g\ sln$		x_1 (compiler)																																																														
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																													
0	0.084	99.506	0.000105	0.96177																																																													
9.5	0.070	99.504	0.000087	0.96162																																																													
19.7	0.043	99.434	0.000054	0.95641																																																													
29.6	0.054	99.270	0.000067	0.94441																																																													
40.0	0.056	99.254	0.000070	0.94325																																																													
50.0	0.047	99.367	0.000059	0.95148																																																													
60.7	0.051	99.305	0.000064	0.94695																																																													
70.0	0.057	99.251	0.000071	0.94303																																																													
80.0	0.063	99.220	0.000079	0.94079																																																													
90.2	0.066	99.215	0.000082	0.94044																																																													
std. dev.	0.003	0.004																																																															
AUXILIARY INFORMATION																																																																	
METHOD/APPARATUS/PROCEDURE: The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial sample; purity 99%; used as received. (2) Not specified. ESTIMATED ERROR: Accuracy of method 0.1 wt% or less, for solubility, see above. REFERENCES:																																																																

<p>COMPONENTS:</p> <p>(1) Acetic acid 1-methylpentyl ester (<i>1-methylpentyl acetate</i>); $C_8H_{16}O_2$; [5953-49-1]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia</p> <p>December, 1988</p>
-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION:

Quantitative solubility data for the acetic acid 1-methylpentyl ester (1) - water (2) system have been reported in the publications listed in Table 1.

TABLE 1: Quantitative Solubility Studies of the Acetic acid 1-methylpentyl ester (1) - Water (2) System

Reference	T/K	Solubility	Method
Park and Hopkins (ref 1)	298	(1) in (2)	unspecified
Doolittle (ref 2)	293	mutual	unspecified

The original data in these publications are compiled in the Data Sheets immediately following this Critical Evaluation.

Unfortunately, the data for the solubility of acetic acid 1-methylpentyl ester (1) in water (2) in both the publications listed in Table 1 are only semiquantitative. According to Park and Hopkins (ref 1) the solubility of (1) in (2) is less than 0.14 g(1)/100 g sln. Doolittle (ref 2) merely indicates (1) is "very slightly soluble" in (2).

For the solubility of water (2) in 1-methylpentyl acetate (1) only the datum of Doolittle (ref 2), 0.68 g(2)/100 g sln, is available and so no Critical Evaluation is possible. This system clearly requires further investigation.

REFERENCES

1. Park, J. G.; Hopkins, M. B. *Ind. Eng. Chem.* 1930, *22*, 826-30.
2. Doolittle, A. K. *Ind. Eng. Chem.* 1935, *27*, 1169-79.

<p>COMPONENTS:</p> <p>(1) Acetic acid 1-methylpentyl ester (1-methylpentyl acetate); $C_8H_{16}O_2$; [5953-49-1]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Park, J.G.; Hopkins, M.B.</p> <p>Ind. Eng. Chem. <u>1930</u>, 22, 826-30.</p>
<p>VARIABLES:</p> <p>$T/K = 298$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of acetic acid 1-methylpentyl ester in water at 25°C was reported to be less than 0.1 mL(1)/100mL(2).</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The method was not specified.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, commercial samples; used as received; 85-88% of ester, b.p. range 146-156°C, n_D^{20} 1.4081.</p> <p>(2) Not specified.</p> <hr/> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <hr/> <p>REFERENCES:</p>

<p>COMPONENTS:</p> <p>(1) Acetic acid 1-methylpentyl ester (<i>1-methylpentyl acetate</i>); $C_8H_{16}O_2$; [5953-49-1]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Doolittle, A.K.</p> <p><i>Ind. End. Chem.</i> <u>1935</u>, 27, 1169-79.</p>
<p>VARIABLES:</p> <p>$T/K = 293$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of acetic acid 1-methylpentyl ester in water at 20°C was reported to be very slightly soluble.</p> <p>The solubility of water in acetic acid 1-methylpentyl ester at 20°C was reported to be 0.68 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.052.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The method was not specified.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, commercial product; purity 98%, b.p. range 129-158°C, d_4^{20} 0.861.</p> <p>(2) Not specified.</p> <hr/> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <hr/> <p>REFERENCES:</p>

<p>COMPONENTS:</p> <p>(1) Butanoic acid butyl ester (<i>butyl butyrate</i>); $C_8H_{16}O_2$; [109-21-7]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988</p>
---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION:

Quantitative solubility data for the butanoic acid butyl ester (1) - water (2) system have been reported in the publications listed in Table 1.

TABLE 1: Quantitative Solubility Studies of the
Butanoic acid butyl ester (1) - Water (2) System

Reference	T/K	Solubility	Method
Bridgman (ref 1)	283-306	(2) in (1)	synthetic
Doolittle (ref 2)	293	mutual	unspecified
Chebotaev <i>et al.</i> (ref 3)	370	(2) in (1)	analytical
Stephenson and Stuart (ref 4)	273-364	mutual	GLC

The original data in these publications are compiled in the Data Sheets immediately following this Critical Evaluation. For convenience, further discussion of this system will be divided into two parts.

1. SOLUBILITY OF BUTANOIC ACID BUTYL ESTER (1) IN WATER (2)

All the available data for the solubility of butanoic acid butyl ester (1) in water (2) are summarized in Table 2. At 293 K, the only temperature where comparison is possible, the data of Doolittle (ref 2) and Stephenson and Stuart (ref 4) are in serious disagreement. In the absence of independent studies it is impossible to prefer either value although it may be noted that the data of Stephenson and Stuart (ref 4) are generally reliable.

(continued next page)

<p>COMPONENTS:</p> <p>(1) Butanoic acid butyl ester (butyl butyrate); $C_8H_{16}O_2$; [109-21-7]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988</p>
--------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

TABLE 2: Reported Solubilities
of Butanoic acid butyl ester (1) in Water (2)

T/K	Reported Solubilities	
	g(1)/100g sln	$10^4 x_1^a$
273	1.22 (ref 4)	15.4
283	0.90* (ref 4)	11.3
293	0.05 (ref 2), 0.66* (ref 4)	8.3
298	0.60* (ref 4)	7.5
303	0.56* (ref 4)	7.0
313	0.52* (ref 4)	6.5
323	0.49* (ref 4)	6.1
333	0.47* (ref 4)	5.9
343	0.47* (ref 4)	5.9
353	0.49* (ref 4)	6.1
363	0.51* (ref 4)	6.3

^a Data of ref 4, see text.

2. SOLUBILITY OF WATER (2) IN BUTANOIC ACID BUTYL ESTER (1)

All the available data for the solubility of water (2) in butanoic acid butyl ester (1) are summarized in Table 3 with the exception of the value of Chebotaev et al. (ref 3) which is much higher than the other (ref 4) study and is rejected. Unlike the H_2O -rich phase the agreement between independent studies over the range 283-303 is good and a number of values have been Recommended. At other temperatures only the data of Stephenson and Stuart (ref 4) are available, excluding the datum of Chebotaev et al. (ref 3) referred to above, and must be considered Tentative pending further studies.

(continued next page)

<p>COMPONENTS:</p> <p>(1) Butanoic acid butyl ester (butyl butyrate); $C_8H_{16}O_2$; [109-21-7]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988</p>
--------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

TABLE 3: Recommended (R) and Tentative Solubilities
of Water (2) in Butanoic acid butyl ester (1)

T/K	Solubilities		
	Reported values	"Best" values ($\pm\sigma_n$) ^a	
	g(2)/100g sln	g(2)/100g sln	10^2x_2
273	0.339 (ref 4)	0.34	2.7
283	0.379 (ref 1), 0.370* (ref 4)	0.37 ± 0.01 (R)	2.9
293	0.450* (ref 1), 0.43 (ref 2), 0.412* (ref 4)	0.43 ± 0.02 (R)	3.3
298	0.490* (ref 1), 0.438* (ref 4)	0.46 ± 0.03 (R)	3.5
303	0.533* (ref 1), 0.466 (ref 4)	0.50 ± 0.03 (R)	3.9
313	0.53* (ref 4)	0.53	4.1
323	0.59* (ref 4)	0.59	4.5
333	0.63* (ref 4)	0.63	4.8
343	0.66* (ref 4)	0.66	5.1
353	0.67* (ref 4)	0.67	5.1
363	0.68* (ref 4)	0.68	5.2

^a Obtained by averaging where appropriate; σ_n has no statistical significance. Mole fraction solubilities (x_2) have the same status and (relative) percentage uncertainties as the mass % solubilities.

REFERENCES

1. Bridgman, J. A. *Ind. Eng. Chem.* 1928, 20, 184-7.
2. Doolittle, A. K. *Ind. Eng. Chem.* 1935, 27, 1169-79.
3. Chebotaev, V. F.; Balashov, M. I.; Serafimov, L. A. *Fiz. Khim. Osn. Rektifikatsii* 1970, 100-3.
4. Stephenson, R.; Stuart, J. *J. Chem. Eng. Data* 1986, 31, 56-70.

<p>COMPONENTS:</p> <p>(1) Butanoic acid butyl ester (butyl butyrate); $C_8H_{16}O_2$; [109-21-7]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Bridgman, J.A. <i>Ind. End. Chem.</i> <u>1928</u>, 20, 184-7.</p>																				
<p>VARIABLES:</p> <p>$T/K = 283 - 306$</p>	<p>PREPARED BY:</p> <p>Z. Maczynska</p>																				
<p>EXPERIMENTAL VALUES:</p> <p style="text-align: center;">Solubility of water in butanoic acid butyl ester</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: center;">$t/^\circ C$</th> <th style="text-align: center;">g(2)/100g(1)</th> <th style="text-align: center;">g(2)/100g solution (compiler)</th> <th style="text-align: center;">x_2 (compiler)</th> </tr> </thead> <tbody> <tr> <td style="text-align: center;">10.0</td> <td style="text-align: center;">0.380</td> <td style="text-align: center;">0.379</td> <td style="text-align: center;">0.0296</td> </tr> <tr> <td style="text-align: center;">16.0</td> <td style="text-align: center;">0.424</td> <td style="text-align: center;">0.422</td> <td style="text-align: center;">0.0328</td> </tr> <tr> <td style="text-align: center;">25.5</td> <td style="text-align: center;">0.497</td> <td style="text-align: center;">0.494</td> <td style="text-align: center;">0.0382</td> </tr> <tr> <td style="text-align: center;">33.0</td> <td style="text-align: center;">0.574</td> <td style="text-align: center;">0.570</td> <td style="text-align: center;">0.0439</td> </tr> </tbody> </table>		$t/^\circ C$	g(2)/100g(1)	g(2)/100g solution (compiler)	x_2 (compiler)	10.0	0.380	0.379	0.0296	16.0	0.424	0.422	0.0328	25.5	0.497	0.494	0.0382	33.0	0.574	0.570	0.0439
$t/^\circ C$	g(2)/100g(1)	g(2)/100g solution (compiler)	x_2 (compiler)																		
10.0	0.380	0.379	0.0296																		
16.0	0.424	0.422	0.0328																		
25.5	0.497	0.494	0.0382																		
33.0	0.574	0.570	0.0439																		
<p>AUXILIARY INFORMATION</p>																					
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The synthetic method similar to that described by Groschuff (ref 1) was used. A glass tube of about 100 mL capacity with a stopcock was filled with weighed (1) and (2). By shaking the tube after it had been warmed somewhat, the water was dissolved completely in the ester, after which the tube was placed in a 2-liter beaker filled with water and equipped with an agitator. By alternate slow cooling and warming, it was possible to determine, within about $0.5^\circ C$, the temperature at which water was precipitated from the solution, as shown by the clouding of the liquid which would clear again when the temperature was slightly increased.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified; distilled; b.p. range $164.5-165.5^\circ C$; boiled for some time in open flask before used.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm 0.25^\circ C$.</p> <p>REFERENCES:</p> <p>1. Groschuff, E. <i>Z. Elektrochem.</i> <u>1911</u>, 17, 348.</p>																				

<p>COMPONENTS:</p> <p>(1) Butanoic acid butyl ester (butyl butyrate); $C_8H_{16}O_2$; [109-21-7]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Doolittle, A.K. <i>Ind. End. Chem.</i> <u>1935</u>, 27, 1169-79.</p>
<p>VARIABLES:</p> <p>$T/K = 293$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of butanoic acid butyl ester in water at 20°C was reported to be 0.05 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 6×10^{-5}.</p> <p>The solubility of water in butanoic acid butyl ester at 20°C was reported to be 0.43 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.033.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The method was not specified.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, commercial product; purity 99%, b.p. range 152-170°C, d_4^{20} 0.874.</p> <p>(2) Not specified.</p>
	<p>ESTIMATED ERROR:</p> <p>Not specified.</p>
	<p>REFERENCES:</p>

COMPONENTS: (1) Butanoic acid butyl ester (butyl butyrate); $C_8H_{16}O_2$; [109-21-7] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Chebotaev, V.F.; Balashov, M.I.; Serafimov, L.A. <i>Fiz. Khim. Osn. Rektifikatsii</i> <u>1970</u> , 100-3.
VARIABLES: $T/K = 370$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: The solubility of water in butanoic acid butyl ester at $97.2^\circ C^a$ was reported to be 2.56 g(2)/100g sln. The corresponding mole fraction, x_2 , value calculated by the compiler is 0.1449. ^a Boiling temperature of two-phase mixture equal to $97.2^\circ C$ was taken from Ogrodnikov, Lesteva and Kogan (ref 2).	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The standard methods were used as described by Alders (ref 1). The phase composition was determined by chemical and refractometry methods. No chemical reaction was observed. The data and method were reported together with ternary system butanoic acid butyl ester-water-butanoic acid (butyl butyrate-water-butyric acid).	SOURCE AND PURITY OF MATERIALS: (1) Source not specified; distilled on a 20 TP column. (2) Not specified. ESTIMATED ERROR: Not specified. REFERENCES: 1. Alders, A. <i>Zhidkostnaya Ekstraktsiya</i> , I.L., Moskva, <u>1962</u> . 2. Ogrodnikov, S.K.; Lesteva, T.M.; Kogan, V.B. <i>Azeotropnye Smesi - spravochnik</i> , Khimiya, Leningrad, <u>1971</u> .

<p>COMPONENTS:</p> <p>(1) Butanoic acid butyl ester (butyl butyrate); $C_8H_{16}O_2$; [109-21-7]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u>, <i>31</i>, 56-70.</p>																																																											
<p>VARIABLES:</p> <p>$T/K = 273 - 364$</p>	<p>PREPARED BY:</p> <p>Z. Maczynska</p>																																																											
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of butanoic acid butyl ester and water</p> <table border="1" data-bbox="251 504 1260 1008"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>0.122</td><td>99.661</td><td>0.000152</td><td>0.97349</td></tr> <tr><td>9.5</td><td>0.102</td><td>99.630</td><td>0.000127</td><td>0.97113</td></tr> <tr><td>20.0</td><td>0.066</td><td>99.560</td><td>0.000082</td><td>0.96583</td></tr> <tr><td>29.7</td><td>0.056</td><td>99.573</td><td>0.000070</td><td>0.96681</td></tr> <tr><td>39.6</td><td>0.052</td><td>99.460</td><td>0.000065</td><td>0.95835</td></tr> <tr><td>50.0</td><td>0.051</td><td>99.423</td><td>0.000064</td><td>0.95560</td></tr> <tr><td>59.9</td><td>0.047</td><td>99.358</td><td>0.000059</td><td>0.95082</td></tr> <tr><td>70.2</td><td>0.047</td><td>99.349</td><td>0.000059</td><td>0.95016</td></tr> <tr><td>80.0</td><td>0.049</td><td>99.320</td><td>0.000061</td><td>0.94804</td></tr> <tr><td>90.3</td><td>0.051</td><td>99.317</td><td>0.000064</td><td>0.94782</td></tr> </tbody> </table> <p>std. dev. 0.001 0.004</p>		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	0.122	99.661	0.000152	0.97349	9.5	0.102	99.630	0.000127	0.97113	20.0	0.066	99.560	0.000082	0.96583	29.7	0.056	99.573	0.000070	0.96681	39.6	0.052	99.460	0.000065	0.95835	50.0	0.051	99.423	0.000064	0.95560	59.9	0.047	99.358	0.000059	0.95082	70.2	0.047	99.349	0.000059	0.95016	80.0	0.049	99.320	0.000061	0.94804	90.3	0.051	99.317	0.000064	0.94782
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																									
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																								
0	0.122	99.661	0.000152	0.97349																																																								
9.5	0.102	99.630	0.000127	0.97113																																																								
20.0	0.066	99.560	0.000082	0.96583																																																								
29.7	0.056	99.573	0.000070	0.96681																																																								
39.6	0.052	99.460	0.000065	0.95835																																																								
50.0	0.051	99.423	0.000064	0.95560																																																								
59.9	0.047	99.358	0.000059	0.95082																																																								
70.2	0.047	99.349	0.000059	0.95016																																																								
80.0	0.049	99.320	0.000061	0.94804																																																								
90.3	0.051	99.317	0.000064	0.94782																																																								
<p>AUXILIARY INFORMATION</p>																																																												
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, commercial sample; purity 99%; used as received.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Accuracy of method 0.1 wt% or less, for solubility, see above.</p> <p>REFERENCES:</p>																																																											

COMPONENTS: (1) Formic acid heptyl ester (heptyl formate); $C_8H_{16}O_2$; [112-23-2] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u> , <i>31</i> , 56-70.																																																											
VARIABLES: T/K = 273 - 364	PREPARED BY: Z. Maczynska																																																											
EXPERIMENTAL VALUES: Mutual solubility of formic acid heptyl ester and water <hr/> <table border="1" data-bbox="137 514 1118 1018"> <thead> <tr> <th rowspan="2">t/°C</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>-</td><td>99.32</td><td>-</td><td>0.9480</td></tr> <tr><td>9.1</td><td>-</td><td>99.37</td><td>-</td><td>0.9517</td></tr> <tr><td>19.5</td><td>0.162</td><td>99.32</td><td>0.000203</td><td>0.9480</td></tr> <tr><td>29.7</td><td>0.142</td><td>-</td><td>0.000178</td><td>-</td></tr> <tr><td>39.6</td><td>0.078</td><td>99.20</td><td>0.000097</td><td>0.9393</td></tr> <tr><td>50.0</td><td>0.071</td><td>99.15</td><td>0.000089</td><td>0.9358</td></tr> <tr><td>60.1</td><td>0.063</td><td>98.88</td><td>0.000079</td><td>0.9168</td></tr> <tr><td>70.1</td><td>0.087</td><td>98.94</td><td>0.000109</td><td>0.9210</td></tr> <tr><td>80.0</td><td>0.075</td><td>98.92</td><td>0.000094</td><td>0.9196</td></tr> <tr><td>90.6</td><td>0.056</td><td>98.81</td><td>0.000070</td><td>0.9120</td></tr> </tbody> </table> <hr/> std. dev. 0.002 0.01		t/°C	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	-	99.32	-	0.9480	9.1	-	99.37	-	0.9517	19.5	0.162	99.32	0.000203	0.9480	29.7	0.142	-	0.000178	-	39.6	0.078	99.20	0.000097	0.9393	50.0	0.071	99.15	0.000089	0.9358	60.1	0.063	98.88	0.000079	0.9168	70.1	0.087	98.94	0.000109	0.9210	80.0	0.075	98.92	0.000094	0.9196	90.6	0.056	98.81	0.000070	0.9120
t/°C	g(1)/100g sln		x_1 (compiler)																																																									
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																								
0	-	99.32	-	0.9480																																																								
9.1	-	99.37	-	0.9517																																																								
19.5	0.162	99.32	0.000203	0.9480																																																								
29.7	0.142	-	0.000178	-																																																								
39.6	0.078	99.20	0.000097	0.9393																																																								
50.0	0.071	99.15	0.000089	0.9358																																																								
60.1	0.063	98.88	0.000079	0.9168																																																								
70.1	0.087	98.94	0.000109	0.9210																																																								
80.0	0.075	98.92	0.000094	0.9196																																																								
90.6	0.056	98.81	0.000070	0.9120																																																								
AUXILIARY INFORMATION																																																												
METHOD/APPARATUS/PROCEDURE: The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial sample; purity 97%; used as received. (2) Not specified. <hr/> ESTIMATED ERROR: Accuracy of method 0.1 wt% or less, for solubility, see above. <hr/> REFERENCES:																																																											

<p>COMPONENTS:</p> <p>(1) Heptanoic acid methyl ester (methyl enanthate); $C_8H_{16}O_2$; [106-73-0]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u>, <i>31</i>, 56-70.</p>																																																											
<p>VARIABLES:</p> <p>$T/K = 273 - 364$</p>	<p>PREPARED BY:</p> <p>Z. Maczynska</p>																																																											
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of heptanoic acid methyl ester and water</p> <table border="1" data-bbox="253 512 1251 1020"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>0.105</td><td>99.45</td><td>0.000131</td><td>0.9576</td></tr> <tr><td>9.5</td><td>0.122</td><td>99.42</td><td>0.000152</td><td>0.9554</td></tr> <tr><td>19.3</td><td>0.094</td><td>99.33</td><td>0.000117</td><td>0.9488</td></tr> <tr><td>30.0</td><td>0.090</td><td>99.30</td><td>0.000112</td><td>0.9466</td></tr> <tr><td>39.6</td><td>0.063</td><td>99.36</td><td>0.000079</td><td>0.9509</td></tr> <tr><td>49.8</td><td>-</td><td>99.24</td><td>-</td><td>0.9422</td></tr> <tr><td>60.5</td><td>0.054</td><td>99.27</td><td>0.000067</td><td>0.9444</td></tr> <tr><td>70.5</td><td>0.087</td><td>99.28</td><td>0.000109</td><td>0.9451</td></tr> <tr><td>80.5</td><td>0.066</td><td>99.24</td><td>0.000082</td><td>0.9422</td></tr> <tr><td>90.5</td><td>0.053</td><td>99.13</td><td>0.000066</td><td>0.9343</td></tr> </tbody> </table> <p>std. dev. 0.001 0.01</p>		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	0.105	99.45	0.000131	0.9576	9.5	0.122	99.42	0.000152	0.9554	19.3	0.094	99.33	0.000117	0.9488	30.0	0.090	99.30	0.000112	0.9466	39.6	0.063	99.36	0.000079	0.9509	49.8	-	99.24	-	0.9422	60.5	0.054	99.27	0.000067	0.9444	70.5	0.087	99.28	0.000109	0.9451	80.5	0.066	99.24	0.000082	0.9422	90.5	0.053	99.13	0.000066	0.9343
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																									
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																								
0	0.105	99.45	0.000131	0.9576																																																								
9.5	0.122	99.42	0.000152	0.9554																																																								
19.3	0.094	99.33	0.000117	0.9488																																																								
30.0	0.090	99.30	0.000112	0.9466																																																								
39.6	0.063	99.36	0.000079	0.9509																																																								
49.8	-	99.24	-	0.9422																																																								
60.5	0.054	99.27	0.000067	0.9444																																																								
70.5	0.087	99.28	0.000109	0.9451																																																								
80.5	0.066	99.24	0.000082	0.9422																																																								
90.5	0.053	99.13	0.000066	0.9343																																																								
<p>AUXILIARY INFORMATION</p>																																																												
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, commercial sample; purity 99%; used as received.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Accuracy of method 0.1 wt% or less, for solubility, see above.</p> <p>REFERENCES:</p>																																																											

<p>COMPONENTS:</p> <p>(1) Hexanoic acid ethyl ester (ethyl caproate); $C_8H_{16}O_2$; [123-66-0]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Sobotka, H.; Kahn, J. <i>J. Am. Chem. Soc.</i> <u>1931</u>, 53, 2935-8.</p>
<p>VARIABLES:</p> <p>$T/K = 293$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of hexanoic acid ethyl ester in water at 20°C was reported to be 0.063 g(1)/100mL(2). The corresponding mass per cent and mole fraction, x_1, values calculated by the compiler are 0.063 g(1)/100g sln and 7.9×10^{-5}.</p> <p>Density of water $d_4^{20} = 0.9982$ (ref 1) was used in the calculation.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The titration method was used. The ester was added dropwise from a micro-burette with a capillary tip to 100, 250, or 500 mL of water in a narrow-mouthed stock bottle with a well-ground glass stopper. The bottle was shaken after each addition of ester. 1-5 mg of Sudan IV dye was put into the water to improve the end-point of the titration. At saturation, one additional drop of ester was sufficient to convert the floating rough indicator particles into dark transparent droplets.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified (Eastman Kodak Laboratories or synthesized); twice-distilled under reduced pressure; n_D^{20} 1.4089.</p> <p>(2) Distilled.</p>
	<p>ESTIMATED ERROR:</p> <p>Not specified.</p>
	<p>REFERENCES:</p> <p>1. <i>Selected Values of Properties of Hydrocarbons and Related Compounds</i>, API Research Project 44, Thermodynamics Research Center, Texas A and M University, Texas, <u>1973</u>.</p>

<p>COMPONENTS:</p> <p>(1) Hexanoic acid ethyl ester (ethyl caproate); $C_8H_{16}O_2$; [123-66-0]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u>, <i>31</i>, 56-70.</p>																																																																
<p>VARIABLES:</p> <p>$T/K = 273 - 364$</p>	<p>PREPARED BY:</p> <p>Z. Maczynska</p>																																																																
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of hexanoic acid ethyl ester and water</p> <table border="1" data-bbox="249 512 1248 1024"> <thead> <tr> <th rowspan="2">t/°C</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>0.107</td><td>99.64</td><td>0.000134</td><td>0.9719</td></tr> <tr><td>9.6</td><td>0.102</td><td>99.61</td><td>0.000127</td><td>0.9696</td></tr> <tr><td>19.7</td><td>0.083</td><td>99.53</td><td>0.000104</td><td>0.9636</td></tr> <tr><td>29.6</td><td>0.075</td><td>99.49</td><td>0.000094</td><td>0.9606</td></tr> <tr><td>39.8</td><td>0.070</td><td>99.44</td><td>0.000087</td><td>0.9568</td></tr> <tr><td>50.0</td><td>0.060</td><td>99.42</td><td>0.000075</td><td>0.9554</td></tr> <tr><td>60.6</td><td>0.049</td><td>99.38</td><td>0.000061</td><td>0.9524</td></tr> <tr><td>70.5</td><td>0.055</td><td>99.34</td><td>0.000069</td><td>0.9495</td></tr> <tr><td>80.4</td><td>0.055</td><td>99.29</td><td>0.000069</td><td>0.9458</td></tr> <tr><td>90.6</td><td>0.059</td><td>99.26</td><td>0.000074</td><td>0.9437</td></tr> <tr> <td>std. dev.</td> <td>0.007</td> <td>0.01</td> <td></td> <td></td> </tr> </tbody> </table>		t/°C	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	0.107	99.64	0.000134	0.9719	9.6	0.102	99.61	0.000127	0.9696	19.7	0.083	99.53	0.000104	0.9636	29.6	0.075	99.49	0.000094	0.9606	39.8	0.070	99.44	0.000087	0.9568	50.0	0.060	99.42	0.000075	0.9554	60.6	0.049	99.38	0.000061	0.9524	70.5	0.055	99.34	0.000069	0.9495	80.4	0.055	99.29	0.000069	0.9458	90.6	0.059	99.26	0.000074	0.9437	std. dev.	0.007	0.01		
t/°C	g(1)/100g sln		x_1 (compiler)																																																														
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																													
0	0.107	99.64	0.000134	0.9719																																																													
9.6	0.102	99.61	0.000127	0.9696																																																													
19.7	0.083	99.53	0.000104	0.9636																																																													
29.6	0.075	99.49	0.000094	0.9606																																																													
39.8	0.070	99.44	0.000087	0.9568																																																													
50.0	0.060	99.42	0.000075	0.9554																																																													
60.6	0.049	99.38	0.000061	0.9524																																																													
70.5	0.055	99.34	0.000069	0.9495																																																													
80.4	0.055	99.29	0.000069	0.9458																																																													
90.6	0.059	99.26	0.000074	0.9437																																																													
std. dev.	0.007	0.01																																																															
<p>AUXILIARY INFORMATION</p>																																																																	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, commercial sample; purity 99%; used as received.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Accuracy of method 0.1 wt% or less, for solubility, see above.</p> <p>REFERENCES:</p>																																																																

COMPONENTS: (1) Propanoic acid, 2-methyl-, butyl ester (butyl isobutyrate); $C_8H_{16}O_2$; [97-87-0] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u> , 31, 56-70.																																																											
VARIABLES: T/K = 273 - 364	PREPARED BY: Z. Maczynska																																																											
EXPERIMENTAL VALUES: Mutual solubility of 2-methylpropanoic acid butyl ester and water <table border="1" data-bbox="137 564 1125 1068"> <thead> <tr> <th rowspan="2">t/°C</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>0.090</td><td>99.64</td><td>0.000112</td><td>0.9719</td></tr> <tr><td>9.8</td><td>0.071</td><td>99.63</td><td>0.000089</td><td>0.9711</td></tr> <tr><td>19.6</td><td>0.077</td><td>99.57</td><td>0.000096</td><td>0.9666</td></tr> <tr><td>30.9</td><td>0.076</td><td>99.61</td><td>0.000095</td><td>0.9696</td></tr> <tr><td>40.1</td><td>0.051</td><td>99.53</td><td>0.000064</td><td>0.9636</td></tr> <tr><td>50.0</td><td>0.043</td><td>99.51</td><td>0.000054</td><td>0.9621</td></tr> <tr><td>60.0</td><td>0.048</td><td>99.42</td><td>0.000060</td><td>0.9554</td></tr> <tr><td>70.0</td><td>0.043</td><td>99.34</td><td>0.000054</td><td>0.9495</td></tr> <tr><td>80.2</td><td>0.044</td><td>99.35</td><td>0.000055</td><td>0.9502</td></tr> <tr><td>90.5</td><td>0.043</td><td>99.32</td><td>0.000054</td><td>0.9480</td></tr> </tbody> </table> std. dev. 0.003 0.01		t/°C	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	0.090	99.64	0.000112	0.9719	9.8	0.071	99.63	0.000089	0.9711	19.6	0.077	99.57	0.000096	0.9666	30.9	0.076	99.61	0.000095	0.9696	40.1	0.051	99.53	0.000064	0.9636	50.0	0.043	99.51	0.000054	0.9621	60.0	0.048	99.42	0.000060	0.9554	70.0	0.043	99.34	0.000054	0.9495	80.2	0.044	99.35	0.000055	0.9502	90.5	0.043	99.32	0.000054	0.9480
t/°C	g(1)/100g sln		x_1 (compiler)																																																									
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																								
0	0.090	99.64	0.000112	0.9719																																																								
9.8	0.071	99.63	0.000089	0.9711																																																								
19.6	0.077	99.57	0.000096	0.9666																																																								
30.9	0.076	99.61	0.000095	0.9696																																																								
40.1	0.051	99.53	0.000064	0.9636																																																								
50.0	0.043	99.51	0.000054	0.9621																																																								
60.0	0.048	99.42	0.000060	0.9554																																																								
70.0	0.043	99.34	0.000054	0.9495																																																								
80.2	0.044	99.35	0.000055	0.9502																																																								
90.5	0.043	99.32	0.000054	0.9480																																																								
AUXILIARY INFORMATION																																																												
METHOD/APPARATUS/PROCEDURE: The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial sample; purity 98%; used as received. (2) Not specified. ESTIMATED ERROR: Accuracy of method 0.1 wt% or less, for solubility, see above. REFERENCES:																																																											

<p>COMPONENTS:</p> <p>(1) Propanoic acid 3-methyl-1-butyl ester (isopentyl propionate); $C_8H_{16}O_2$; [105-68-0]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u>, <i>31</i>, 56-70.</p>																																																																
<p>VARIABLES:</p> <p>$T/K = 273 - 364$</p>	<p>PREPARED BY:</p> <p>Z. Maczynska</p>																																																																
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of propanoic acid 3-methyl-1-butyl ester and water</p> <table border="1" data-bbox="249 580 1251 1084"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">$g(1)/100g\ sln$</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>0.097</td><td>99.57</td><td>0.000121</td><td>0.9666</td></tr> <tr><td>9.3</td><td>0.100</td><td>99.57</td><td>0.000125</td><td>0.9666</td></tr> <tr><td>19.3</td><td>0.065</td><td>99.52</td><td>0.000081</td><td>0.9628</td></tr> <tr><td>29.7</td><td>0.056</td><td>99.46</td><td>0.000070</td><td>0.9583</td></tr> <tr><td>40.0</td><td>0.063</td><td>99.40</td><td>0.000079</td><td>0.9539</td></tr> <tr><td>49.8</td><td>0.045</td><td>99.47</td><td>0.000056</td><td>0.9591</td></tr> <tr><td>60.0</td><td>0.056</td><td>99.40</td><td>0.000070</td><td>0.9539</td></tr> <tr><td>70.0</td><td>0.075</td><td>-</td><td>0.000094</td><td>-</td></tr> <tr><td>79.7</td><td>0.057</td><td>99.22</td><td>0.000071</td><td>0.9408</td></tr> <tr><td>90.6</td><td>0.064</td><td>99.68</td><td>0.000080</td><td>0.9032</td></tr> <tr> <td>std. dev.</td> <td>0.001</td> <td>0.01</td> <td></td> <td></td> </tr> </tbody> </table>		$t/^\circ C$	$g(1)/100g\ sln$		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	0.097	99.57	0.000121	0.9666	9.3	0.100	99.57	0.000125	0.9666	19.3	0.065	99.52	0.000081	0.9628	29.7	0.056	99.46	0.000070	0.9583	40.0	0.063	99.40	0.000079	0.9539	49.8	0.045	99.47	0.000056	0.9591	60.0	0.056	99.40	0.000070	0.9539	70.0	0.075	-	0.000094	-	79.7	0.057	99.22	0.000071	0.9408	90.6	0.064	99.68	0.000080	0.9032	std. dev.	0.001	0.01		
$t/^\circ C$	$g(1)/100g\ sln$		x_1 (compiler)																																																														
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																													
0	0.097	99.57	0.000121	0.9666																																																													
9.3	0.100	99.57	0.000125	0.9666																																																													
19.3	0.065	99.52	0.000081	0.9628																																																													
29.7	0.056	99.46	0.000070	0.9583																																																													
40.0	0.063	99.40	0.000079	0.9539																																																													
49.8	0.045	99.47	0.000056	0.9591																																																													
60.0	0.056	99.40	0.000070	0.9539																																																													
70.0	0.075	-	0.000094	-																																																													
79.7	0.057	99.22	0.000071	0.9408																																																													
90.6	0.064	99.68	0.000080	0.9032																																																													
std. dev.	0.001	0.01																																																															
<p>AUXILIARY INFORMATION</p>																																																																	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, commercial sample; purity 99%; used as received.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Accuracy of method 0.1 wt% or less, for solubility, see above.</p> <p>REFERENCES:</p>																																																																

COMPONENTS: (1) Propanoic acid, 2-methyl-, 2-methylpropyl ester (isobutyl isobutyrate); $C_8H_{16}O_2$; [97-85-8] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u> , <i>31</i> , 56-70.																																																																
VARIABLES: $T/K = 273 - 363$	PREPARED BY: Z. Maczynska																																																																
EXPERIMENTAL VALUES: Mutual solubility of 2-methylpropanoic acid 2-methylpropyl ester and water <table border="1" data-bbox="141 572 1126 1076"> <thead> <tr> <th rowspan="2">t/°C</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>0.098</td><td>99.57</td><td>0.000122</td><td>0.9666</td></tr> <tr><td>9.3</td><td>-</td><td>99.48</td><td>-</td><td>0.9598</td></tr> <tr><td>19.4</td><td>0.050</td><td>99.46</td><td>0.000062</td><td>0.9583</td></tr> <tr><td>29.6</td><td>0.049</td><td>99.29</td><td>0.000061</td><td>0.9458</td></tr> <tr><td>39.5</td><td>0.050</td><td>99.43</td><td>0.000062</td><td>0.9561</td></tr> <tr><td>50.0</td><td>0.051</td><td>99.40</td><td>0.000064</td><td>0.9539</td></tr> <tr><td>59.8</td><td>0.046</td><td>99.18</td><td>0.000057</td><td>0.9379</td></tr> <tr><td>70.2</td><td>0.045</td><td>99.26</td><td>0.000056</td><td>0.9437</td></tr> <tr><td>80.1</td><td>0.041</td><td>99.24</td><td>0.000051</td><td>0.9422</td></tr> <tr><td>90.3</td><td>0.056</td><td>99.20</td><td>0.000070</td><td>0.9393</td></tr> <tr> <td>std. dev.</td> <td>0.001</td> <td>0.02</td> <td></td> <td></td> </tr> </tbody> </table>		t/°C	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	0.098	99.57	0.000122	0.9666	9.3	-	99.48	-	0.9598	19.4	0.050	99.46	0.000062	0.9583	29.6	0.049	99.29	0.000061	0.9458	39.5	0.050	99.43	0.000062	0.9561	50.0	0.051	99.40	0.000064	0.9539	59.8	0.046	99.18	0.000057	0.9379	70.2	0.045	99.26	0.000056	0.9437	80.1	0.041	99.24	0.000051	0.9422	90.3	0.056	99.20	0.000070	0.9393	std. dev.	0.001	0.02		
t/°C	g(1)/100g sln		x_1 (compiler)																																																														
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																													
0	0.098	99.57	0.000122	0.9666																																																													
9.3	-	99.48	-	0.9598																																																													
19.4	0.050	99.46	0.000062	0.9583																																																													
29.6	0.049	99.29	0.000061	0.9458																																																													
39.5	0.050	99.43	0.000062	0.9561																																																													
50.0	0.051	99.40	0.000064	0.9539																																																													
59.8	0.046	99.18	0.000057	0.9379																																																													
70.2	0.045	99.26	0.000056	0.9437																																																													
80.1	0.041	99.24	0.000051	0.9422																																																													
90.3	0.056	99.20	0.000070	0.9393																																																													
std. dev.	0.001	0.02																																																															
AUXILIARY INFORMATION																																																																	
METHOD/APPARATUS/PROCEDURE: The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial sample; purity 98%; used as received. (2) Not specified. ESTIMATED ERROR: Accuracy of method 0.1 wt% or less, for solubility, see above. REFERENCES:																																																																

COMPONENTS: (1) Propanoic acid pentyl ester (pentyl propionate); $C_8H_{16}O_2$; [624-54-4] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Hemptinne, A. <i>Z. Phys. Chem.</i> <u>1894</u> , 13, 561-9.
VARIABLES: $T/K = 298$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: The solubility of propanoic acid pentyl ester in water at 25°C was reported to be 0.908 g(1)/L sln.	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The analytical method was used. The mixture of water with excess ester was heated for some time in a water bath and the ester phase was filtered. A sample of known volume was then transferred to a smaller flask, heated with the known amount of baryta until complete saponification was obtained and then titrated. No further details were reported in the paper.	SOURCE AND PURITY OF MATERIALS: (1) Not specified. (2) Not specified.
	ESTIMATED ERROR: Not specified.
	REFERENCES:

COMPONENTS: (1) Acetic acid 2-butoxyethyl ester (2-butoxyethyl acetate); $C_8H_{16}O_3$; [112-07-2] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Doolittle, A.K. <i>Ind. End. Chem.</i> <u>1935</u> , 27, 1169-79.
VARIABLES: $T/K = 293$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: <p>The solubility of acetic acid 2-butoxyethyl ester in water at 20°C was reported to be 0.9 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 0.0010.</p> <p>The solubility of water in acetic acid 2-butoxyethyl ester at 20°C was reported to be 1.9 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.15.</p>	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The method was not specified.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial product; purity 99%, b.p. range 188-192°C, d_4^{20} 0.943. (2) Not specified.
	ESTIMATED ERROR: Not specified.
	REFERENCES:

<p>COMPONENTS:</p> <p>(1) Acetic acid phenylmethyl ester (benzyl acetate); $C_9H_{10}O_2$; [140-11-4]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988</p>
-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION:

Quantitative solubility data for the acetic acid phenylmethyl ester (1) - water (2) system have been reported in the publications listed in Table 1.

TABLE 1: Quantitative Solubility Studies of the
Acetic acid phenylmethyl ester (1) - Water (2) System

Reference	T/K	Solubility	Method
Goto et al. (ref 1)	363	(1) in (2)	GLC
Stephenson and Stuart (ref 2)	273-364	mutual	GLC

The original data in these publications are compiled in the Data Sheets immediately following this Critical Evaluation. For convenience, further discussion of this system will be divided into two parts.

1. SOLUBILITY OF ACETIC ACID PHENYLMETHYL ESTER (1) IN WATER (2)

All the available data for the solubility of acetic acid phenylmethyl ester (1) in water (2) are summarized in Table 2. At 363 K, the only temperature where comparison is possible, the data of Goto et al. (ref 1) and Stephenson and Stuart (ref 2) are in fair agreement. At other temperatures only the data of Stephenson and Stuart (ref 2) are available and in the absence of other studies must be regarded as Tentative, although it may be noted that the solubilities reported by these authors are usually reliable.

(continued next page)

<p>COMPONENTS:</p> <p>(1) Acetic acid phenylmethyl ester (benzyl acetate); $C_9H_{10}O_2$; [140-11-4]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988</p>
-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

TABLE 2: Tentative Solubilities
of Acetic acid phenylmethyl ester (1) in Water (2)

T/K	Solubilities		
	Reported values g(1)/100g sln	"Best" values ^a g(1)/100g sln	$10^4 x_1$
273	0.253 (ref 2)	0.25	3.0
283	0.253 (ref 2)	0.25	3.0
293	0.255* (ref 2)	0.26	3.1
298	0.258* (ref 2)	0.26	3.1
303	0.262* (ref 2)	0.26	3.1
313	0.277* (ref 2)	0.28	3.3
323	0.305* (ref 2)	0.31	3.6
333	0.338* (ref 2)	0.34	4.0
343	0.372* (ref 2)	0.37	4.5
353	0.409 (ref 2)	0.41	4.9
363	0.58 (ref 1), 0.447* (ref 2)	0.45	5.4

^a Rounded values of ref 2, see text.

2. SOLUBILITY OF WATER (2) IN ACETIC ACID PHENYLMETHYL ESTER (1)

The only data available for the solubility of water (2) in acetic acid phenylmethyl ester (1) are those of Stephenson and Stuart (ref 2) and so no Critical Evaluation is possible. The interested user is referred to the relevant Data Sheet for the experimental results but it may be noted that solubilities reported by these authors are usually reliable.

REFERENCES

- Goto, S.; Matsubara, M.; Washino, K. *Kagaku Kogaku* 1974, 38, 869-73.
- Stephenson, R.; Stuart, J. *J. Chem. Eng. Data* 1986, 31, 56-70.

<p>COMPONENTS:</p> <p>(1) Acetic acid phenylmethyl ester (benzyl acetate); C₉H₁₀O₂; [140-11-4]</p> <p>(2) Water; H₂O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Goto, S.; Matsubara, M.; Washino, K.</p> <p>Kagaku Kogaku <u>1974</u>, 38(12), 869-73.</p>
<p>VARIABLES:</p> <p>T/K = 363</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of acetic acid phenylmethyl ester in water at 90°C was reported to be 0.57 and 0.58 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 6.9×10^{-4}.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method was used. The samples of total volume about 50 cm³ containing acetic acid phenylmethyl ester, water and tetralin were thermostated at 90°C, mixed and separated over 1 h. Then, the phases were analysed by glc. The method and data were reported together with the ternary system acetic acid phenylmethyl ester-water-tetralin. Tetralin was not detected in the water phase.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified; purity >99 wt%; used as received.</p> <p>(2) Distilled.</p> <p>ESTIMATED ERROR:</p> <p>Soly. $<\pm 0.01$ g(1)/100g sln (compiler).</p> <p>REFERENCES:</p>

COMPONENTS: (1) Acetic acid phenylmethyl ester (benzyl acetate); $C_9H_{10}O_2$; [140-11-4] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u> , 31, 56-70.																																																											
VARIABLES: $T/K = 273 - 364$	PREPARED BY: Z. Maczynska																																																											
EXPERIMENTAL VALUES: Mutual solubility of acetic acid phenylmethyl ester and water <hr/> <table border="1" data-bbox="148 512 1126 1016"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>0.253</td><td>99.38</td><td>0.000304</td><td>0.9506</td></tr> <tr><td>9.7</td><td>0.253</td><td>99.36</td><td>0.000304</td><td>0.9490</td></tr> <tr><td>20.0</td><td>0.266</td><td>99.22</td><td>0.000320</td><td>0.9385</td></tr> <tr><td>30.7</td><td>0.240</td><td>99.08</td><td>0.000288</td><td>0.9281</td></tr> <tr><td>40.3</td><td>0.278</td><td>98.94</td><td>0.000334</td><td>0.9180</td></tr> <tr><td>49.8</td><td>0.305</td><td>98.79</td><td>0.000367</td><td>0.9073</td></tr> <tr><td>60.1</td><td>0.336</td><td>98.61</td><td>0.000404</td><td>0.8948</td></tr> <tr><td>70.4</td><td>0.375</td><td>98.41</td><td>0.000451</td><td>0.8813</td></tr> <tr><td>80.1</td><td>0.409</td><td>98.17</td><td>0.000492</td><td>0.8655</td></tr> <tr><td>90.4</td><td>0.448</td><td>97.83</td><td>0.000539</td><td>0.8439</td></tr> </tbody> </table> <hr/> std. dev. 0.002 0.03		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	0.253	99.38	0.000304	0.9506	9.7	0.253	99.36	0.000304	0.9490	20.0	0.266	99.22	0.000320	0.9385	30.7	0.240	99.08	0.000288	0.9281	40.3	0.278	98.94	0.000334	0.9180	49.8	0.305	98.79	0.000367	0.9073	60.1	0.336	98.61	0.000404	0.8948	70.4	0.375	98.41	0.000451	0.8813	80.1	0.409	98.17	0.000492	0.8655	90.4	0.448	97.83	0.000539	0.8439
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																									
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																								
0	0.253	99.38	0.000304	0.9506																																																								
9.7	0.253	99.36	0.000304	0.9490																																																								
20.0	0.266	99.22	0.000320	0.9385																																																								
30.7	0.240	99.08	0.000288	0.9281																																																								
40.3	0.278	98.94	0.000334	0.9180																																																								
49.8	0.305	98.79	0.000367	0.9073																																																								
60.1	0.336	98.61	0.000404	0.8948																																																								
70.4	0.375	98.41	0.000451	0.8813																																																								
80.1	0.409	98.17	0.000492	0.8655																																																								
90.4	0.448	97.83	0.000539	0.8439																																																								
AUXILIARY INFORMATION																																																												
METHOD/APPARATUS/PROCEDURE: The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial sample; purity 99%; used as received. (2) Not specified. <hr/> ESTIMATED ERROR: Accuracy of method 0.1 wt% or less, for solubility, see above. <hr/> REFERENCES:																																																											

<p>COMPONENTS:</p> <p>(1) Benzoic acid ethyl ester (ethyl benzoate); $C_9H_{10}O_2$; [93-89-0]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988</p>
------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION:

Quantitative solubility data for the benzoic acid ethyl ester (1) - water (2) system have been reported in the publications listed in Table 1.

TABLE 1: Quantitative Solubility Studies of the Benzoic acid ethyl ester (1) - Water (2) System

Reference	T/K	Solubility	Method
Krupatkin and Glagoleva (ref 1)	298	mutual	titration
Stephenson and Stuart (ref 2)	273-363	mutual	GLC

The original data in these publications are compiled in the Data Sheets immediately following this Critical Evaluation. For convenience, further discussion of this system will be divided into two parts.

1. SOLUBILITY OF BENZOIC ACID ETHYL ESTER (1) IN WATER (2)

All the available data for the solubility of benzoic acid ethyl ester (1) in water (2) are summarized in Table 2. At 298 K, the only temperature where comparison is possible, the data of Krupatkin and Glagoleva (ref 1) and Stephenson and Stuart (ref 2) are in only fair agreement. At other temperatures only the data of Stephenson and Stuart (ref 2) are available. Although the data reported by these authors are usually reliable their results for the present system are very scattered and, pending further studies, should be regarded as very Tentative.

(continued next page)

COMPONENTS: (1) Benzoic acid ethyl ester (ethyl benzoate); $C_9H_{10}O_2$; [93-89-0] (2) Water; H_2O ; [7732-18-5]	EVALUATOR: G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988
--------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

TABLE 2: Tentative Solubilities
of Benzoic acid ethyl ester (1) in Water (2)

T/K	Solubilities		
	Reported values	"Best" values ^a	
	g(1)/100g sln	g(1)/100g sln	$10^4 x_1$
273	0.108 (ref 2)	0.1	1
283	0.095* (ref 2)	0.1	1
293	0.085* (ref 2)	0.09	1
298	0.15 ^b (ref 1), 0.085* (ref 2)	0.09	1
303	0.090 (ref 2)	0.09	1
313	0.10* (ref 2)	0.1	1
323	0.11* (ref 2)	0.1	1
333	0.11* (ref 2)	0.1	1
343	0.12* (ref 2)	0.1	1
353	0.13* (ref 2)	0.1	1
363	0.14* (ref 2)	0.1	1

^a Rounded values of ref 2, but see text.

^b Not included in estimation of "Best" value.

2. SOLUBILITY OF WATER (2) IN BENZOIC ACID ETHYL ESTER (1)

All the available data for the solubility of water (2) in benzoic acid ethyl ester (1) are summarized in Table 3. The situation for the organic-rich phase is equivalent to that of the H_2O -rich phase and the user is referred to the comments in Part 1. of this Evaluation.

(continued next page)

<p>COMPONENTS:</p> <p>(1) Benzoic acid ethyl ester (ethyl benzoate); C₉H₁₀O₂; [93-89-0]</p> <p>(2) Water; H₂O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988</p>
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

TABLE 3: Tentative Solubilities
of Water (2) in Benzoic acid ethyl ester (1)

T/K	Solubilities		
	Reported values	"Best" values ^a	
	g(2)/100g sln	g(2)/100g sln	10 ² x ₂
283	1.1* (ref 2)	1.1	8
293	1.0* (ref 2)	1.0	7
298	0.57 ^b (ref 1), 1.0* (ref 2)	1.0	7
303	0.9* (ref 2)	0.9	7
313	0.9* (ref 2)	0.9	7
323	1.1* (ref 2)	1.1	8
333	1.0* (ref 2)	1.0	7

^a Rounded values of ref 2, but see text.

^b Not included in calculation of "Best" value.

REFERENCES

1. Krupatkin, I. L.; Glagoleva, M. F. *Zh. Prikl. Khim.* 1972, *45*, 1795-9.
2. Stephenson, R.; Stuart, J. *J. Chem. Eng. Data* 1986, *31*, 56-70.

COMPONENTS: (1) Benzoic acid ethyl ester (ethyl benzoate); $C_9H_{10}O_2$; [93-89-0] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Krupatkin, I.L.; Glagoleva, M.F. <i>Zh. Prikl. Khim.</i> <u>1972</u> , 45, 1795-9.
VARIABLES: $T/K = 298$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: <p>The solubility of benzoic acid ethyl ester in water at 25°C was reported to be 0.15 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 1.8×10^{-4}.</p> <p>The solubility of water in benzoic acid ethyl ester at 25°C was reported to be 0.57 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.046.</p>	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: <p>The titration method was used as described in the publication of Krupatkin and Glagoleva (ref 1). The samples were titrated up to turbidity. The data were reported together with the ternary system benzoic acid ethyl ester-water-2-furancarbal (ethyl benzoate-water-furfural). No further details were reported in the paper.</p>	SOURCE AND PURITY OF MATERIALS: (1) Source not specified; b.p. 212°C, $d_{25}^{25} 1.050$, $n_D^{25} 1.5050$. (2) Twice distilled. ESTIMATED ERROR: Not specified. REFERENCES: 1. Krupatkin, I.L.; Glagoleva, M.F. <i>Zh. Prikl. Khim.</i> <u>1969</u> , 42, 880.

<p>COMPONENTS:</p> <p>(1) Benzoic acid ethyl ester (ethyl benzoate); C₉H₁₀O₂; [93-89-0]</p> <p>(2) Water; H₂O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u>, <i>31</i>, 56-70.</p>																																																																
<p>VARIABLES:</p> <p>T/K = 273 - 363</p>	<p>PREPARED BY:</p> <p>Z. Maczynska</p>																																																																
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of benzoic acid ethyl ester and water</p> <table border="1" data-bbox="249 512 1247 1024"> <thead> <tr> <th rowspan="2">t/°C</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x₁ (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr> <td>0</td> <td>0.108</td> <td>98.81</td> <td>0.000130</td> <td>0.9088</td> </tr> <tr> <td>9.5</td> <td>-</td> <td>99.18</td> <td>-</td> <td>0.9355</td> </tr> <tr> <td>19.6</td> <td>0.085</td> <td>99.01</td> <td>0.000102</td> <td>0.9230</td> </tr> <tr> <td>30.5</td> <td>0.081</td> <td>99.08</td> <td>0.000097</td> <td>0.9281</td> </tr> <tr> <td>40.0</td> <td>0.106</td> <td>99.02</td> <td>0.000127</td> <td>0.9238</td> </tr> <tr> <td>50.0</td> <td>0.108</td> <td>98.88</td> <td>0.000130</td> <td>0.9137</td> </tr> <tr> <td>60.1</td> <td>0.117</td> <td>98.97</td> <td>0.000140</td> <td>0.9202</td> </tr> <tr> <td>70.5</td> <td>0.121</td> <td>-</td> <td>0.000145</td> <td>-</td> </tr> <tr> <td>80.2</td> <td>0.121</td> <td>-</td> <td>0.000145</td> <td>-</td> </tr> <tr> <td>90.3</td> <td>0.143</td> <td>-</td> <td>0.000172</td> <td>-</td> </tr> <tr> <td>std. dev.</td> <td>0.002</td> <td>0.01</td> <td></td> <td></td> </tr> </tbody> </table>		t/°C	g(1)/100g sln		x ₁ (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	0.108	98.81	0.000130	0.9088	9.5	-	99.18	-	0.9355	19.6	0.085	99.01	0.000102	0.9230	30.5	0.081	99.08	0.000097	0.9281	40.0	0.106	99.02	0.000127	0.9238	50.0	0.108	98.88	0.000130	0.9137	60.1	0.117	98.97	0.000140	0.9202	70.5	0.121	-	0.000145	-	80.2	0.121	-	0.000145	-	90.3	0.143	-	0.000172	-	std. dev.	0.002	0.01		
t/°C	g(1)/100g sln		x ₁ (compiler)																																																														
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																													
0	0.108	98.81	0.000130	0.9088																																																													
9.5	-	99.18	-	0.9355																																																													
19.6	0.085	99.01	0.000102	0.9230																																																													
30.5	0.081	99.08	0.000097	0.9281																																																													
40.0	0.106	99.02	0.000127	0.9238																																																													
50.0	0.108	98.88	0.000130	0.9137																																																													
60.1	0.117	98.97	0.000140	0.9202																																																													
70.5	0.121	-	0.000145	-																																																													
80.2	0.121	-	0.000145	-																																																													
90.3	0.143	-	0.000172	-																																																													
std. dev.	0.002	0.01																																																															
<p>AUXILIARY INFORMATION</p>																																																																	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, commercial sample; purity 97%; used as received.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Accuracy of method 0.1 wt% or less, for solubility, see above.</p> <p>REFERENCES:</p>																																																																

COMPONENTS: (1) Benzoic acid, 2-hydroxy-, ethyl ester (ethyl salicylate); $C_9H_{10}O_3$; [118-61-6] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u> , 31, 56-70.																																																																
VARIABLES: $T/K = 273 - 364$	PREPARED BY: Z. Maczynska																																																																
EXPERIMENTAL VALUES: Mutual solubility of 2-hydroxybenzoic acid ethyl ester and water <table border="1" data-bbox="159 568 1159 1072"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr> <td>0</td> <td>-</td> <td>99.80</td> <td>-</td> <td>0.9818</td> </tr> <tr> <td>9.2</td> <td>0.067</td> <td>99.76</td> <td>0.000073</td> <td>0.9783</td> </tr> <tr> <td>19.5</td> <td>0.036</td> <td>99.79</td> <td>0.000039</td> <td>0.9809</td> </tr> <tr> <td>29.8</td> <td>0.037</td> <td>99.68</td> <td>0.000040</td> <td>0.9712</td> </tr> <tr> <td>39.7</td> <td>0.029</td> <td>99.74</td> <td>0.000031</td> <td>0.9765</td> </tr> <tr> <td>49.8</td> <td>0.050</td> <td>99.73</td> <td>0.000054</td> <td>0.9756</td> </tr> <tr> <td>60.1</td> <td>0.040</td> <td>99.76</td> <td>0.000043</td> <td>0.9783</td> </tr> <tr> <td>70.1</td> <td>0.040</td> <td>99.54</td> <td>0.000043</td> <td>0.9591</td> </tr> <tr> <td>80.1</td> <td>0.069</td> <td>99.37</td> <td>0.000075</td> <td>0.9447</td> </tr> <tr> <td>90.5</td> <td>0.078</td> <td>99.33</td> <td>0.000085</td> <td>0.9414</td> </tr> <tr> <td>std. dev.</td> <td>0.003</td> <td>0.01</td> <td></td> <td></td> </tr> </tbody> </table>		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	-	99.80	-	0.9818	9.2	0.067	99.76	0.000073	0.9783	19.5	0.036	99.79	0.000039	0.9809	29.8	0.037	99.68	0.000040	0.9712	39.7	0.029	99.74	0.000031	0.9765	49.8	0.050	99.73	0.000054	0.9756	60.1	0.040	99.76	0.000043	0.9783	70.1	0.040	99.54	0.000043	0.9591	80.1	0.069	99.37	0.000075	0.9447	90.5	0.078	99.33	0.000085	0.9414	std. dev.	0.003	0.01		
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																														
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																													
0	-	99.80	-	0.9818																																																													
9.2	0.067	99.76	0.000073	0.9783																																																													
19.5	0.036	99.79	0.000039	0.9809																																																													
29.8	0.037	99.68	0.000040	0.9712																																																													
39.7	0.029	99.74	0.000031	0.9765																																																													
49.8	0.050	99.73	0.000054	0.9756																																																													
60.1	0.040	99.76	0.000043	0.9783																																																													
70.1	0.040	99.54	0.000043	0.9591																																																													
80.1	0.069	99.37	0.000075	0.9447																																																													
90.5	0.078	99.33	0.000085	0.9414																																																													
std. dev.	0.003	0.01																																																															
AUXILIARY INFORMATION																																																																	
METHOD/APPARATUS/PROCEDURE: The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial sample; purity 97%; used as received. (2) Not specified. ESTIMATED ERROR: Accuracy of method 0.1 wt% or less, for solubility, see above. REFERENCES:																																																																

<p>COMPONENTS:</p> <p>(1) 1,2,3-Propanetriol triacetate (glycerol triacetate); C₉H₁₄O₆; [102-76-1]</p> <p>(2) Water; H₂O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988</p>
-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION:

Quantitative solubility data for the 1,2,3-propanetriol triacetate (1) - water (2) system have been reported in the publications listed in Table 1.

TABLE 1: Quantitative Solubility Studies of the
1,2,3-Propanetriol triacetate (1) - Water (2) System

Reference	T/K	Solubility	Method
Othmer <i>et al.</i> (ref 1)	297 ^a	mutual	unspecified
Kraus <i>et al.</i> (ref 2)	298	mutual	analytical, Karl Fischer

^a 23.5 °C.

The original data in these publications are compiled in the Data Sheets immediately following this Critical Evaluation and are summarized in Tables 2 and 3 below. Unfortunately, these two independent studies are in poor agreement in both the H₂O-rich and the organic-rich phases (the slight temperature difference should not be important) and the average "Best" values must be regarded as very Tentative, pending further studies.

TABLE 2: Tentative Solubilities
of 1,2,3-Propanetriol triacetate (1) in Water (2)

T/K	Solubilities		
	Reported values	"Best" value ^a	
	g(1)/100g sln	g(1)/100g sln	10 ³ x ₁
298	5.5 ^b (ref 1), 7.4 (ref 2)	6.5 ± 1.0	5.7

^a Average value, but see text. x₁ has the same (relative) percentage uncertainty as the mass % solubility.

^b 296.7 K.

(continued next page)

COMPONENTS: (1) 1,2,3-Propanetriol triacetate (glycerol triacetate); $C_9H_{14}O_6$; [102-76-1] (2) Water; H_2O ; [7732-18-5]	EVALUATOR: G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988
-----------------------------------------------------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

TABLE 3: Tentative Solubilities
of Water (2) in 1,2,3-Propanetriol triacetate (1)

T/K	Solubilities		
	Reported values	"Best" value ^a	
	g(2)/100g sln	g(2)/100g sln	x_2
298	2.5 ^b (ref 1), 4.60 (ref 2)	3.5 ± 1.0	0.31

^a Average value, but see text. x_2 has the same status and (relative) percentage uncertainty as the mass % solubility.

^b 296.7 K.

REFERENCES

- Othmer, D. F.; White, R. E.; Trueger, E. *Ind. Eng. Chem.* 1941, 33, 1240-8.
- Kraus, K. A.; Rardon, R. J.; Baldwin, W. H. *J. Am. Chem. Soc.* 1964, 86, 2571-6.

<p>COMPONENTS:</p> <p>(1) 1,2,3-Propanetriol triacetate (glycerol triacetate); $C_9H_{14}O_6$; [102-76-1]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Othmer, D.F.; White, R.E.; Trueger, E.</p> <p>Ind. Eng. Chem. <u>1941</u>, 33, 1240-8.</p>
<p>VARIABLES:</p> <p>$T/K = 298$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of 1,2,3-propanetriol triacetate in water at 24.5°C was reported to be 5.5 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 0.0048.</p> <p>The solubility of water in 1,2,3-propanetriol triacetate at 24.5°C was reported to be 2.5 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.24.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>Nothing was specified in the paper.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Not specified.</p> <p>(2) Not specified.</p>
	<p>ESTIMATED ERROR:</p> <p>Temp. $\pm 0.5^\circ C$ (mean of reported range).</p>
	<p>REFERENCES:</p>

COMPONENTS: (1) 1,2,3-Propanetriol triacetate (glycerol triacetate); $C_9H_{14}O_6$; [102-76-1] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Kraus, K.A.; Raridon, R.J.; Baldwin, W.H. <i>J. Am. Chem. Soc.</i> <u>1964</u> , 86, 2571-6.
VARIABLES: $T/K = 298$	PREPARED BY: Z. Maczynska
EXPERIMENTAL VALUES: The solubility of 1,2,3-propanetriol triacetate in water at 25°C was reported to be 7.4 g(1)/100g sln. The corresponding mole fraction, x_1 , value calculated by the compiler is 0.0066. The solubility of water in 1,2,3-propanetriol triacetate at 25°C was reported to be 4.60 g(2)/100g sln. The corresponding mole fraction, x_2 , value calculated by the compiler is 0.369.	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The analytical methods were used. The mutual solubilities of (1) and (2) were determined by shaking mixtures in stoppered tubes in a constant temperature bath. The (1)-rich phases were analyzed for water by the use of Karl Fischer titrations. The (2)-rich phases were analyzed for acetate by a semimicro adaptation of the standard saponification technique whereby back titration of excess alkali is performed on a boiling solution to minimize carbonate error, and in a flask under a reflux condenser to minimize loss of acetic acid.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified; commercial product; distilled under vacuum and middle fraction 60-80% collected. (2) Not specified. ESTIMATED ERROR: Soly. ± 5 g(1)/100g sln and ± 1 g(2)/100g sln (precision). REFERENCES:

COMPONENTS: (1) Propanoic acid cyclohexyl ester (<i>cyclohexyl propionate</i>); $C_9H_{16}O_2$; [6222-35-1] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u> , <i>31</i> , 56-70.																																																											
VARIABLES: $T/K = 273 - 364$	PREPARED BY: Z. Maczynska																																																											
EXPERIMENTAL VALUES: Mutual solubility of propanoic acid cyclohexyl ester and water <table border="1" data-bbox="256 520 1250 1024"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>0.109</td><td>99.17</td><td>0.000126</td><td>0.9323</td></tr> <tr><td>9.5</td><td>0.096</td><td>99.29</td><td>0.000111</td><td>0.9416</td></tr> <tr><td>20.2</td><td>0.079</td><td>99.21</td><td>0.000091</td><td>0.9354</td></tr> <tr><td>30.6</td><td>0.073</td><td>99.06</td><td>0.000084</td><td>0.9239</td></tr> <tr><td>39.6</td><td>0.065</td><td>99.18</td><td>0.000075</td><td>0.9331</td></tr> <tr><td>50.0</td><td>0.061</td><td>99.32</td><td>0.000070</td><td>0.9439</td></tr> <tr><td>60.1</td><td>0.073</td><td>99.28</td><td>0.000084</td><td>0.9408</td></tr> <tr><td>70.2</td><td>0.066</td><td>99.06</td><td>0.000076</td><td>0.9239</td></tr> <tr><td>80.2</td><td>0.080</td><td>98.91</td><td>0.000092</td><td>0.9127</td></tr> <tr><td>90.6</td><td>0.075</td><td>98.78</td><td>0.000086</td><td>0.9032</td></tr> </tbody> </table> std. dev. 0.001 0.05		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	0.109	99.17	0.000126	0.9323	9.5	0.096	99.29	0.000111	0.9416	20.2	0.079	99.21	0.000091	0.9354	30.6	0.073	99.06	0.000084	0.9239	39.6	0.065	99.18	0.000075	0.9331	50.0	0.061	99.32	0.000070	0.9439	60.1	0.073	99.28	0.000084	0.9408	70.2	0.066	99.06	0.000076	0.9239	80.2	0.080	98.91	0.000092	0.9127	90.6	0.075	98.78	0.000086	0.9032
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																									
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																								
0	0.109	99.17	0.000126	0.9323																																																								
9.5	0.096	99.29	0.000111	0.9416																																																								
20.2	0.079	99.21	0.000091	0.9354																																																								
30.6	0.073	99.06	0.000084	0.9239																																																								
39.6	0.065	99.18	0.000075	0.9331																																																								
50.0	0.061	99.32	0.000070	0.9439																																																								
60.1	0.073	99.28	0.000084	0.9408																																																								
70.2	0.066	99.06	0.000076	0.9239																																																								
80.2	0.080	98.91	0.000092	0.9127																																																								
90.6	0.075	98.78	0.000086	0.9032																																																								
AUXILIARY INFORMATION																																																												
METHOD/APPARATUS/PROCEDURE: The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial sample; purity 99%; used as received. (2) Not specified. ESTIMATED ERROR: Accuracy of method 0.1 wt% or less, for solubility, see above. REFERENCES:																																																											

COMPONENTS: (1) Pentanedioic acid diethyl ester (<i>diethyl glutarate</i>); $C_9H_{16}O_4$; [818-38-2] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Sobotka, H.; Kahn, J. <i>J. Am. Chem. Soc.</i> <u>1931</u> , 53, 2935-8.
VARIABLES: $T/K = 293$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: The solubility of pentanedioic acid diethyl ester in water at 20°C was reported to be 0.882 g(1)/100mL(2). The corresponding mass per cent and mole fraction, x_1 , values calculated by the compiler are 0.876 g(1)/100g sln and 8.45×10^{-4} . Density of water $d_4^{20} = 0.9982$ (ref 1) was used in the calculation.	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The titration method was used. The ester was added dropwise from a micro-burette with a capillary tip to 100, 250, or 500 mL of water in a narrow-mouthed stock bottle with a well-ground glass stopper. The bottle was shaken after each addition of ester. 1-5 mg of Sudan IV dye was put into the water to improve the end-point of the titration. At saturation, one additional drop of ester was sufficient to convert the floating rough indicator particles into dark transparent droplets.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified (Eastman Kodak Laboratories or synthesized); twice-distilled under reduced pressure; d_4^{20} 1.0220, n_D^{20} 1.4242. (2) Distilled. ESTIMATED ERROR: Not specified. REFERENCES: 1. <i>Selected Values of Properties of Hydrocarbons and Related Compounds</i> , API Research Project 44, Thermodynamics Research Center, Texas A and M University, Texas, 1973.

COMPONENTS: (1) Acetic acid heptyl ester (heptyl acetate); $C_9H_{18}O_2$; [112-06-1] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u> , 31, 56-70.																																																											
VARIABLES: $T/K = 273 - 364$	PREPARED BY: Z. Maczynska																																																											
EXPERIMENTAL VALUES: Mutual solubility of acetic acid heptyl ester and water <hr/> <table border="1" data-bbox="249 520 1243 1024"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>0.023</td><td>99.66</td><td>0.000026</td><td>0.9709</td></tr> <tr><td>9.2</td><td>0.018</td><td>99.62</td><td>0.000020</td><td>0.9676</td></tr> <tr><td>19.6</td><td>0.020</td><td>99.55</td><td>0.000023</td><td>0.9618</td></tr> <tr><td>30.6</td><td>0.023</td><td>99.52</td><td>0.000026</td><td>0.9593</td></tr> <tr><td>40.3</td><td>0.021</td><td>99.41</td><td>0.000024</td><td>0.9504</td></tr> <tr><td>50.0</td><td>0.020</td><td>99.41</td><td>0.000023</td><td>0.9504</td></tr> <tr><td>60.2</td><td>0.019</td><td>99.37</td><td>0.000022</td><td>0.9472</td></tr> <tr><td>70.2</td><td>0.017</td><td>99.35</td><td>0.000019</td><td>0.9456</td></tr> <tr><td>80.4</td><td>0.015</td><td>99.33</td><td>0.000017</td><td>0.9440</td></tr> <tr><td>90.6</td><td>0.019</td><td>99.33</td><td>0.000022</td><td>0.9440</td></tr> </tbody> </table> <hr/> std. dev. 0.001 0.01		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	0.023	99.66	0.000026	0.9709	9.2	0.018	99.62	0.000020	0.9676	19.6	0.020	99.55	0.000023	0.9618	30.6	0.023	99.52	0.000026	0.9593	40.3	0.021	99.41	0.000024	0.9504	50.0	0.020	99.41	0.000023	0.9504	60.2	0.019	99.37	0.000022	0.9472	70.2	0.017	99.35	0.000019	0.9456	80.4	0.015	99.33	0.000017	0.9440	90.6	0.019	99.33	0.000022	0.9440
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																									
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																								
0	0.023	99.66	0.000026	0.9709																																																								
9.2	0.018	99.62	0.000020	0.9676																																																								
19.6	0.020	99.55	0.000023	0.9618																																																								
30.6	0.023	99.52	0.000026	0.9593																																																								
40.3	0.021	99.41	0.000024	0.9504																																																								
50.0	0.020	99.41	0.000023	0.9504																																																								
60.2	0.019	99.37	0.000022	0.9472																																																								
70.2	0.017	99.35	0.000019	0.9456																																																								
80.4	0.015	99.33	0.000017	0.9440																																																								
90.6	0.019	99.33	0.000022	0.9440																																																								
AUXILIARY INFORMATION																																																												
METHOD/APPARATUS/PROCEDURE: The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial sample; purity 98%; used as received. (2) Not specified. <hr/> ESTIMATED ERROR: Accuracy of method 0.1 wt% or less, for solubility, see above. <hr/> REFERENCES:																																																											

COMPONENTS: (1) Butanoic acid 3-methyl-1-butyl ester (<i>isopentyl butyrate</i>); $C_9H_{18}O_2$; [106-27-4] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u> , <i>31</i> , 56-70.																																																											
VARIABLES: $T/K = 273 - 364$	PREPARED BY: Z. Maczynska																																																											
EXPERIMENTAL VALUES: Mutual solubility of butanoic acid 3-methyl-1-butyl ester and water <hr/> <table border="1" data-bbox="154 524 1155 1018"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr> <td>0</td> <td>-</td> <td>99.75</td> <td>-</td> <td>0.9784</td> </tr> <tr> <td>9.4</td> <td>0.035</td> <td>99.71</td> <td>0.000040</td> <td>0.9751</td> </tr> <tr> <td>19.6</td> <td>0.022</td> <td>99.76</td> <td>0.000025</td> <td>0.9793</td> </tr> <tr> <td>30.9</td> <td>0.027</td> <td>99.62</td> <td>0.000031</td> <td>0.9676</td> </tr> <tr> <td>39.9</td> <td>0.015</td> <td>99.59</td> <td>0.000017</td> <td>0.9651</td> </tr> <tr> <td>49.8</td> <td>0.037</td> <td>99.69</td> <td>0.000042</td> <td>0.9734</td> </tr> <tr> <td>59.7</td> <td>0.013</td> <td>99.68</td> <td>0.000015</td> <td>0.9726</td> </tr> <tr> <td>70.2</td> <td>0.014</td> <td>99.61</td> <td>0.000016</td> <td>0.9667</td> </tr> <tr> <td>80.1</td> <td>0.024</td> <td>99.52</td> <td>0.000027</td> <td>0.9593</td> </tr> <tr> <td>90.7</td> <td>0.020</td> <td>99.50</td> <td>0.000023</td> <td>0.9577</td> </tr> </tbody> </table> <hr/> std. dev. 0.004 0.01		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	-	99.75	-	0.9784	9.4	0.035	99.71	0.000040	0.9751	19.6	0.022	99.76	0.000025	0.9793	30.9	0.027	99.62	0.000031	0.9676	39.9	0.015	99.59	0.000017	0.9651	49.8	0.037	99.69	0.000042	0.9734	59.7	0.013	99.68	0.000015	0.9726	70.2	0.014	99.61	0.000016	0.9667	80.1	0.024	99.52	0.000027	0.9593	90.7	0.020	99.50	0.000023	0.9577
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																									
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																								
0	-	99.75	-	0.9784																																																								
9.4	0.035	99.71	0.000040	0.9751																																																								
19.6	0.022	99.76	0.000025	0.9793																																																								
30.9	0.027	99.62	0.000031	0.9676																																																								
39.9	0.015	99.59	0.000017	0.9651																																																								
49.8	0.037	99.69	0.000042	0.9734																																																								
59.7	0.013	99.68	0.000015	0.9726																																																								
70.2	0.014	99.61	0.000016	0.9667																																																								
80.1	0.024	99.52	0.000027	0.9593																																																								
90.7	0.020	99.50	0.000023	0.9577																																																								
AUXILIARY INFORMATION																																																												
METHOD/APPARATUS/PROCEDURE: The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial sample; purity 99%; used as received. (2) Not specified. <hr/> ESTIMATED ERROR: Accuracy of method 0.1 wt% or less, for solubility, see above. <hr/> REFERENCES:																																																											

<p>COMPONENTS:</p> <p>(1) Butanoic acid pentyl ester (<i>pentyl butyrate</i>); $C_9H_{18}O_2$; [540-18-1]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988</p>
-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION:

Quantitative solubility data for the butanoic acid pentyl ester (1) - water (2) system have been reported in the publications listed in Table 1.

TABLE 1: Quantitative Solubility Studies of the
Butanoic acid pentyl ester (1) - Water (2) System

Reference	T/K	Solubility	Method
Hemptinne (ref 1)	298	(1) in (2)	analytical
Stephenson and Stuart (ref 2)	273-364	mutual	GLC

The original data in these publications are compiled in the Data Sheets immediately following this Critical Evaluation. For convenience, further discussion of this system will be divided into two parts.

1. SOLUBILITY OF BUTANOIC ACID PENTYL ESTER (1) IN WATER (2)

All the available data for the solubility of butanoic acid pentyl ester (1) in water (2) are summarized in Table 2. At 298 K, the only temperature where comparison is possible, the data of Hemptinne (ref 1) and Stephenson and Stuart (ref 2) are in only fair agreement. At other temperatures only the data of Stephenson and Stuart (ref 2) are available. Although the solubilities reported by these authors are usually reliable it should be noted that their results for this system are very scattered. Consequently, all the values in Table 2 should be regarded as very Tentative, pending further studies.

(continued next page)

COMPONENTS: (1) Butanoic acid pentyl ester (pentyl butyrate); $C_9H_{18}O_2$; [540-18-1] (2) Water; H_2O ; [7732-18-5]	EVALUATOR: G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988
----------------------------------------------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

TABLE 2: Tentative Solubilities
of Butanoic acid pentyl ester (1) in Water (2)

T/K	Solubilities		
	Reported values ^a		"Best" values ^b
	g(1)/100g sln		g(1)/100g sln $10^5 x_1$
273	0.03* (ref 2)		0.03 3
283	0.03* (ref 2)		0.03 3
293	0.03* (ref 2)		0.03 3
298	0.050 (ref 1), 0.03* (ref 2)		0.04 ± 0.01^c 5
303	0.03* (ref 2)		0.03 3
313	0.03* (ref 2)		0.03 3
323	0.03* (ref 2)		0.03 3
333	0.03* (ref 2)		0.03 3
343	0.03* (ref 2)		0.03 3
353	0.02* (ref 2)		0.02 2
363	0.02* (ref 2)		0.02 2

^a Data in ref 2 very scattered.

^b Rounded values of ref 2, except at 298 K.

^c Average value; x_1 has the same (relative) percentage uncertainty as the mass % solubility.

2. SOLUBILITY OF WATER (2) IN BUTANOIC ACID PENTYL ESTER (1)

Only the data of Stephenson and Stuart (ref 2) are available for the solubility of water (2) in butanoic acid pentyl ester (1) and so no Critical Evaluation is possible. The interested user is referred to the relevant Data Sheet for the experimental solubilities. It should be noted, however, that whilst the solubilities reported by these authors are generally reliable, their results for this system are very scattered. Further studies are clearly required.

REFERENCES

- Hemptinne, A. *Z. Phys. Chem.* 1894, 13, 561-9.
- Stephenson, R.; Stuart, J. *J. Chem. Eng. Data* 1986, 31, 56-70.

<p>COMPONENTS:</p> <p>(1) Butanoic acid pentyl ester (<i>pentyl butyrate</i>); $C_9H_{18}O_2$; [540-18-1]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Hemptinne, A. <i>Z. Phys. Chem.</i> <u>1894</u>, 13, 561-9.</p>
<p>VARIABLES:</p> <p>$T/K = 298$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of butanoic acid pentyl ester in water at 25°C was reported to be 0.505 g(1)/L sln.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method was used. The mixture of water with excess ester was heated for some time in a water bath and the ester phase was filtered. A sample of known volume was then transferred to a smaller flask, heated with the known amount of baryta until complete saponification was obtained and then titrated. No further details were reported in the paper.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Not specified. (2) Not specified.</p>
	<p>ESTIMATED ERROR:</p> <p>Not specified.</p>
	<p>REFERENCES:</p>

COMPONENTS: (1) Butanoic acid pentyl ester (pentyl butyrate); $C_9H_{18}O_2$; [540-18-1] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u> , 31, 56-70.																																																											
VARIABLES: $T/K = 273 - 364$	PREPARED BY: Z. Maczynska																																																											
EXPERIMENTAL VALUES: Mutual solubility of butanoic acid pentyl ester and water <hr/> <table border="1" data-bbox="148 506 1131 1010"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>0.026</td><td>99.54</td><td>0.000030</td><td>0.9610</td></tr> <tr><td>9.4</td><td>0.034</td><td>99.66</td><td>0.000039</td><td>0.9709</td></tr> <tr><td>19.2</td><td>0.025</td><td>99.60</td><td>0.000028</td><td>0.9659</td></tr> <tr><td>29.6</td><td>0.024</td><td>99.65</td><td>0.000027</td><td>0.9701</td></tr> <tr><td>40.1</td><td>0.030</td><td>99.48</td><td>0.000034</td><td>0.9561</td></tr> <tr><td>50.0</td><td>0.032</td><td>99.61</td><td>0.000036</td><td>0.9667</td></tr> <tr><td>60.0</td><td>0.028</td><td>99.56</td><td>0.000032</td><td>0.9626</td></tr> <tr><td>70.3</td><td>-</td><td>99.38</td><td>-</td><td>0.9480</td></tr> <tr><td>80.0</td><td>0.019</td><td>99.52</td><td>0.000022</td><td>0.9593</td></tr> <tr><td>90.5</td><td>0.025</td><td>99.44</td><td>0.000028</td><td>0.9529</td></tr> </tbody> </table> <hr/> std. dev. 0.002 0.01		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	0.026	99.54	0.000030	0.9610	9.4	0.034	99.66	0.000039	0.9709	19.2	0.025	99.60	0.000028	0.9659	29.6	0.024	99.65	0.000027	0.9701	40.1	0.030	99.48	0.000034	0.9561	50.0	0.032	99.61	0.000036	0.9667	60.0	0.028	99.56	0.000032	0.9626	70.3	-	99.38	-	0.9480	80.0	0.019	99.52	0.000022	0.9593	90.5	0.025	99.44	0.000028	0.9529
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																									
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																								
0	0.026	99.54	0.000030	0.9610																																																								
9.4	0.034	99.66	0.000039	0.9709																																																								
19.2	0.025	99.60	0.000028	0.9659																																																								
29.6	0.024	99.65	0.000027	0.9701																																																								
40.1	0.030	99.48	0.000034	0.9561																																																								
50.0	0.032	99.61	0.000036	0.9667																																																								
60.0	0.028	99.56	0.000032	0.9626																																																								
70.3	-	99.38	-	0.9480																																																								
80.0	0.019	99.52	0.000022	0.9593																																																								
90.5	0.025	99.44	0.000028	0.9529																																																								
AUXILIARY INFORMATION																																																												
METHOD/APPARATUS/PROCEDURE: The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial sample; purity 99%; used as received. (2) Not specified. <hr/> ESTIMATED ERROR: Accuracy of method 0.1 wt% or less, for solubility, see above. <hr/> REFERENCES:																																																											

COMPONENTS: (1) Formic acid octyl ester (octyl formate); $C_9H_{18}O_2$; [112-32-3] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u> , <i>31</i> , 56-70.																																																											
VARIABLES: $T/K = 273 - 364$	PREPARED BY: Z. Maczynska																																																											
EXPERIMENTAL VALUES: Mutual solubility of formic acid octyl ester and water <hr/> <table border="1" data-bbox="268 533 1268 1044"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>-</td><td>99.579</td><td>-</td><td>0.96419</td></tr> <tr><td>9.2</td><td>0.078</td><td>99.555</td><td>0.000089</td><td>0.96222</td></tr> <tr><td>19.0</td><td>0.064</td><td>99.583</td><td>0.000073</td><td>0.96452</td></tr> <tr><td>29.4</td><td>0.039</td><td>99.572</td><td>0.000044</td><td>0.96361</td></tr> <tr><td>39.5</td><td>0.040</td><td>99.555</td><td>0.000045</td><td>0.96222</td></tr> <tr><td>50.0</td><td>0.071</td><td>99.545</td><td>0.000081</td><td>0.96140</td></tr> <tr><td>60.0</td><td>0.028</td><td>99.522</td><td>0.000032</td><td>0.95952</td></tr> <tr><td>70.1</td><td>0.044</td><td>99.506</td><td>0.000050</td><td>0.95821</td></tr> <tr><td>80.2</td><td>0.036</td><td>99.426</td><td>0.000041</td><td>0.95174</td></tr> <tr><td>90.5</td><td>0.061</td><td>99.461</td><td>0.000069</td><td>0.95456</td></tr> </tbody> </table> <hr/> std. dev. 0.003 0.003		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	-	99.579	-	0.96419	9.2	0.078	99.555	0.000089	0.96222	19.0	0.064	99.583	0.000073	0.96452	29.4	0.039	99.572	0.000044	0.96361	39.5	0.040	99.555	0.000045	0.96222	50.0	0.071	99.545	0.000081	0.96140	60.0	0.028	99.522	0.000032	0.95952	70.1	0.044	99.506	0.000050	0.95821	80.2	0.036	99.426	0.000041	0.95174	90.5	0.061	99.461	0.000069	0.95456
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																									
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																								
0	-	99.579	-	0.96419																																																								
9.2	0.078	99.555	0.000089	0.96222																																																								
19.0	0.064	99.583	0.000073	0.96452																																																								
29.4	0.039	99.572	0.000044	0.96361																																																								
39.5	0.040	99.555	0.000045	0.96222																																																								
50.0	0.071	99.545	0.000081	0.96140																																																								
60.0	0.028	99.522	0.000032	0.95952																																																								
70.1	0.044	99.506	0.000050	0.95821																																																								
80.2	0.036	99.426	0.000041	0.95174																																																								
90.5	0.061	99.461	0.000069	0.95456																																																								
AUXILIARY INFORMATION																																																												
METHOD/APPARATUS/PROCEDURE: The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial sample; purity 99%; used as received. (2) Not specified. <hr/> ESTIMATED ERROR: Accuracy of method 0.1 wt% or less, for solubility, see above. <hr/> REFERENCES:																																																											

COMPONENTS: (1) Heptanoic acid ethyl ester (ethyl heptanoate); $C_9H_{18}O_2$; [106-30-9] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Sobotka, H.; Kahn, J. <i>J. Am. Chem. Soc.</i> <u>1931</u> , 53, 2935-8.
VARIABLES: $T/K = 293$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: The solubility of heptanoic acid ethyl ester in water at 20°C was reported to be 0.029 g(1)/100mL(2). The corresponding mass per cent and mole fraction, x_1 , values calculated by the compiler are 0.029 g(1)/100g sln and 3.3×10^{-5} . Density of water $d_4^{20} = 0.9982$ (ref 1) was used in the calculation.	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The titration method was used. The ester was added dropwise from a micro-burette with a capillary tip to 100, 250, or 500 mL of water in a narrow-mouthed stock bottle with a well-ground glass stopper. The bottle was shaken after each addition of ester. 1-5 mg of Sudan IV dye was put into the water to improve the end-point of the titration. At saturation, one additional drop of ester was sufficient to convert the floating rough indicator particles into dark transparent droplets.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified (Eastman Kodak Laboratories or synthesized); twice-distilled under reduced pressure; n_D^{20} 1.4137. (2) Distilled. ESTIMATED ERROR: Not specified. REFERENCES: 1. <i>Selected Values of Properties of Hydrocarbons and Related Compounds</i> , API Research Project 44, Thermodynamics Research Center, Texas A and M University, Texas, <u>1973</u> .

<p>COMPONENTS:</p> <p>(1) Pentanoic acid 1-butyl ester (butyl valerate); $C_9H_{18}O_2$; [591-68-4]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Bomshtein, A.L.; Trofimov, A.N.; Serafimov, L.A.</p> <p>Zh. Prikl. Khim. <u>1978</u>, 51, 1280-2.</p>																																																	
<p>VARIABLES:</p> <p>$T/K = 293 - 363$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>																																																	
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of pentanoic acid 1-butyl ester and water</p> <table border="1" data-bbox="263 520 1257 951"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">x_1</th> <th colspan="2">g(1)/100g sln (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr> <td>20</td> <td>0.00002</td> <td>0.9037</td> <td>0.02</td> <td>98.80</td> </tr> <tr> <td>30</td> <td>0.00009</td> <td>0.8821</td> <td>0.08</td> <td>98.50</td> </tr> <tr> <td>40</td> <td>0.00015</td> <td>0.8751</td> <td>0.13</td> <td>98.40</td> </tr> <tr> <td>50</td> <td>0.00021</td> <td>0.8514</td> <td>0.18</td> <td>98.05</td> </tr> <tr> <td>60</td> <td>0.00027</td> <td>0.8319</td> <td>0.24</td> <td>97.75</td> </tr> <tr> <td>70</td> <td>0.00037</td> <td>0.8163</td> <td>0.32</td> <td>97.50</td> </tr> <tr> <td>80</td> <td>0.00039</td> <td>0.8011</td> <td>0.34</td> <td>97.25</td> </tr> <tr> <td>90</td> <td>0.00046</td> <td>0.7865</td> <td>0.40</td> <td>97.00</td> </tr> </tbody> </table>		$t/^\circ C$	x_1		g(1)/100g sln (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	20	0.00002	0.9037	0.02	98.80	30	0.00009	0.8821	0.08	98.50	40	0.00015	0.8751	0.13	98.40	50	0.00021	0.8514	0.18	98.05	60	0.00027	0.8319	0.24	97.75	70	0.00037	0.8163	0.32	97.50	80	0.00039	0.8011	0.34	97.25	90	0.00046	0.7865	0.40	97.00
$t/^\circ C$	x_1		g(1)/100g sln (compiler)																																															
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																														
20	0.00002	0.9037	0.02	98.80																																														
30	0.00009	0.8821	0.08	98.50																																														
40	0.00015	0.8751	0.13	98.40																																														
50	0.00021	0.8514	0.18	98.05																																														
60	0.00027	0.8319	0.24	97.75																																														
70	0.00037	0.8163	0.32	97.50																																														
80	0.00039	0.8011	0.34	97.25																																														
90	0.00046	0.7865	0.40	97.00																																														
<p>AUXILIARY INFORMATION</p>																																																		
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The titration method was used at constant temperature. No further details were reported in the paper.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified; distilled; without impurities by glc, b.p. $185.3^\circ C$, n_D^{20} 1.412.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <p>REFERENCES:</p>																																																	

COMPONENTS: (1) Propanoic acid hexyl ester (hexyl propionate); $C_9H_{18}O_2$; [2445-76-3] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u> , <i>31</i> , 56-70.																																																											
VARIABLES: T/K = 273 - 364	PREPARED BY: Z. Maczynska																																																											
EXPERIMENTAL VALUES: Mutual solubility of propanoic acid hexyl ester and water <hr/> <table border="1" data-bbox="157 504 1145 1008"> <thead> <tr> <th rowspan="2">t/°C</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>0.029</td><td>99.61</td><td>0.000033</td><td>0.9667</td></tr> <tr><td>9.7</td><td>0.022</td><td>99.66</td><td>0.000025</td><td>0.9709</td></tr> <tr><td>20.0</td><td>0.024</td><td>99.66</td><td>0.000027</td><td>0.9709</td></tr> <tr><td>29.6</td><td>0.012</td><td>99.55</td><td>0.000014</td><td>0.9618</td></tr> <tr><td>39.8</td><td>0.012</td><td>99.52</td><td>0.000014</td><td>0.9593</td></tr> <tr><td>50.0</td><td>0.014</td><td>99.54</td><td>0.000016</td><td>0.9610</td></tr> <tr><td>60.0</td><td>0.016</td><td>99.50</td><td>0.000018</td><td>0.9577</td></tr> <tr><td>70.3</td><td>0.018</td><td>99.51</td><td>0.000020</td><td>0.9585</td></tr> <tr><td>80.3</td><td>0.017</td><td>99.49</td><td>0.000019</td><td>0.9569</td></tr> <tr><td>90.6</td><td>0.018</td><td>99.17</td><td>0.000020</td><td>0.9315</td></tr> </tbody> </table> <hr/> std. dev. 0.001 0.02		t/°C	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	0.029	99.61	0.000033	0.9667	9.7	0.022	99.66	0.000025	0.9709	20.0	0.024	99.66	0.000027	0.9709	29.6	0.012	99.55	0.000014	0.9618	39.8	0.012	99.52	0.000014	0.9593	50.0	0.014	99.54	0.000016	0.9610	60.0	0.016	99.50	0.000018	0.9577	70.3	0.018	99.51	0.000020	0.9585	80.3	0.017	99.49	0.000019	0.9569	90.6	0.018	99.17	0.000020	0.9315
t/°C	g(1)/100g sln		x_1 (compiler)																																																									
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																								
0	0.029	99.61	0.000033	0.9667																																																								
9.7	0.022	99.66	0.000025	0.9709																																																								
20.0	0.024	99.66	0.000027	0.9709																																																								
29.6	0.012	99.55	0.000014	0.9618																																																								
39.8	0.012	99.52	0.000014	0.9593																																																								
50.0	0.014	99.54	0.000016	0.9610																																																								
60.0	0.016	99.50	0.000018	0.9577																																																								
70.3	0.018	99.51	0.000020	0.9585																																																								
80.3	0.017	99.49	0.000019	0.9569																																																								
90.6	0.018	99.17	0.000020	0.9315																																																								
AUXILIARY INFORMATION																																																												
METHOD/APPARATUS/PROCEDURE: The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial sample; purity 99%; used as received. (2) Not specified. <hr/> ESTIMATED ERROR: Accuracy of method 0.1 wt% or less, for solubility, see above. <hr/> REFERENCES:																																																											

COMPONENTS: (1) Dibutylphosphinic acid methyl ester (methyl dibutylphosphinate); $C_9H_{21}O_2P$; [7163-67-9] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Nikolaev, A.V.; Dyadin, Yu.A.; Yakovlev, I.I.; Durasov, V.B.; Yakovleva, N.I.; Khol'kina, I.D. Zh. Fiz. Khim. 1966, 40, 221-3.																																																																					
VARIABLES: $T/K = 309 - 464$	PREPARED BY: A. Skrzecz																																																																					
EXPERIMENTAL VALUES: Mutual solubility of dibutylphosphinic acid methyl ester and water																																																																						
<table border="1"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr> <td>35.5 LCST</td> <td>19.83</td> <td>19.83</td> <td>0.02265</td> <td>0.02265</td> </tr> <tr> <td>36.2</td> <td>12.75</td> <td>-</td> <td>0.01351</td> <td>-</td> </tr> <tr> <td>36.6</td> <td>-</td> <td>31.05</td> <td>-</td> <td>0.04049</td> </tr> <tr> <td>37.0</td> <td>-</td> <td>36.87</td> <td>-</td> <td>0.05189</td> </tr> <tr> <td>38.0</td> <td>-</td> <td>42.02</td> <td>-</td> <td>0.06360</td> </tr> <tr> <td>40.2</td> <td>7.40</td> <td>-</td> <td>0.00743</td> <td>-</td> </tr> <tr> <td>41.0</td> <td>-</td> <td>46.14</td> <td>-</td> <td>0.07431</td> </tr> <tr> <td>47.2</td> <td>-</td> <td>52.49</td> <td>-</td> <td>0.09382</td> </tr> <tr> <td>50.0</td> <td>4.85</td> <td>-</td> <td>0.00475</td> <td>-</td> </tr> <tr> <td>56.3</td> <td>-</td> <td>57.58</td> <td>-</td> <td>0.11285</td> </tr> <tr> <td>58.0</td> <td>3.79</td> <td>-</td> <td>0.00368</td> <td>-</td> </tr> <tr> <td>60.0</td> <td>3.65</td> <td>-</td> <td>0.00354</td> <td>-</td> </tr> </tbody> </table>		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	35.5 LCST	19.83	19.83	0.02265	0.02265	36.2	12.75	-	0.01351	-	36.6	-	31.05	-	0.04049	37.0	-	36.87	-	0.05189	38.0	-	42.02	-	0.06360	40.2	7.40	-	0.00743	-	41.0	-	46.14	-	0.07431	47.2	-	52.49	-	0.09382	50.0	4.85	-	0.00475	-	56.3	-	57.58	-	0.11285	58.0	3.79	-	0.00368	-	60.0	3.65	-	0.00354	-
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																																			
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																																		
35.5 LCST	19.83	19.83	0.02265	0.02265																																																																		
36.2	12.75	-	0.01351	-																																																																		
36.6	-	31.05	-	0.04049																																																																		
37.0	-	36.87	-	0.05189																																																																		
38.0	-	42.02	-	0.06360																																																																		
40.2	7.40	-	0.00743	-																																																																		
41.0	-	46.14	-	0.07431																																																																		
47.2	-	52.49	-	0.09382																																																																		
50.0	4.85	-	0.00475	-																																																																		
56.3	-	57.58	-	0.11285																																																																		
58.0	3.79	-	0.00368	-																																																																		
60.0	3.65	-	0.00354	-																																																																		
(continued next page)																																																																						
AUXILIARY INFORMATION																																																																						
METHOD/APPARATUS/PROCEDURE: The Alekseev's synthetic method was used. A water or glycerine bath was used for temperature control. Ester hydrolysis was not observed. No further details were reported in the paper.	SOURCE AND PURITY OF MATERIALS: (1) Synthesized; distilled 5-6 times under vacuum; d_4^{25} 0.9501, n_D^{25} 1.4441. (2) Not specified.																																																																					
ESTIMATED ERROR: Not specified.																																																																						
REFERENCES:																																																																						

<p>COMPONENTS:</p> <p>(1) Dibutylphosphinic acid methyl ester (methyl dibutylphosphinate); $C_9H_{21}O_2P$; [7163-67-9]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Nikolaev, A.V.; Dyadin, Yu.A.; Yakovlev, I.I.; Durasov, V.B.; Yakovleva, N.I.; Khol'kina, I.D. <i>Zh. Fiz. Khim.</i> <u>1966</u>, <i>40</i>, 221-3.</p>
-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

EXPERIMENTAL VALUES: (continued)

Mutual solubility of dibutylphosphinic acid methyl ester and water

$t/^\circ C$	g(1)/100g sln		x_1 (compiler)	
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase
64.0	3.46	-	0.00335	-
68.0	3.29	-	0.00318	-
71.0	-	62.20	-	0.13360
90.5	-	65.14	-	0.14902
120.0	3.29	-	0.00318	-
133.0	-	65.14	-	0.14902
150.0	4.85	-	0.00475	-
155.0	-	62.20	-	0.13360
171.0	-	57.58	-	0.11285
172.0	7.40	-	0.00743	-
181.0	-	52.49	-	0.09382
186.0	12.75	-	0.01351	-
190.0	19.83	42.02	0.02265	0.06360
191.0 UCST	31.05	31.05	0.04049	0.04049

<p>COMPONENTS:</p> <p>(1) Methylphosphonic acid dibutyl ester (dibutyl methylphosphonate); $C_9H_{21}O_3P$; [2404-73-1]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Nikolaev, A.V.; Dyadin, Yu.A.; Yakovlev, I.I.; Durasov, V.B.; Yakovleva, N.I.; Khol'kina, I.D. <i>Zh. Fiz. Khim.</i> <u>1967</u>, 41, 1815-7.</p>																																																																
<p>VARIABLES:</p> <p>$T/K = 275 - 348$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>																																																																
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of methylphosphonic acid dibutyl ester and water</p> <table border="1" data-bbox="243 592 1237 1137"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">$g(1)/100g\ sln$</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>1.6</td><td>-</td><td>60.82</td><td>-</td><td>0.11840</td></tr> <tr><td>4.3</td><td>3.49</td><td>-</td><td>0.00312</td><td>-</td></tr> <tr><td>7.5</td><td>3.02</td><td>-</td><td>0.00269</td><td>-</td></tr> <tr><td>8.4</td><td>-</td><td>64.88</td><td>-</td><td>0.13780</td></tr> <tr><td>8.7</td><td>2.94</td><td>-</td><td>0.00261</td><td>-</td></tr> <tr><td>18.2</td><td>-</td><td>69.20</td><td>-</td><td>0.16274</td></tr> <tr><td>19.7</td><td>2.00</td><td>-</td><td>0.00176</td><td>-</td></tr> <tr><td>28.0</td><td>-</td><td>72.09</td><td>-</td><td>0.18265</td></tr> <tr><td>34.0</td><td>1.49</td><td>-</td><td>0.00131</td><td>-</td></tr> <tr><td>44.5</td><td>-</td><td>75.30</td><td>-</td><td>0.20870</td></tr> <tr><td>75.0</td><td>-</td><td>78.34</td><td>-</td><td>0.23833</td></tr> </tbody> </table>		$t/^\circ C$	$g(1)/100g\ sln$		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	1.6	-	60.82	-	0.11840	4.3	3.49	-	0.00312	-	7.5	3.02	-	0.00269	-	8.4	-	64.88	-	0.13780	8.7	2.94	-	0.00261	-	18.2	-	69.20	-	0.16274	19.7	2.00	-	0.00176	-	28.0	-	72.09	-	0.18265	34.0	1.49	-	0.00131	-	44.5	-	75.30	-	0.20870	75.0	-	78.34	-	0.23833
$t/^\circ C$	$g(1)/100g\ sln$		x_1 (compiler)																																																														
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																													
1.6	-	60.82	-	0.11840																																																													
4.3	3.49	-	0.00312	-																																																													
7.5	3.02	-	0.00269	-																																																													
8.4	-	64.88	-	0.13780																																																													
8.7	2.94	-	0.00261	-																																																													
18.2	-	69.20	-	0.16274																																																													
19.7	2.00	-	0.00176	-																																																													
28.0	-	72.09	-	0.18265																																																													
34.0	1.49	-	0.00131	-																																																													
44.5	-	75.30	-	0.20870																																																													
75.0	-	78.34	-	0.23833																																																													
<p>AUXILIARY INFORMATION</p>																																																																	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The Alekseev's synthetic method was used. No further details were reported in the paper.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Synthesized; distilled 5-6 times under vacuum; d_4^{20} 0.9788, n_D^{20} 1.4255.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <p>REFERENCES:</p>																																																																

<p>COMPONENTS:</p> <p>(1) 1,2-Benzenedicarboxylic acid dimethyl ester (dimethyl phthalate); $C_{10}H_{10}O_4$; [131-11-13]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988</p>
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION:

Quantitative solubility data for the 1,2-benzenedicarboxylic acid dimethyl ester (1) - water (2) system have been reported in the publications listed in Table 1.

TABLE 1: Quantitative Solubility Studies of the 1,2-Benzenedicarboxylic acid dimethyl ester (1) - Water (2) System

Reference	T/K	Solubility	Method
Shanley and Greenspan (ref 1)	RT ^a	(2) in (1)	unspecified
Leyder and Boulanger (ref 2)	293	(1) in (2)	GLC, spectrophotometric

^a RT - room temperature.

As can be seen from Table 1, there are no independent studies for solubilities in either the H_2O -rich or organic-rich phases and so no Critical Evaluation is possible. The interested user is referred to the relevant Data Sheets immediately following this Evaluation for the experimental data.

REFERENCES

1. Shanley, E. S.; Greenspan, F. P. *Ind. Eng. Chem.* 1947, *39*, 1536-43.
2. Leyder, F.; Boulanger, P. *Bull. Environ. Contam. Toxicol.* 1983, *30*, 152-7.

<p>COMPONENTS:</p> <p>(1) 1,2-Benzenedicarboxylic acid dimethyl ester (dimethyl phthalate); $C_{10}H_{10}O_4$; [131-11-13]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Shanley, E.S.; Greenspan, F.P. <i>Ind. Eng. Chem.</i> <u>1947</u>, 39, 1536-43.</p>
<p>VARIABLES:</p> <p>T/K = room temperature</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of water in 1,2-benzenedicarboxylic acid dimethyl ester at room temperature was reported to be 1.6 g(2)/100g(1). The corresponding mass per cent and mole fraction, x_2, values calculated by the compiler are 1.6 g(2)/100g sln and 0.15.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The method was not specified.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Not specified. (2) Not specified.</p> <hr/> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <hr/> <p>REFERENCES:</p>

<p>COMPONENTS:</p> <p>(1) 1,2-Benzenedicarboxylic acid dimethyl ester (dimethyl phthalate); $C_{10}H_{10}O_4$; [131-11-13]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Leyder, F.; Boulanger, P. <i>Bull. Environ. Contam. Toxicol.</i> <u>1983</u>, 30(2), 152-7.</p>
<p>VARIABLES:</p> <p>$T/K = 293$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of 1,2-benzenedicarboxylic acid dimethyl ester in water at 20°C was reported to be 2.21×10^{-2} mol(1)/L and 4.290 g(1)/L.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method as described in the <i>OECD Guidelines for Testing of Chemicals</i> (ref 1) was used. Gas chromatography analyses were conducted on a Girdel 3000 FFLE apparatus with flame-ionization detection. A 4 mm x 180 m stainless steel column packed with 3% OV-1 on 80-100 mesh Chromosorb W-AW-DMCS and helium at a 20mL/min flow rate were used. Samples were fortified with an internal standard, extracted, and concentrated to 1 mL of hexane solution. "Solvent-flush" and "hot-needle" injection techniques were used. The two methods gave similar results. Peak heights were measured throughout. UV measurements were made by using 1-cm cells in a Perkin-Elmer model 552 uv-Visible spectrophotometer.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Fluka; purity >99% ; used as received.</p> <p>(2) Deionized and distilled from $KMnO_4$.</p> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm 1^\circ C$.</p> <p>REFERENCES:</p> <p>1. <i>OECD Guidelines for Testing of Chemicals</i>, Paris, OECD, <u>1981</u>. (methods 101, 105).</p>

<p>COMPONENTS:</p> <p>(1) Hexanedioic acid diethyl ester (<i>diethyl adipate</i>); $C_{10}H_{18}O_4$; [141-28-6]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988</p>
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION:

Quantitative solubility data for the hexanedioic acid diethyl ester (1) - water (2) system have been reported in the publications listed in Table 1.

TABLE 1: Quantitative Solubility Studies of the
Hexanedioic acid diethyl ester (1) - Water (2) System

Reference	T/K	Solubility	Method
Sobotka and Kahn (ref 1)	293	(1) in (2)	titration
Gross et al. (ref 2)	303	(1) in (2)	interferometric
Stephenson and Stuart (ref 3)	273-363	mutual	GLC

The original data in these publications are compiled in the Data Sheets immediately following this Critical Evaluation. For convenience, further discussion of this system will be divided into two parts.

1. SOLUBILITY OF HEXANEDIOIC ACID DIETHYL ESTER (1) IN WATER (2)

All the available data for the solubility of hexanedioic acid diethyl ester (1) in water (2) are summarized in Table 2. At 293 and 303 K, the only temperatures where comparison is possible, the studies of Sobotka and Kahn (ref 1) and Gross et al. (ref 2) respectively, are in only fair agreement with that of Stephenson and Stuart (ref 3). At other temperatures only the data of Stephenson and Stuart (ref 3) are available. Consequently, all data must be considered Tentative pending further studies although it may be noted that the data of Stephenson and Stuart (ref 3) are generally reliable.

(continued next page)

COMPONENTS: (1) Hexanedioic acid diethyl ester (diethyl adipate); C ₁₀ H ₁₈ O ₄ ; [141-28-6] (2) Water; H ₂ O; [7732-18-5]	EVALUATOR: G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988
-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

TABLE 2: Tentative Solubilities
of Hexanedioic acid diethyl ester (1) in Water (2)

T/K	Solubilities		
	Reported values	"Best" values ^a	
	g(1)/100g sln	g(1)/100g sln	10 ⁴ x ₁
273	1.07 (ref 3)	1.0	9
283	0.87* (ref 3)	0.9	8
293	0.422 (ref 1), 0.69* (ref 3)	0.6 ± 0.1 ^b	5
298	0.64* (ref 3)	0.6	5
303	0.424 (ref 2), 0.60* (ref 3)	0.5 ± 0.1 ^b	4
313	0.54* (ref 3)	0.5	4
323	0.51* (ref 3)	0.5	4
333	0.50* (ref 3)	0.5	4
343	0.50* (ref 3)	0.5	4
353	0.51* (ref 3)	0.5	4
363	0.55* (ref 3)	0.6	5

^a Rounded values of ref 3 unless otherwise indicated.

^b Average value; x₁ values have the same (relative) percentage uncertainty as the mass % solubilities.

2. SOLUBILITY OF WATER (2) IN HEXANEDIOIC ACID DIETHYL ESTER (1)

As the only available data for the solubility of water (2) in hexanedioic acid diethyl ester (1) are those of Stephenson and Stuart (ref 3) no Critical Evaluation is possible. The interested user is referred to the relevant Data Sheet for the experimental solubilities but it may be noted that the data of these authors are generally reliable (but see Part 1. above).

REFERENCES

- Sobotka, H.; Kahn, J. *J. Am. Chem. Soc.* 1931, *53*, 2935-8.
- Gross, P. M.; Saylor, J. H.; Gorman, M. A. *J. Am. Chem. Soc.* 1953, *55*, 650-2.
- Stephenson, R.; Stuart, J. *J. Chem. Eng. Data* 1986, *31*, 56-70.

<p>COMPONENTS:</p> <p>(1) Hexanedioic acid diethyl ester (diethyl adipate); $C_{10}H_{18}O_4$; [141-28-6]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Sobotka, H.; Kahn, J. <i>J. Am. Chem. Soc.</i> <u>1931</u>, 53, 2935-8.</p>
<p>VARIABLES:</p> <p>$T/K = 293$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of hexanedioic acid diethyl ester in water at 20°C was reported to be 0.423 g(1)/100mL(2). The corresponding mass per cent and mole fraction, x_1, values calculated by the compiler are 0.422 g(1)/100g sln and 3.77×10^{-4}.</p> <p>Density of water $d_4^{20} = 0.9982$ (ref 1) was used in the calculation.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The titration method was used. The ester was added dropwise from a micro-burette with a capillary tip to 100, 250, or 500 mL of water in a narrow-mouthed stock bottle with a well-ground glass stopper. The bottle was shaken after each addition of ester. 1-5 mg of Sudan IV dye was put into the water to improve the end-point of the titration. At saturation, one additional drop of ester was sufficient to convert the floating rough indicator particles into dark transparent droplets.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified (Eastman Kodak Laboratories or synthesized); twice-distilled under reduced pressure; d_4^{20} 1.0086, n_D^{20} 1.4278.</p> <p>(2) Distilled.</p> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <p>REFERENCES:</p> <p>1. <i>Selected Values of Properties of Hydrocarbons and Related Compounds</i>, API Research Project 44, Thermodynamics Research Center, Texas A and M University, Texas, <u>1973</u>.</p>

COMPONENTS: (1) Hexanedioic acid diethyl ester (diethyl adipate); $C_{10}H_{18}O_4$; [141-28-6] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Gross, P.M.; Saylor, J.H.; Gorman, M.A. J. Am. Chem. Soc. <u>1933</u> , 55, 650-2.
VARIABLES: $T/K = 303$	PREPARED BY: Z. Maczynska
EXPERIMENTAL VALUES: The solubility of hexanedioic acid diethyl ester in water at 30°C was reported to be 4.26 g(1)/kg(2) and 0.0211 mol(1)/kg(2). The corresponding mass per cent and mole fraction, x_1 , values calculated by the compiler are 0.424 g(1)/100g sln and 3.79×10^{-4} .	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The analytical method was used. The saturated solutions of (1) in (2) were prepared in thin-walled cylindrical flask in a large water thermostat. The saturated solutions were analyzed by means of a Zeiss combination liquid and gas interferometer. The procedure has been previously described in detail by Gross and Saylor (ref 1).	SOURCE AND PURITY OF MATERIALS: (1) Eastman Kodak Co. best grade; distilled; b.p. range 129.0-129.4°C. (2) Distilled. ESTIMATED ERROR: Temp. $\pm 0.02^\circ C$. Soly. $\pm 0.5\%$ (from the concordance of duplicate determinations). REFERENCES: 1. Gross, P.M.; Saylor, J.H. J. Am. Chem. Soc. <u>1931</u> , 53, 1744.

COMPONENTS: (1) Hexanedioic acid diethyl ester (diethyl adipate); $C_{10}H_{18}O_4$; [141-28-6] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u> , 31, 56-70.																																																											
VARIABLES: T/K = 273 - 363	PREPARED BY: Z. Maczynska																																																											
EXPERIMENTAL VALUES: Mutual solubility of hexanedioic acid diethyl ester and water <hr/> <table border="1" data-bbox="249 539 1233 1064"> <thead> <tr> <th rowspan="2">t/°C</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr> <td>0</td> <td>1.07</td> <td>98.92</td> <td>0.00096</td> <td>0.8908</td> </tr> <tr> <td>9.8</td> <td>1.26</td> <td>98.89</td> <td>0.00113</td> <td>0.8881</td> </tr> <tr> <td>20.5</td> <td>0.69</td> <td>-</td> <td>0.00062</td> <td>-</td> </tr> <tr> <td>39.5</td> <td>0.54</td> <td>98.05</td> <td>0.00048</td> <td>0.8175</td> </tr> <tr> <td>50.0</td> <td>0.52</td> <td>97.89</td> <td>0.00046</td> <td>0.8051</td> </tr> <tr> <td>60.1</td> <td>0.50</td> <td>97.62</td> <td>0.00045</td> <td>0.7851</td> </tr> <tr> <td>69.7</td> <td>0.50</td> <td>97.56</td> <td>0.00045</td> <td>0.7808</td> </tr> <tr> <td>80.2</td> <td>0.49</td> <td>97.17</td> <td>0.00045</td> <td>0.7536</td> </tr> <tr> <td>90.3</td> <td>0.56</td> <td>97.12</td> <td>0.00050</td> <td>0.7502</td> </tr> <tr> <td>std. dev.</td> <td>0.01</td> <td>0.03</td> <td></td> <td></td> </tr> </tbody> </table>		t/°C	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	1.07	98.92	0.00096	0.8908	9.8	1.26	98.89	0.00113	0.8881	20.5	0.69	-	0.00062	-	39.5	0.54	98.05	0.00048	0.8175	50.0	0.52	97.89	0.00046	0.8051	60.1	0.50	97.62	0.00045	0.7851	69.7	0.50	97.56	0.00045	0.7808	80.2	0.49	97.17	0.00045	0.7536	90.3	0.56	97.12	0.00050	0.7502	std. dev.	0.01	0.03		
t/°C	g(1)/100g sln		x_1 (compiler)																																																									
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																								
0	1.07	98.92	0.00096	0.8908																																																								
9.8	1.26	98.89	0.00113	0.8881																																																								
20.5	0.69	-	0.00062	-																																																								
39.5	0.54	98.05	0.00048	0.8175																																																								
50.0	0.52	97.89	0.00046	0.8051																																																								
60.1	0.50	97.62	0.00045	0.7851																																																								
69.7	0.50	97.56	0.00045	0.7808																																																								
80.2	0.49	97.17	0.00045	0.7536																																																								
90.3	0.56	97.12	0.00050	0.7502																																																								
std. dev.	0.01	0.03																																																										
AUXILIARY INFORMATION																																																												
METHOD/APPARATUS/PROCEDURE: <p>The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.</p>	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial sample; purity 99%; used as received. (2) Not specified. <hr/> ESTIMATED ERROR: Accuracy of method 0.1 wt% or less, for solubility, see above. <hr/> REFERENCES:																																																											

COMPONENTS: (1) Acetic acid 2-ethylhexyl ester (2-ethylhexyl acetate); $C_{10}H_{20}O_2$; [103-09-3] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Doolittle, A.K. <i>Ind. End. Chem.</i> <u>1935</u> , 27, 1169-79.
VARIABLES: $T/K = 293$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: <p>The solubility of acetic acid 2-ethylhexyl ester in water at 20°C was reported to be very slightly soluble.</p> <p>The solubility of water in acetic acid 2-ethylhexyl ester at 20°C was reported to be 0.39 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.036.</p>	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The method was not specified.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial product; purity 99%, b.p. range 195-203°C, d_4^{20} 0.873, n_D^{20} 1.4300. (2) Not specified.
	ESTIMATED ERROR: Not specified.
	REFERENCES:

COMPONENTS: (1) Acetic acid octyl ester (octyl acetate); $C_{10}H_{20}O_2$; [112-14-1] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Carbide and Carbon Chemicals Corporation, Solvent Chart. Othmer, D.F.; White, R.E.; Trueger, E. <i>Ind. Eng. Chem.</i> <u>1941</u> , 33, 1240-8, 1513.
VARIABLES: $T/K = 296$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: The solubility of acetic acid octyl ester in water at 23°C was reported to be less than 0.3 g(1)/100g sln. The corresponding mole fraction, x_1 , value calculated by the compiler is less than 3×10^{-5} . The solubility of water in acetic acid octyl ester at 23°C was reported to be 0.55 g(2)/100g sln. The corresponding mole fraction, x_2 , value calculated by the compiler is 0.050.	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: Nothing was specified in the paper.	SOURCE AND PURITY OF MATERIALS: (1) Not specified. (2) Not specified. ESTIMATED ERROR: Not specified. REFERENCES:

COMPONENTS: (1) Nonanoic acid methyl ester (methyl nonanoate); $C_{10}H_{20}O_2$; [1731-84-6] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Tewari, Y.B.; Miller, M.M.; Wasik, S.P.; Martire, D.E. <i>J. Chem. Eng. Data</i> <u>1982</u> , 27, 451-4.
VARIABLES: $T/K = 298$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: The solubility of nonanoic acid methyl ester in water at 25°C was reported to be 1.33×10^{-4} mol(1)/L sln. The corresponding value on a mass/volume basis is 0.0229 g(1)/L sln (compiler).	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The analytical method was used. The aqueous phase was generated by pumping water into the inlet of coated generator column which was thermostated, either by using a minipump or by means of water reservoir using compressed air at 5 psi. The aqueous solution was extracted by the use of a known amount of immiscible with water solution and then analyzed by a gas chromatographic technique.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified; purity >99% by glc. (2) Not specified.
	ESTIMATED ERROR: Temp. $\pm 0.1^\circ C$. Soly. $\pm 1.0\%$.
	REFERENCES:

<p>COMPONENTS:</p> <p>(1) Octanoic acid ethyl ester (ethyl caprylate); $C_{10}H_{20}O_2$; [106-32-1]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988</p>
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION:

Quantitative solubility data for the octanoic acid ethyl ester (1) - water (2) system have been reported in the publications listed in Table 1.

TABLE 1: Quantitative Solubility Studies of the Octanoic acid ethyl ester (1) - Water (2) System

Reference	T/K	Solubility	Method
Sobotka and Kahn (ref 1)	293	(1) in (2)	titration
Stephenson and Stuart (ref 2)	283-364	mutual	GLC

The original data in these publications are compiled in the Data Sheets immediately following this Critical Evaluation. For convenience, further discussion of this system will be divided into two parts.

1. SOLUBILITY OF OCTANOIC ACID ETHYL ESTER (1) IN WATER (2)

All the available data for the solubility of octanoic acid ethyl ester (1) in water (2) are summarized in Table 2. At 293 K, the only temperature where comparison is possible, the data of Sobotka and Kahn (ref 1) and Stephenson and Stuart (ref 2) disagree by almost an order of magnitude (Table 2). In the absence of other independent studies it is not possible to determine which value is more reliable. At other temperatures, only the data of Stephenson and Stuart (ref 2) are available.

Although the solubilities reported by these authors are usually reliable it should be noted that their results for this system are very scattered. For these reasons the data in Table 2 should be regarded as very Tentative pending further studies.

(continued next page)

COMPONENTS: (1) Octanoic acid ethyl ester (ethyl caprylate); $C_{10}H_{20}O_2$; [106-32-1] (2) Water; H_2O ; [7732-18-5]	EVALUATOR: G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988
------------------------------------------------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

TABLE 2: Tentative Solubilities
of Octanoic acid ethyl ester (1) in Water (2)

T/K	Solubilities		
	Reported values	"Best" values ^a	
	g(1)/100g sln	g(1)/100g sln	$10^5 x_1$
283	0.05* (ref 2)	0.05	5
293	0.007 (ref 1), 0.05 (ref 2)	0.05	5
298	0.04* (ref 2)	0.04	4
303	0.04* (ref 2)	0.04	4
313	0.04* (ref 2)	0.04	4
323	0.04* (ref 2)	0.04	4
333	0.04* (ref 2)	0.04	4
343	0.04* (ref 2)	0.04	4
353	0.04* (ref 2)	0.04	4
363	0.04* (ref 2)	0.04	4

^a Rounded values of ref 2, but see text.

2. SOLUBILITY OF WATER (2) IN OCTANOIC ACID ETHYL ESTER (1)

As the only available data for the solubility of water (2) in octanoic acid ethyl ester (1) are those of Stephenson and Stuart (ref 2) no Critical Evaluation is possible. The interested user is referred to the relevant Data Sheet for the experimental solubilities. However, it may be noted that the solubilities reported by these authors are generally reliable although the values reported for this particular system are rather scattered (see also Part 1. above).

REFERENCES

- Sobotka, H.; Kahn, J. *J. Am. Chem. Soc.* **1931**, *53*, 2935-8.
- Stephenson, R.; Stuart, J. *J. Chem. Eng. Data* **1986**, *31*, 56-70.

<p>COMPONENTS:</p> <p>(1) Octanoic acid ethyl ester (ethyl caprylate); $C_{10}H_{20}O_2$; [106-32-1]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Sobotka, H.; Kahn, J. <i>J. Am. Chem. Soc.</i> <u>1931</u>, 53, 2935-8.</p>
<p>VARIABLES:</p> <p>$T/K = 293$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of octanoic acid ethyl ester in water at 20°C was reported to be 0.007 g(1)/100mL(2). The corresponding mass per cent and mole fraction, x_1, values calculated by the compiler are 0.007 g(1)/100g sln and 7×10^{-6}.</p> <p>Density of water $d_4^{20} = 0.9982$ (ref 1) was used in the calculation.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The titration method was used. The ester was added dropwise from a micro-burette with a capillary tip to 100, 250, or 500 mL of water in a narrow-mouthed stock bottle with a well-ground glass stopper. The bottle was shaken after each addition of ester. 1-5 mg of Sudan IV dye was put into the water to improve the end-point of the titration. At saturation, one additional drop of ester was sufficient to convert the floating rough indicator particles into dark transparent droplets.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified (Eastman Kodak Laboratories or synthesized); twice-distilled under reduced pressure; $d_4^{20} 0.8674$, $n_D^{20} 1.4197$.</p> <p>(2) Distilled.</p> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <p>REFERENCES:</p> <p>1. <i>Selected Values of Properties of Hydrocarbons and Related Compounds</i>, API Research Project 44, Thermodynamics Research Center, Texas A and M University, Texas, <u>1973</u>.</p>

COMPONENTS: (1) Octanoic acid ethyl ester (ethyl caprylate); $C_{10}H_{20}O_2$; [106-32-1] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u> , <i>31</i> , 56-70.																																																											
VARIABLES: $T/K = 273 - 364$	PREPARED BY: Z. Maczynska																																																											
EXPERIMENTAL VALUES: Mutual solubility of octanoic acid ethyl ester and water <hr/> <table border="1" data-bbox="148 520 1126 1024"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>-</td><td>99.18</td><td>-</td><td>0.9267</td></tr> <tr><td>9.5</td><td>0.054</td><td>98.85</td><td>0.000056</td><td>0.8999</td></tr> <tr><td>19.6</td><td>0.051</td><td>99.10</td><td>0.000053</td><td>0.9201</td></tr> <tr><td>30.6</td><td>0.038</td><td>99.55</td><td>0.000040</td><td>0.9585</td></tr> <tr><td>39.5</td><td>-</td><td>99.61</td><td>-</td><td>0.9639</td></tr> <tr><td>50.0</td><td>0.037</td><td>99.35</td><td>0.000039</td><td>0.9411</td></tr> <tr><td>60.1</td><td>0.044</td><td>99.49</td><td>0.000046</td><td>0.9533</td></tr> <tr><td>70.2</td><td>0.040</td><td>99.57</td><td>0.000042</td><td>0.9603</td></tr> <tr><td>80.0</td><td>0.042</td><td>99.59</td><td>0.000044</td><td>0.9621</td></tr> <tr><td>90.6</td><td>-</td><td>99.61</td><td>-</td><td>0.9639</td></tr> </tbody> </table> <hr/> std. dev. 0.003 0.01		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	-	99.18	-	0.9267	9.5	0.054	98.85	0.000056	0.8999	19.6	0.051	99.10	0.000053	0.9201	30.6	0.038	99.55	0.000040	0.9585	39.5	-	99.61	-	0.9639	50.0	0.037	99.35	0.000039	0.9411	60.1	0.044	99.49	0.000046	0.9533	70.2	0.040	99.57	0.000042	0.9603	80.0	0.042	99.59	0.000044	0.9621	90.6	-	99.61	-	0.9639
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																									
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																								
0	-	99.18	-	0.9267																																																								
9.5	0.054	98.85	0.000056	0.8999																																																								
19.6	0.051	99.10	0.000053	0.9201																																																								
30.6	0.038	99.55	0.000040	0.9585																																																								
39.5	-	99.61	-	0.9639																																																								
50.0	0.037	99.35	0.000039	0.9411																																																								
60.1	0.044	99.49	0.000046	0.9533																																																								
70.2	0.040	99.57	0.000042	0.9603																																																								
80.0	0.042	99.59	0.000044	0.9621																																																								
90.6	-	99.61	-	0.9639																																																								
AUXILIARY INFORMATION																																																												
METHOD/APPARATUS/PROCEDURE: The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial sample; purity 99%; used as received. (2) Not specified. <hr/> ESTIMATED ERROR: Accuracy of method 0.1 wt% or less, for solubility, see above. <hr/> REFERENCES:																																																											

<p>COMPONENTS:</p> <p>(1) Propanoic acid, 2-methyl-, hexyl ester (hexyl isobutyrate); $C_{10}H_{20}O_2$; [2349-07-7]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u>, <i>31</i>, 56-70.</p>																																																																
<p>VARIABLES:</p> <p>$T/K = 273 - 364$</p>	<p>PREPARED BY:</p> <p>Z. Maczynska</p>																																																																
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of 2-methylpropanoic acid hexyl ester and water</p> <table border="1" data-bbox="244 594 1239 1098"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>0.0116</td><td>99.50</td><td>0.0000121</td><td>0.9541</td></tr> <tr><td>9.4</td><td>0.0061</td><td>99.69</td><td>0.0000064</td><td>0.9711</td></tr> <tr><td>19.5</td><td>0.0038</td><td>99.63</td><td>0.0000040</td><td>0.9657</td></tr> <tr><td>29.7</td><td>-</td><td>99.74</td><td>-</td><td>0.9757</td></tr> <tr><td>39.5</td><td>0.0032</td><td>-</td><td>0.0000033</td><td>-</td></tr> <tr><td>50.0</td><td>0.0076</td><td>99.56</td><td>0.0000079</td><td>0.9594</td></tr> <tr><td>59.8</td><td>0.0081</td><td>-</td><td>0.0000085</td><td>-</td></tr> <tr><td>70.1</td><td>-</td><td>99.60</td><td>-</td><td>0.9630</td></tr> <tr><td>80.0</td><td>0.0040</td><td>99.59</td><td>0.0000042</td><td>0.9621</td></tr> <tr><td>90.6</td><td>0.0087</td><td>99.58</td><td>0.0000091</td><td>0.9612</td></tr> <tr> <td>std. dev.</td> <td>0.0005</td> <td>0.02</td> <td></td> <td></td> </tr> </tbody> </table>		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	0.0116	99.50	0.0000121	0.9541	9.4	0.0061	99.69	0.0000064	0.9711	19.5	0.0038	99.63	0.0000040	0.9657	29.7	-	99.74	-	0.9757	39.5	0.0032	-	0.0000033	-	50.0	0.0076	99.56	0.0000079	0.9594	59.8	0.0081	-	0.0000085	-	70.1	-	99.60	-	0.9630	80.0	0.0040	99.59	0.0000042	0.9621	90.6	0.0087	99.58	0.0000091	0.9612	std. dev.	0.0005	0.02		
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																														
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																													
0	0.0116	99.50	0.0000121	0.9541																																																													
9.4	0.0061	99.69	0.0000064	0.9711																																																													
19.5	0.0038	99.63	0.0000040	0.9657																																																													
29.7	-	99.74	-	0.9757																																																													
39.5	0.0032	-	0.0000033	-																																																													
50.0	0.0076	99.56	0.0000079	0.9594																																																													
59.8	0.0081	-	0.0000085	-																																																													
70.1	-	99.60	-	0.9630																																																													
80.0	0.0040	99.59	0.0000042	0.9621																																																													
90.6	0.0087	99.58	0.0000091	0.9612																																																													
std. dev.	0.0005	0.02																																																															
<p>AUXILIARY INFORMATION</p>																																																																	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, commercial sample; purity 98%; used as received.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Accuracy of method 0.1 wt% or less, for solubility, see above.</p> <p>REFERENCES:</p>																																																																

COMPONENTS: (1) Acetic acid 2-(2-butoxyethoxy)ethyl ester (2-(2-butoxyethoxy)ethyl acetate); C ₁₀ H ₂₀ O ₄ ; [124-17-4] (2) Water; H ₂ O; [7732-18-5]	EVALUATOR: G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988
-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION:

Quantitative solubility data for the acetic acid 2-(2-butoxyethoxy)ethyl ester (1) - water (2) system have been reported in the publications listed in Table 1.

TABLE 1: Quantitative Solubility Studies of the Acetic acid 2-(2-butoxyethoxy)ethyl ester (1) - Water (2) System

Reference	T/K	Solubility	Method
Doolittle (ref 1)	293	mutual	unspecified
Othmer et al. (ref 2)	297	mutual	unspecified

The original data in these publications are compiled in the Data Sheets immediately following this Critical Evaluation. They are also summarized in Tables 2 and 3 below.

As there have been only two independent studies at slightly different temperatures, Critical Evaluation is difficult. If the temperature difference is assumed to be negligible (ester-water mutual solubilities do not generally vary dramatically ca. 295 K) then the data for the H₂O-rich phase are in poor agreement whilst the ester-rich phase values are in good agreement. The latter may be regarded as Tentative.

TABLE 2: Reported Solubilities of Acetic acid 2-(2-butoxyethoxy)ethyl ester (1) in Water (2)

T/K	Reported Solubilities	
	g(1)/100g sln	10 ³ x ₁
293	1.6 (ref 1)	1.4
297	6.5 (ref 2)	6.1

(continued next page)

<p>COMPONENTS:</p> <p>(1) Acetic acid 2-(2-butoxyethoxy)ethyl ester (2-(2-butoxyethoxy)ethyl acetate); C₁₀H₂₀O₄; [124-17-4]</p> <p>(2) Water; H₂O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988</p>									
<p>CRITICAL EVALUATION: (continued)</p>										
<p style="text-align: center;"><u>TABLE 3: Tentative Solubilities of Water (2) in Acetic acid 2-(2-butoxyethoxy)ethyl ester (1)</u></p>										
<p>T/K</p>	<p style="text-align: center;">Tentative Solubilities</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="width: 15%;"></th> <th style="width: 45%; text-align: center;">g(2)/100g sln</th> <th style="width: 40%; text-align: center;">x₂</th> </tr> </thead> <tbody> <tr> <td style="text-align: center;">293</td> <td style="text-align: center;">3.4 (ref 1)</td> <td style="text-align: center;">0.29</td> </tr> <tr> <td style="text-align: center;">297</td> <td style="text-align: center;">3.7 (ref 2)</td> <td style="text-align: center;">0.30</td> </tr> </tbody> </table>		g(2)/100g sln	x ₂	293	3.4 (ref 1)	0.29	297	3.7 (ref 2)	0.30
	g(2)/100g sln	x ₂								
293	3.4 (ref 1)	0.29								
297	3.7 (ref 2)	0.30								
<p>REFERENCES</p>										
<p>1. Doolittle, A. K. <i>Ind. Eng. Chem.</i> <u>1935</u>, 27, 1169-79.</p> <p>2. Othmer, D. F.; White, R. E.; Trueger, E. <i>Ind. Eng. Chem.</i> <u>1941</u>, 33, 1240-8; <i>ibid.</i> 1513; see also Carbide and Carbon Chemical Corporation, Solvent Chart, before 1941.</p>										

COMPONENTS: (1) Acetic acid 2-(2-butoxyethoxy)ethyl ester (2-(2-butoxyethoxy)ethyl acetate); C ₁₀ H ₂₀ O ₄ ; [124-17-4] (2) Water; H ₂ O; [7732-18-5]	ORIGINAL MEASUREMENTS: Doolittle, A.K. <i>Ind. End. Chem.</i> <u>1935</u> , 27, 1169-79.
VARIABLES: T/K = 293	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: The solubility of acetic acid 2-(2-butoxyethoxy)ethyl ester in water at 20°C was reported to be 1.6 g(1)/100g sln. The corresponding mole fraction, x_1 , value calculated by the compiler is 0.0014. The solubility of water in acetic acid 2-(2-butoxyethoxy)ethyl ester at 20°C was reported to be 3.4 g(2)/100g sln. The corresponding mole fraction, x_2 , value calculated by the compiler is 0.29.	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The method was not specified.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial product; purity 99%, b.p. range 236-249°C, d_4^{20} 0.987. (2) Not specified.
	ESTIMATED ERROR: Not specified.
	REFERENCES:

<p>COMPONENTS:</p> <p>(1) Acetic acid 2-(2-butoxyethoxy)ethyl ester (2-(2-butoxyethoxy)ethyl acetate); C₁₀H₂₀O₄; [124-17-4]</p> <p>(2) Water; H₂O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Carbide and Carbon Chemicals Corporation, Solvent Chart.</p> <p>Othmer, D.F.; White, R.E.; Trueger, E.</p> <p><i>Ind. Eng. Chem.</i> <u>1941</u>, 33, 1240-8, 1513.</p>
<p>VARIABLES:</p> <p>T/K = 297</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of acetic acid 2-(2-butoxyethoxy)ethyl ester in water at 23.5°C was reported to be 6.5 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 0.00609.</p> <p>The solubility of water in acetic acid 2-(2-butoxyethoxy)ethyl ester at 23.5°C was reported to be 3.7 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.303.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>Nothing was specified in the paper.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Not specified.</p> <p>(2) Not specified.</p> <hr/> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm 0.5^\circ\text{C}$ (mean of reported range).</p> <hr/> <p>REFERENCES:</p>

COMPONENTS: (1) Dibutylphosphinic acid ethyl ester (ethyl dibutylphosphinate); $C_{10}H_{23}O_2P$; [7100-92-7] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Nikolaev, A.V.; Dyadin, Yu.A.; Yakovlev, I.I.; Durasov, V.B.; Yakovleva, N.I.; Khol'kina, I.D. Zh. Fiz. Khim. <u>1966</u> , 40, 221-3.																																																																					
VARIABLES: $T/K = 287 - 599$	PREPARED BY: A. Skrzecz																																																																					
EXPERIMENTAL VALUES: Mutual solubility of dibutylphosphinic acid ethyl ester and water <hr/> <table border="1" data-bbox="124 566 1126 1130"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>14.0 LCST</td><td>23.36</td><td>23.36</td><td>0.02593</td><td>0.02593</td></tr> <tr><td>14.6</td><td>-</td><td>34.57</td><td>-</td><td>0.04411</td></tr> <tr><td>15.4</td><td>9.99</td><td>-</td><td>0.00960</td><td>-</td></tr> <tr><td>18.0</td><td>-</td><td>46.94</td><td>-</td><td>0.07172</td></tr> <tr><td>18.6</td><td>5.96</td><td>-</td><td>0.00550</td><td>-</td></tr> <tr><td>22.5</td><td>-</td><td>53.17</td><td>-</td><td>0.09022</td></tr> <tr><td>23.4</td><td>4.43</td><td>-</td><td>0.00403</td><td>-</td></tr> <tr><td>26.5</td><td>3.88</td><td>-</td><td>0.00351</td><td>-</td></tr> <tr><td>28.0</td><td>-</td><td>58.65</td><td>-</td><td>0.11023</td></tr> <tr><td>29.5</td><td>3.58</td><td>-</td><td>0.00323</td><td>-</td></tr> <tr><td>41.0</td><td>2.44</td><td>-</td><td>0.00218</td><td>-</td></tr> <tr><td>52.0</td><td>2.02</td><td>-</td><td>0.00180</td><td>-</td></tr> </tbody> </table> <p style="text-align: right;">(continued next page)</p>		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	14.0 LCST	23.36	23.36	0.02593	0.02593	14.6	-	34.57	-	0.04411	15.4	9.99	-	0.00960	-	18.0	-	46.94	-	0.07172	18.6	5.96	-	0.00550	-	22.5	-	53.17	-	0.09022	23.4	4.43	-	0.00403	-	26.5	3.88	-	0.00351	-	28.0	-	58.65	-	0.11023	29.5	3.58	-	0.00323	-	41.0	2.44	-	0.00218	-	52.0	2.02	-	0.00180	-
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																																			
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																																		
14.0 LCST	23.36	23.36	0.02593	0.02593																																																																		
14.6	-	34.57	-	0.04411																																																																		
15.4	9.99	-	0.00960	-																																																																		
18.0	-	46.94	-	0.07172																																																																		
18.6	5.96	-	0.00550	-																																																																		
22.5	-	53.17	-	0.09022																																																																		
23.4	4.43	-	0.00403	-																																																																		
26.5	3.88	-	0.00351	-																																																																		
28.0	-	58.65	-	0.11023																																																																		
29.5	3.58	-	0.00323	-																																																																		
41.0	2.44	-	0.00218	-																																																																		
52.0	2.02	-	0.00180	-																																																																		
AUXILIARY INFORMATION																																																																						
METHOD/APPARATUS/PROCEDURE: The Alekseev's synthetic method was used. A water or glycerine bath was used for temperature control. Ester hydrolysis was not observed. No further details were reported in the paper.	SOURCE AND PURITY OF MATERIALS: (1) Synthesized; distilled 5-6 times under vacuum; d_4^{20} 0.9394, n_D^{20} 1.4420. (2) Not specified.																																																																					
	ESTIMATED ERROR: Not specified.																																																																					
	REFERENCES:																																																																					

COMPONENTS:		ORIGINAL MEASUREMENTS:		
(1) Dibutylphosphinic acid ethyl ester (ethyl dibutylphosphinate); $C_{10}H_{23}O_2P$; [7100-92-7]		Nikolaev, A.V.; Dyadin, Yu.A.; Yakovlev, I.I.; Durasov, V.B.; Yakovleva, N.I.; Khol'kina, I.D.		
(2) Water; H_2O ; [7732-18-5]		<i>Zh. Fiz. Khim.</i> 1966, 40, 221-3.		
EXPERIMENTAL VALUES: (continued)				
Mutual solubility of dibutylphosphinic acid ethyl ester and water				
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)	
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase
54.8	-	70.28	-	0.17118
58.0	1.79	-	0.00159	-
89.0	-	74.91	-	0.20683
127.0	1.79	-	0.00159	-
159.0	2.44	-	0.00218	-
168.0	-	70.28	-	0.17118
189.0	4.43	-	0.00403	-
211.0	-	58.65	-	0.11023
216.0	9.99	-	0.00960	-
217.0	-	53.17	-	0.09022
223.0	-	46.94	-	0.07172
225.0	-	34.57	-	0.04411
226.0 UCST	23.36	23.36	0.02593	0.02593

<p>COMPONENTS:</p> <p>(1) Dipropylphosphinic acid butyl ester (butyl dipropylphosphinate); $C_{10}H_{23}O_2P$; [16984-10-4]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Nikolaev, A.V.; Dyadin, Yu.A.; Yakovlev, I.I.; Durasov, V.B.; Yakovleva, N.I.; Khol'kina, I.D.</p> <p>Zh. Fiz. Khim. 1967, 41, 1815-7.</p>																																																											
<p>VARIABLES:</p> <p>$T/K = 273 - 448$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>																																																											
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of dipropylphosphinic acid butyl ester and water</p> <table border="1" data-bbox="116 564 1118 1068"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr> <td>0.0</td> <td>3.3^b</td> <td>13.1^b</td> <td>0.003^a</td> <td>0.013^a</td> </tr> <tr> <td>0.3</td> <td>-</td> <td>14.36</td> <td>-</td> <td>0.01443</td> </tr> <tr> <td>2.4</td> <td>3.12</td> <td>-</td> <td>0.002805</td> <td>-</td> </tr> <tr> <td>3.15</td> <td>-</td> <td>15.76</td> <td>-</td> <td>0.01608</td> </tr> <tr> <td>4.2</td> <td>-</td> <td>16.62</td> <td>-</td> <td>0.01711</td> </tr> <tr> <td>5.8</td> <td>2.57</td> <td>-</td> <td>0.002299</td> <td>-</td> </tr> <tr> <td>7.4</td> <td>-</td> <td>19.70</td> <td>-</td> <td>0.02098</td> </tr> <tr> <td>8.3</td> <td>-</td> <td>19.95</td> <td>-</td> <td>0.02130</td> </tr> <tr> <td>13.6</td> <td>-</td> <td>30.08</td> <td>-</td> <td>0.03621</td> </tr> <tr> <td>14.2</td> <td>1.88</td> <td>-</td> <td>0.001671</td> <td>-</td> </tr> </tbody> </table> <p>^a Experimental value. ^b Calculated by the compiler.</p> <p style="text-align: right;">(continued next page)</p>		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0.0	3.3 ^b	13.1 ^b	0.003 ^a	0.013 ^a	0.3	-	14.36	-	0.01443	2.4	3.12	-	0.002805	-	3.15	-	15.76	-	0.01608	4.2	-	16.62	-	0.01711	5.8	2.57	-	0.002299	-	7.4	-	19.70	-	0.02098	8.3	-	19.95	-	0.02130	13.6	-	30.08	-	0.03621	14.2	1.88	-	0.001671	-
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																									
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																								
0.0	3.3 ^b	13.1 ^b	0.003 ^a	0.013 ^a																																																								
0.3	-	14.36	-	0.01443																																																								
2.4	3.12	-	0.002805	-																																																								
3.15	-	15.76	-	0.01608																																																								
4.2	-	16.62	-	0.01711																																																								
5.8	2.57	-	0.002299	-																																																								
7.4	-	19.70	-	0.02098																																																								
8.3	-	19.95	-	0.02130																																																								
13.6	-	30.08	-	0.03621																																																								
14.2	1.88	-	0.001671	-																																																								
<p>AUXILIARY INFORMATION</p>																																																												
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The Alekseev's synthetic method was used. No further details were reported in the paper.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Synthesized; distilled 5-6 times under vacuum; d_4^{20} 0.9372, n_D^{20} 1.4420.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <p>REFERENCES:</p>																																																											

COMPONENTS:			ORIGINAL MEASUREMENTS:	
(1) Dipropylphosphinic acid butyl ester (butyl dipropylphosphinate); $C_{10}H_{23}O_2P$; [16984-10-4]			Nikolaev, A.V.; Dyadin, Yu.A.; Yakovlev, I.I.; Durasov, V.B.; Yakovleva, N.I.; Khol'kina, I.D.	
(2) Water; H_2O ; [7732-18-5]			Zh. Fiz. Khim. 1967, 41, 1815-7.	
EXPERIMENTAL VALUES: (continued)				
Mutual solubility of dipropylphosphinic acid butyl ester and water				
t/°C	g(1)/100g sln		x_1 (compiler)	
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase
17.2	1.69	-	0.001499	-
18.1	-	39.70	-	0.05438
20.8	1.56	-	0.001382	-
23.9	-	49.44	-	0.07869
33.8	1.04	-	0.000917	-
34.0	-	59.62	-	0.11423
53.0	0.651	-	0.000572	-
57.0	-	69.98	-	0.16916
60.7	-	70.76	-	0.17448
89.8	-	74.49	-	0.20321
139.0	-	74.49	-	0.20321
175.0	-	70.76	-	0.17448

<p>COMPONENTS:</p> <p>(1) Ethylphosphonic acid dibutyl ester (dibutyl ethylphosphonate); $C_{10}H_{23}O_3P$; [2404-58-2]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Nikolaev, A.V.; Dyadin, Yu.A.; Yakovlev, I.I.; Durasov, V.B.; Yakovleva, N.I.; Khol'kina, I.D.</p> <p>Zh. Fiz. Khim. 1967, 41, 1815-7.</p>																																																											
<p>VARIABLES:</p> <p>$T/K = 278 - 346$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>																																																											
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of ethylphosphonic acid dibutyl ester and water</p> <table border="1" data-bbox="137 574 1125 1078"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr> <td>5.2</td> <td>-</td> <td>73.28</td> <td>-</td> <td>0.1819</td> </tr> <tr> <td>8.6</td> <td>1.51</td> <td>-</td> <td>0.001241</td> <td>-</td> </tr> <tr> <td>9.8</td> <td>-</td> <td>74.87</td> <td>-</td> <td>0.1945</td> </tr> <tr> <td>16.2</td> <td>-</td> <td>77.06</td> <td>-</td> <td>0.2140</td> </tr> <tr> <td>19.5</td> <td>1.02</td> <td>-</td> <td>0.000835</td> <td>-</td> </tr> <tr> <td>29.5</td> <td>-</td> <td>79.94</td> <td>-</td> <td>0.2441</td> </tr> <tr> <td>34.5</td> <td>0.703</td> <td>-</td> <td>0.000574</td> <td>-</td> </tr> <tr> <td>44.4</td> <td>-</td> <td>83.31</td> <td>-</td> <td>0.2880</td> </tr> <tr> <td>53.0</td> <td>0.504</td> <td>-</td> <td>0.000410</td> <td>-</td> </tr> <tr> <td>73.0</td> <td>-</td> <td>84.62</td> <td>-</td> <td>0.3084</td> </tr> </tbody> </table>		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	5.2	-	73.28	-	0.1819	8.6	1.51	-	0.001241	-	9.8	-	74.87	-	0.1945	16.2	-	77.06	-	0.2140	19.5	1.02	-	0.000835	-	29.5	-	79.94	-	0.2441	34.5	0.703	-	0.000574	-	44.4	-	83.31	-	0.2880	53.0	0.504	-	0.000410	-	73.0	-	84.62	-	0.3084
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																									
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																								
5.2	-	73.28	-	0.1819																																																								
8.6	1.51	-	0.001241	-																																																								
9.8	-	74.87	-	0.1945																																																								
16.2	-	77.06	-	0.2140																																																								
19.5	1.02	-	0.000835	-																																																								
29.5	-	79.94	-	0.2441																																																								
34.5	0.703	-	0.000574	-																																																								
44.4	-	83.31	-	0.2880																																																								
53.0	0.504	-	0.000410	-																																																								
73.0	-	84.62	-	0.3084																																																								
<p>AUXILIARY INFORMATION</p>																																																												
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The Alekseev's synthetic method was used. No further details were reported in the paper.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Synthesized; distilled 5-6 times under vacuum; d_4^{20} 0.9658, n_D^{20} 1.4290.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <p>REFERENCES:</p>																																																											

<p>COMPONENTS:</p> <p>(1) Heptanedioic acid diethyl ester (<i>diethyl heptanedioate</i>); $C_{11}H_{20}O_4$; [2050-20-6]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Sobotka, H.; Kahn, J. <i>J. Am. Chem. Soc.</i> <u>1931</u>, 53, 2935-8.</p>
<p>VARIABLES:</p> <p>$T/K = 293$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of heptanedioic acid diethyl ester in water at 20°C was reported to be 0.199 g(1)/100mL(2). The corresponding mass per cent and mole fraction, x_1, values calculated by the compiler are 0.199 g(1)/100g sln and 1.66×10^{-4}.</p> <p>Density of water $d_4^{20} = 0.9982$ (ref 1) was used in the calculation.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The titration method was used. The ester was added dropwise from a micro-burette with a capillary tip to 100, 250, or 500 mL of water in a narrow-mouthed stock bottle with a well-ground glass stopper. The bottle was shaken after each addition of ester. 1-5 mg of Sudan IV dye was put into the water to improve the end-point of the titration. At saturation, one additional drop of ester was sufficient to convert the floating rough indicator particles into dark transparent droplets.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified (Eastman Kodak Laboratories or synthesized); twice-distilled under reduced pressure; d_4^{20} 0.9945, n_D^{20} 1.4303.</p> <p>(2) Distilled.</p> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <p>REFERENCES:</p> <p>1. <i>Selected Values of Properties of Hydrocarbons and Related Compounds</i>, API Research Project 44, Thermodynamics Research Center, Texas A and M University, Texas, <u>1973</u>.</p>

COMPONENTS: (1) Acetic acid isononyl ester (isononyl acetate); $C_{11}H_{22}O_2$; [40379-24-6] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Stephenson, R.; Stuart, J. <i>J. Chem. Eng. Data</i> <u>1986</u> , 31, 56-70.																																																											
VARIABLES: T/K = 273 - 364	PREPARED BY: Z. Maczynska																																																											
EXPERIMENTAL VALUES: Mutual solubility of acetic acid isononyl ester and water <hr/> <table border="1" data-bbox="137 504 1125 1008"> <thead> <tr> <th rowspan="2">t/°C</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0</td><td>0.018</td><td>99.666</td><td>0.000017</td><td>0.96651</td></tr> <tr><td>9.2</td><td>0.023</td><td>99.687</td><td>0.000022</td><td>0.96855</td></tr> <tr><td>19.4</td><td>0.020</td><td>99.708</td><td>0.000019</td><td>0.97061</td></tr> <tr><td>29.6</td><td>0.018</td><td>99.628</td><td>0.000017</td><td>0.96282</td></tr> <tr><td>39.8</td><td>0.021</td><td>99.608</td><td>0.000020</td><td>0.96089</td></tr> <tr><td>49.8</td><td>0.021</td><td>99.629</td><td>0.000020</td><td>0.96292</td></tr> <tr><td>60.3</td><td>0.013</td><td>99.631</td><td>0.000012</td><td>0.96311</td></tr> <tr><td>70.2</td><td>0.018</td><td>99.613</td><td>0.000017</td><td>0.96138</td></tr> <tr><td>80.3</td><td>0.026</td><td>99.637</td><td>0.000025</td><td>0.96369</td></tr> <tr><td>90.5</td><td>0.021</td><td>99.598</td><td>0.000020</td><td>0.95993</td></tr> </tbody> </table> <hr/> std. dev. 0.002 0.003		t/°C	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0	0.018	99.666	0.000017	0.96651	9.2	0.023	99.687	0.000022	0.96855	19.4	0.020	99.708	0.000019	0.97061	29.6	0.018	99.628	0.000017	0.96282	39.8	0.021	99.608	0.000020	0.96089	49.8	0.021	99.629	0.000020	0.96292	60.3	0.013	99.631	0.000012	0.96311	70.2	0.018	99.613	0.000017	0.96138	80.3	0.026	99.637	0.000025	0.96369	90.5	0.021	99.598	0.000020	0.95993
t/°C	g(1)/100g sln		x_1 (compiler)																																																									
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																								
0	0.018	99.666	0.000017	0.96651																																																								
9.2	0.023	99.687	0.000022	0.96855																																																								
19.4	0.020	99.708	0.000019	0.97061																																																								
29.6	0.018	99.628	0.000017	0.96282																																																								
39.8	0.021	99.608	0.000020	0.96089																																																								
49.8	0.021	99.629	0.000020	0.96292																																																								
60.3	0.013	99.631	0.000012	0.96311																																																								
70.2	0.018	99.613	0.000017	0.96138																																																								
80.3	0.026	99.637	0.000025	0.96369																																																								
90.5	0.021	99.598	0.000020	0.95993																																																								
AUXILIARY INFORMATION																																																												
METHOD/APPARATUS/PROCEDURE: The analytical method was used. Component (1) was equilibrated with component (2) at a given temperature in a thermostat. Each layer was sampled with a syringe; (1) was determined by adding a weighed amount of acetonitrile (or sometimes propanol) to the organic layer sample and measuring by a Gow-Mac thermal conductivity gc the (1)/acetonitrile peak ratio (Chromosorb 101 packing and a HP 3390 A recorder-integrator). A similar procedure but a higher boiling material (e.g. 1-hexanol) was used to determine (2) in the water layer.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, commercial sample; purity 98%; used as received. (2) Not specified. <hr/> ESTIMATED ERROR: Accuracy of method 0.1 wt% or less, for solubility, see above. <hr/> REFERENCES:																																																											

<p>COMPONENTS:</p> <p>(1) Decanoic acid methyl ester (methyl decanoate); $C_{11}H_{22}O_2$; [110-42-9]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Tewari, Y.B.; Miller, M.M.; Wasik, S.P.; Martire, D.E.</p> <p><i>J. Chem. Eng. Data</i> <u>1982</u>, <i>27</i>, 451-4.</p>
<p>VARIABLES:</p> <p>$T/K = 298$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of decanoic acid methyl ester in water at 25°C was reported to be 2.05×10^{-5} mol(1)/L sln. The corresponding value on mass/volume basis is 0.00382 g(1)/L sln (compiler).</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method was used. The aqueous phase was generated by pumping water into the inlet of a coated generator column which was thermostated, either by using a minipump or by means of a water reservoir using compressed air at 5 psi. The aqueous solution was extracted by the use of a known amount of immiscible-with-water solution and then analyzed by a gas chromatographic technique.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified; purity >99% by glc.</p> <p>(2) Not specified.</p> <hr/> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm 0.1^\circ C$. Soly. $\pm 1.0\%$.</p> <hr/> <p>REFERENCES:</p>

COMPONENTS: (1) Nonanoic acid ethyl ester (ethyl nonanoate); $C_{11}H_{22}O_2$; [123-29-5] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Sobotka, H.; Kahn, J. <i>J. Am. Chem. Soc.</i> <u>1931</u> , <i>53</i> , 2935-8.
VARIABLES: $T/K = 293$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: The solubility of Nonanoic acid ethyl ester in water at 20°C was reported to be 0.003 g(1)/100mL(2). The corresponding mass per cent and mole fraction, x_1 , values calculated by the compiler are 0.003 g(1)/100g sln and 3×10^{-6} . Density of water $d_4^{20} = 0.9982$ (ref 1) was used in the calculation.	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The titration method was used. The ester was added dropwise from a micro-burette with a capillary tip to 100, 250, or 500 mL of water in a narrow-mouthed stock bottle with a well-ground glass stopper. The bottle was shaken after each addition of ester. 1-5 mg of Sudan IV dye was put into the water to improve the end-point of the titration. At saturation, one additional drop of ester was sufficient to convert the floating rough indicator particles into dark transparent droplets.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified (Eastman Kodak Laboratories or synthesized); twice-distilled under reduced pressure; $n_D^{20} 1.4223$. (2) Distilled. ESTIMATED ERROR: Not specified. REFERENCES: 1. <i>Selected Values of Properties of Hydrocarbons and Related Compounds</i> , API Research Project 44, Thermodynamics Research Center, Texas A and M University, Texas, <u>1973</u> .

<p>COMPONENTS:</p> <p>(1) Dibutylphosphinic acid propyl ester (propyl dibutylphosphinate); $C_{11}H_{25}O_2P$; [7100-93-8]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Nikolaev, A.V.; Dyadin, Yu.A.; Yakovlev, I.I.; Durasov, V.B.; Yakovleva, N.I.; Khol'kina, I.D.</p> <p>Zh. Fiz. Khim. <u>1966</u>, 40, 221-3.</p>																																																																					
<p>VARIABLES:</p> <p>$T/K = 279 - 516$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>																																																																					
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of dibutylphosphinic acid propyl ester and water</p> <table border="1" data-bbox="230 580 1239 1139"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>6.1</td><td>-</td><td>65.22</td><td>-</td><td>0.13296</td></tr> <tr><td>10.0</td><td>3.01</td><td>-</td><td>0.00253</td><td>-</td></tr> <tr><td>13.7</td><td>-</td><td>69.63</td><td>-</td><td>0.15789</td></tr> <tr><td>15.2</td><td>-</td><td>70.48</td><td>-</td><td>0.16336</td></tr> <tr><td>20.5</td><td>2.01</td><td>-</td><td>0.00167</td><td>-</td></tr> <tr><td>27.5</td><td>-</td><td>74.76</td><td>-</td><td>0.19499</td></tr> <tr><td>29.0</td><td>1.50</td><td>-</td><td>0.00124</td><td>-</td></tr> <tr><td>29.8</td><td>-</td><td>75.31</td><td>-</td><td>0.19964</td></tr> <tr><td>41.0</td><td>1.01</td><td>-</td><td>0.00083</td><td>-</td></tr> <tr><td>64.0</td><td>0.75</td><td>-</td><td>0.00062</td><td>-</td></tr> <tr><td>66.5</td><td>-</td><td>80.58</td><td>-</td><td>0.25336</td></tr> <tr><td>143.0</td><td>-</td><td>80.58</td><td>-</td><td>0.25336</td></tr> </tbody> </table> <p style="text-align: right;">(continued next page)</p>		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	6.1	-	65.22	-	0.13296	10.0	3.01	-	0.00253	-	13.7	-	69.63	-	0.15789	15.2	-	70.48	-	0.16336	20.5	2.01	-	0.00167	-	27.5	-	74.76	-	0.19499	29.0	1.50	-	0.00124	-	29.8	-	75.31	-	0.19964	41.0	1.01	-	0.00083	-	64.0	0.75	-	0.00062	-	66.5	-	80.58	-	0.25336	143.0	-	80.58	-	0.25336
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																																			
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																																		
6.1	-	65.22	-	0.13296																																																																		
10.0	3.01	-	0.00253	-																																																																		
13.7	-	69.63	-	0.15789																																																																		
15.2	-	70.48	-	0.16336																																																																		
20.5	2.01	-	0.00167	-																																																																		
27.5	-	74.76	-	0.19499																																																																		
29.0	1.50	-	0.00124	-																																																																		
29.8	-	75.31	-	0.19964																																																																		
41.0	1.01	-	0.00083	-																																																																		
64.0	0.75	-	0.00062	-																																																																		
66.5	-	80.58	-	0.25336																																																																		
143.0	-	80.58	-	0.25336																																																																		
<p>AUXILIARY INFORMATION</p>																																																																						
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The Alekseev's synthetic method was used. A water or glycerine bath was used for temperature control. No further details were reported in the paper.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Synthesized; distilled 5-6 times under vacuum; d_4^{20} 0.9298, n_D^{20} 1.4439.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <p>REFERENCES:</p>																																																																					

<p>COMPONENTS:</p> <p>(1) Dibutylphosphinic acid propyl ester (propyl dibutylphosphinate); $C_{11}H_{25}O_2P$; [7100-93-8]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Nikolaev, A.V.; Dyadin, Yu.A.; Yakovlev, I.I.; Durasov, V.B.; Yakovleva, N.I.; Khol'kina, I.D.</p> <p>Zh. Fiz. Khim. 1966, 40, 221-3.</p>
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

EXPERIMENTAL VALUES: (continued)

Mutual solubility of dibutylphosphinic acid propyl ester and water

$t/^\circ C$	g(1)/100g sln		x_1 (compiler)	
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase
176.0	1.41	-	0.00117	-
189.0	-	75.31	-	0.19964
197.0	2.24	-	0.00187	-
208.0	-	70.48	-	0.16336
217.0	4.05	-	0.00344	-
221.0	-	65.22	-	0.13296
230.0	6.86	-	0.00599	-
242.0	17.09	50.32	0.01658	0.07650
243.0 UCST	35.08	35.08	0.04232	0.04232

The ester hydrolysis began at temperature 150-170°C.

<p>COMPONENTS:</p> <p>(1) Propylphosphonic acid dibutyl ester (dibutyl propylphosphonate); $C_{11}H_{25}O_3P$; [4628-12-0]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Nikolaev, A.V.; Dyadin, Yu.A.; Yakovlev, I.I.; Durasov, V.B.; Yakovleva, N.I.; Khol'kina, I.D. Zh. Fiz. Khim. <u>1967</u>, 41, 1815-7.</p>																																																											
<p>VARIABLES:</p> <p>$T/K = 274 - 316$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>																																																											
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of propylphosphonic acid dibutyl ester and water</p> <table border="1" data-bbox="249 609 1234 1120"> <thead> <tr> <th rowspan="2">t/°C</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr> <td>1.2</td> <td>-</td> <td>81.62</td> <td>-</td> <td>0.2529</td> </tr> <tr> <td>9.5</td> <td>-</td> <td>83.43</td> <td>-</td> <td>0.2774</td> </tr> <tr> <td>11.5</td> <td>0.596</td> <td>-</td> <td>4.569×10^{-4}</td> <td>-</td> </tr> <tr> <td>15.7</td> <td>0.446</td> <td>-</td> <td>3.414×10^{-4}</td> <td>-</td> </tr> <tr> <td>19.4</td> <td>0.386</td> <td>-</td> <td>2.953×10^{-4}</td> <td>-</td> </tr> <tr> <td>22.4</td> <td>-</td> <td>85.38</td> <td>-</td> <td>0.3081</td> </tr> <tr> <td>23.8</td> <td>0.324</td> <td>-</td> <td>2.478×10^{-4}</td> <td>-</td> </tr> <tr> <td>33.0</td> <td>-</td> <td>86.36</td> <td>-</td> <td>0.3256</td> </tr> <tr> <td>34.5</td> <td>0.230</td> <td>-</td> <td>1.757×10^{-4}</td> <td>-</td> </tr> <tr> <td>43.0</td> <td>-</td> <td>87.06</td> <td>-</td> <td>0.3390</td> </tr> </tbody> </table>		t/°C	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	1.2	-	81.62	-	0.2529	9.5	-	83.43	-	0.2774	11.5	0.596	-	4.569×10^{-4}	-	15.7	0.446	-	3.414×10^{-4}	-	19.4	0.386	-	2.953×10^{-4}	-	22.4	-	85.38	-	0.3081	23.8	0.324	-	2.478×10^{-4}	-	33.0	-	86.36	-	0.3256	34.5	0.230	-	1.757×10^{-4}	-	43.0	-	87.06	-	0.3390
t/°C	g(1)/100g sln		x_1 (compiler)																																																									
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																								
1.2	-	81.62	-	0.2529																																																								
9.5	-	83.43	-	0.2774																																																								
11.5	0.596	-	4.569×10^{-4}	-																																																								
15.7	0.446	-	3.414×10^{-4}	-																																																								
19.4	0.386	-	2.953×10^{-4}	-																																																								
22.4	-	85.38	-	0.3081																																																								
23.8	0.324	-	2.478×10^{-4}	-																																																								
33.0	-	86.36	-	0.3256																																																								
34.5	0.230	-	1.757×10^{-4}	-																																																								
43.0	-	87.06	-	0.3390																																																								
<p>AUXILIARY INFORMATION</p>																																																												
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The Alekseev's synthetic method was used. No further details were reported in the paper.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Synthesized; distilled 5-6 times under vacuum; d_4^{20} 0.9569, n_D^{20} 1.4309.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <p>REFERENCES:</p>																																																											

COMPONENTS: (1) 1,2-Benzenedicarboxylic acid diethyl ester (diethyl phthalate); $C_{12}H_{14}O_4$; [84-66-2] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Shanley, E.S.; Greenspan, F.P. <i>Ind. Eng. Chem.</i> <u>1947</u> , 39, 1536-43.
VARIABLES: T/K = room temperature	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: The solubility of water in 1,2-benzenedicarboxylic acid diethyl ester at room temperature was reported to be 1.0 g(2)/100g(1). The corresponding mass per cent and mole fraction, x_2 , values calculated by the compiler are 0.99 g(2)/100g sln and 0.11.	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The method was not specified. No further details were reported in the paper.	SOURCE AND PURITY OF MATERIALS: (1) Not specified. (2) Not specified.
	ESTIMATED ERROR: Not specified.
	REFERENCES:

<p>COMPONENTS:</p> <p>(1) 1,2-Benzenedicarboxylic acid diethyl ester (diethyl phthalate); $C_{12}H_{14}O_4$; [84-66-2]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Leyder, F.; Boulanger, P. <i>Bull. Environ. Contam. Toxicol.</i> <u>1983</u>, 30(2), 152-7.</p>
<p>VARIABLES:</p> <p>T/K = 293</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of 1,2-benzenedicarboxylic acid diethyl ester in water at 20°C was reported to be 4.18×10^{-3} mol(1)/L and 0.928 g(1)/L.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method as described in the <i>OECD Guidelines for Testing of Chemicals</i> (ref 1) was used. Gas chromatography analyses were conducted on a Girdel 3000 FFLE apparatus with flame-ionization detection. A 4 mm x 180 m stainless steel column packed with 3% OV-1 on 80-100 mesh Chromosorb W-AW-DMCS and helium at a 20mL/min flow rate were used. Samples were fortified with an internal standard, extracted, and concentrated to 1 mL of hexane solution. "Solvent flush" and "hot needle" injection techniques were used. The two methods gave similar results. Peak heights were measured throughout. UV measurements were made by using 1-cm cells in a Perkin-Elmer model 552 uv-Visible spectrophotometer.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Merck; purity 99% ; used as received.</p> <p>(2) Deionized and distilled from $KMnO_4$.</p> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm 1^\circ C$.</p> <p>REFERENCES:</p> <p>1. <i>OECD Guidelines for Testing of Chemicals</i>, Paris, OECD, <u>1981</u>. (methods 101, 105).</p>

<p>COMPONENTS:</p> <p>(1) Octanedioic acid diethyl ester (diethyl octanedioate); $C_{12}H_{22}O_4$; [2050-23-9]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Sobotka, H.; Kahn, J. <i>J. Am. Chem. Soc.</i> <u>1931</u>, 53, 2935-8.</p>
<p>VARIABLES:</p> <p>$T/K = 293$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of octanedioic acid diethyl ester in water at 20°C was reported to be 0.068 g(1)/100mL(2). The corresponding mass per cent and mole fraction, x_1, values calculated by the compiler are 0.068 g(1)/100g sln and 5.3×10^{-5}.</p> <p>Density of water $d_4^{20} = 0.9982$ (ref 1) was used in the calculation.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The titration method was used. The ester was added dropwise from a micro-burette with a capillary tip to 100, 250, or 500 mL of water in a narrow-mouthed stock bottle with a well-ground glass stopper. The bottle was shaken after each addition of ester. 1-5 mg of Sudan IV dye was put into the water to improve the end-point of the titration. At saturation, one additional drop of ester was sufficient to convert the floating rough indicator particles into dark transparent droplets.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified (Eastman Kodak Laboratories or synthesized); twice-distilled under reduced pressure; $d_4^{20} 0.9822$, $n_D^{20} 1.4334$.</p> <p>(2) Distilled.</p> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <p>REFERENCES:</p> <p>1. <i>Selected Values of Properties of Hydrocarbons and Related Compounds</i>, API Research Project 44, Thermodynamics Research Center, Texas A and M University, Texas, <u>1973</u>.</p>

<p>COMPONENTS:</p> <p>(1) Decanoic acid ethyl ester (ethyl decanoate); $C_{12}H_{24}O_2$; [110-38-3]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Sobotka, H.; Kahn, J. <i>J. Am. Chem. Soc.</i> <u>1931</u>, 53, 2935-8.</p>
<p>VARIABLES:</p> <p>$T/K = 293$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of decanoic acid ethyl ester in water at 20°C was reported to be 0.0015 g(1)/100mL(2). The corresponding mass per cent and mole fraction, x_1, values calculated by the compiler are 0.0015 g(1)/100g sln and 1.3×10^{-6}.</p> <p>Density of water $d_4^{20} = 0.9982$ (ref 1) was used in the calculation.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The titration method was used. The ester was added dropwise from a micro-burette with a capillary tip to 100, 250, or 500 mL of water in a narrow-mouthed stock bottle with a well-ground glass stopper. The bottle was shaken after each addition of ester. 1-5 mg of Sudan IV dye was put into the water to improve the end-point of the titration. At saturation, one additional drop of ester was sufficient to convert the floating rough indicator particles into dark transparent droplets.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified (Eastman Kodak Laboratories or synthesized); twice-distilled under reduced pressure; d_4^{20} 0.8637, n_D^{20} 1.4269.</p> <p>(2) Distilled.</p> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <p>REFERENCES:</p> <p>1. <i>Selected Values of Properties of Hydrocarbons and Related Compounds</i>, API Research Project 44, Thermodynamics Research Center, Texas A and M University, Texas, <u>1973</u>.</p>

<p>COMPONENTS:</p> <p>(1) Dibutylphosphinic acid butyl ester (butyl dibutylphosphinate); $C_{12}H_{27}O_2P$; [2950-47-2]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Nikolaev, A.V.; Dyadin, Yu.A.; Yakovlev, I.I.; Durasov, V.B.; Mironova, Z.N.</p> <p><i>Izv. Sib. Otd. Akad. Nauk SSSR,</i> <i>Ser. Khim. Nauk</i> <u>1965</u>, (3), 27-31.</p>																																																																					
<p>VARIABLES:</p> <p>$T/K = 273 - 429$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>																																																																					
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of dibutylphosphinic acid butyl ester and water</p> <hr/> <table border="1" data-bbox="134 580 1131 1118"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr> <td>0.</td> <td>-</td> <td>73.65^a</td> <td>-</td> <td>0.1769</td> </tr> <tr> <td>3.7</td> <td>1.30</td> <td>-</td> <td>10.12×10^{-4}</td> <td>-</td> </tr> <tr> <td>6.1</td> <td>-</td> <td>75.46</td> <td>-</td> <td>0.1912</td> </tr> <tr> <td>7.7</td> <td>1.03</td> <td>-</td> <td>8.00×10^{-4}</td> <td>-</td> </tr> <tr> <td>12.0</td> <td>-</td> <td>77.27</td> <td>-</td> <td>0.2072</td> </tr> <tr> <td>14.4</td> <td>0.716</td> <td>-</td> <td>5.54×10^{-4}</td> <td>-</td> </tr> <tr> <td>17.5</td> <td>-</td> <td>78.54</td> <td>-</td> <td>0.2196</td> </tr> <tr> <td>22.9</td> <td>0.490</td> <td>-</td> <td>3.78×10^{-4}</td> <td>-</td> </tr> <tr> <td>24.4</td> <td>-</td> <td>79.95</td> <td>-</td> <td>0.2346</td> </tr> <tr> <td>25.0</td> <td>0.49^b</td> <td>80.45^a</td> <td>3.78×10^{-4}</td> <td>0.2403</td> </tr> <tr> <td>25.0</td> <td>0.445</td> <td>-</td> <td>3.44×10^{-4}</td> <td>-</td> </tr> <tr> <td>26.5</td> <td>0.409</td> <td>-</td> <td>3.16×10^{-4}</td> <td>-</td> </tr> </tbody> </table> <p style="text-align: right;">(continued next page)</p>		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0.	-	73.65 ^a	-	0.1769	3.7	1.30	-	10.12×10^{-4}	-	6.1	-	75.46	-	0.1912	7.7	1.03	-	8.00×10^{-4}	-	12.0	-	77.27	-	0.2072	14.4	0.716	-	5.54×10^{-4}	-	17.5	-	78.54	-	0.2196	22.9	0.490	-	3.78×10^{-4}	-	24.4	-	79.95	-	0.2346	25.0	0.49 ^b	80.45 ^a	3.78×10^{-4}	0.2403	25.0	0.445	-	3.44×10^{-4}	-	26.5	0.409	-	3.16×10^{-4}	-
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																																			
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																																		
0.	-	73.65 ^a	-	0.1769																																																																		
3.7	1.30	-	10.12×10^{-4}	-																																																																		
6.1	-	75.46	-	0.1912																																																																		
7.7	1.03	-	8.00×10^{-4}	-																																																																		
12.0	-	77.27	-	0.2072																																																																		
14.4	0.716	-	5.54×10^{-4}	-																																																																		
17.5	-	78.54	-	0.2196																																																																		
22.9	0.490	-	3.78×10^{-4}	-																																																																		
24.4	-	79.95	-	0.2346																																																																		
25.0	0.49 ^b	80.45 ^a	3.78×10^{-4}	0.2403																																																																		
25.0	0.445	-	3.44×10^{-4}	-																																																																		
26.5	0.409	-	3.16×10^{-4}	-																																																																		
<p>AUXILIARY INFORMATION</p>																																																																						
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The synthetic method proposed by Aleksejev was used. Light filters and a high intensity light source were used to measure the cloudiness point with good accuracy. As a check, two analytical methods were used: the Karl Fischer method for water in the organic phase and the extraction of dibutylphosphinic acid butyl ester with diethyl ether, stripping, and drying over P_2O_5.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Synthesized and purified as given by Nikolaev, Dyadin, Yakovlev and Mironova (ref 1); b.p. $138^\circ C$ at 6 mm Hg, d_4^{20} 0.9262, n_D^{20} 1.4440.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm(0.2-0.6)^\circ C$ below $30^\circ C$, $\pm(1-3)^\circ C$ at high temperature.</p> <p>REFERENCES:</p> <p>1. Nikolaev, A.V.; Dyadin, Yu.A.; Yakovlev, I.I.; Mironova, Z.N. <i>Dokl. Akad. Nauk SSSR</i> <u>1963</u>, 153, 118.</p>																																																																					

<p>COMPONENTS:</p> <p>(1) Dibutylphosphinic acid butyl ester (butyl dibutylphosphinate); $C_{12}H_{27}O_2P$; [2950-47-2]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Nikolaev, A.V.; Dyadin, Yu.A.; Yakolvlev, I.I.; Durasov, V.B.; Mironova, Z.N.</p> <p>Izv. Sib. Otd. Akad. Nauk SSSR, Ser. Khim. Nauk <u>1965</u>, (3), 27-31.</p>
--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

EXPERIMENTAL VALUES: (continued)

Mutual solubility of dibutylphosphinic acid butyl ester and water

$t/^\circ C$	g(1)/100g sln		x_1 (compiler)	
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase
31.5	0.341	-	2.63×10^{-4}	-
32.5	-	81.19	-	0.2492
41.0	0.259	-	2.00×10^{-4}	-
43.2	-	82.38	-	0.2644
55.0	-	83.29	-	0.2771
57.5	0.200	-	1.54×10^{-4}	-
69.5	0.180	-	1.39×10^{-4}	-
70.7	-	84.12	-	0.2894
87.0	0.180	-	1.39×10^{-4}	-
124.0	-	84.12	-	0.2894
141.0	-	83.29	-	0.2771
156.0	-	82.38	-	0.2644

^a Karl Fischer method.

^b Extraction of dibutylphosphinic acid butyl ester with diethyl ether.

<p>COMPONENTS:</p> <p>(1) Dibutylphosphinic acid butyl ester (butyl dibutylphosphinate); $C_{12}H_{27}O_2P$; [2950-47-2]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Nikolaev, A.V.; Dyadin, Yu.A.; Yakovlev, I.I.; Durasov, V.B.; Yakovleva, N.I.; Khol'kina, I.D.</p> <p>Zh. Fiz. Khim. <u>1966</u>, 40, 221-3.</p>
<p>VARIABLES:</p> <p>$T/K = 526$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The UCST was reported to be 253°C.</p> <p>The dibutylphosphinic acid butyl ester hydrolysis began at temperature 150-170°C. The hydrolysis measured in the sample containing 19.89wt% of ester after heating at 260°C through 20 min was 5-10%.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The Alekseev's synthetic method was used. A water or glycerine bath was used for temperature control. No further details were reported in the paper.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Synthesized; distilled 5-6 times under vacuum; d_4^{25} 0.9179, n_D^{25} 1.4442.</p> <p>(2) Not specified.</p> <hr/> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <hr/> <p>REFERENCES:</p>

<p>COMPONENTS:</p> <p>(1) Butylphosphonic acid dibutyl ester (dibutyl butylphosphonate); $C_{12}H_{27}O_3P$; [78-46-6]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Nikolaev, A.V.; Dyadin, Yu.A.; Yakovlev, I.I.; Durasov, V.B.; Mironova, Z.N.</p> <p><i>Izv. Sib. Otd. Akad. Nauk SSSR, Ser. Khim. Nauk</i> <u>1965</u>, (3), 27-31.</p>																																																											
<p>VARIABLES:</p> <p>$T/K = 275 - 419$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>																																																											
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of butylphosphonic acid dibutyl ester and water</p> <table border="1" data-bbox="232 576 1240 1070"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr> <td>2.2</td> <td>-</td> <td>86.00</td> <td>-</td> <td>0.3066</td> </tr> <tr> <td>6.0</td> <td>0.247</td> <td>-</td> <td>17.82×10^{-5}</td> <td>-</td> </tr> <tr> <td>13.3</td> <td>-</td> <td>87.47</td> <td>-</td> <td>0.3344</td> </tr> <tr> <td>13.4</td> <td>0.168</td> <td>-</td> <td>12.11×10^{-5}</td> <td>-</td> </tr> <tr> <td>20.5</td> <td>-</td> <td>88.12</td> <td>-</td> <td>0.3480</td> </tr> <tr> <td>21.0</td> <td>0.121</td> <td>-</td> <td>8.72×10^{-5}</td> <td>-</td> </tr> <tr> <td>24.1</td> <td>-</td> <td>88.44</td> <td>-</td> <td>0.3551</td> </tr> <tr> <td>28.6</td> <td>0.0985</td> <td>-</td> <td>7.10×10^{-5}</td> <td>-</td> </tr> <tr> <td>35.0</td> <td>0.0853</td> <td>-</td> <td>6.14×10^{-5}</td> <td>-</td> </tr> <tr> <td>39.5</td> <td>-</td> <td>88.29</td> <td>-</td> <td>0.3750</td> </tr> </tbody> </table> <p style="text-align: right;">(continued next page)</p>		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	2.2	-	86.00	-	0.3066	6.0	0.247	-	17.82×10^{-5}	-	13.3	-	87.47	-	0.3344	13.4	0.168	-	12.11×10^{-5}	-	20.5	-	88.12	-	0.3480	21.0	0.121	-	8.72×10^{-5}	-	24.1	-	88.44	-	0.3551	28.6	0.0985	-	7.10×10^{-5}	-	35.0	0.0853	-	6.14×10^{-5}	-	39.5	-	88.29	-	0.3750
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																									
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																								
2.2	-	86.00	-	0.3066																																																								
6.0	0.247	-	17.82×10^{-5}	-																																																								
13.3	-	87.47	-	0.3344																																																								
13.4	0.168	-	12.11×10^{-5}	-																																																								
20.5	-	88.12	-	0.3480																																																								
21.0	0.121	-	8.72×10^{-5}	-																																																								
24.1	-	88.44	-	0.3551																																																								
28.6	0.0985	-	7.10×10^{-5}	-																																																								
35.0	0.0853	-	6.14×10^{-5}	-																																																								
39.5	-	88.29	-	0.3750																																																								
<p>AUXILIARY INFORMATION</p>																																																												
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The synthetic method proposed by Aleksejev was used. Light filters and a high intensity light source were used to measure the cloudiness point with good accuracy. No further details were reported in the paper.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Synthesized and purified as given by Nikolaev, Dyadin, Yakovlev and Mironova (ref 1); b.p. $133-5^\circ C$ at 6 mm Hg, d_4^{20} 0.9472, n_D^{20} 1.4323.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm(0.2-0.6)^\circ C$ below $30^\circ C$, $\pm(1-3)^\circ C$ at high temperature.</p> <p>REFERENCES:</p> <p>1. Nikolaev, A.V.; Dyadin, Yu.A.; Yakovlev, I.I.; Mironova, Z.N. <i>Dokl. Akad. Nauk SSSR</i> <u>1963</u>, 153, 118.</p>																																																											

COMPONENTS:

- (1) Butylphosphonic acid dibutyl ester
(dibutyl butylphosphonate);
 $C_{12}H_{27}O_3P$; [78-46-6]
(2) Water; H_2O ; [7732-18-5]

ORIGINAL MEASUREMENTS:

Nikolaev, A.V.; Dyadin, Yu.A.;
Yakolvlev, I.I.; Durasov, V.B.;
Mironova, Z.N.

Izv. Sib. Otd. Akad. Nauk SSSR,
Ser. Khim. Nauk 1965, (3),
27-31.

EXPERIMENTAL VALUES: (continued)

Mutual solubility of butylphosphonic acid dibutyl ester and water

t/°C	g(1)/100g sln		x_1 (compiler)	
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase
43.9	0.0688	-	4.95×10^{-5}	-
48.0	0.0591	-	4.26×10^{-5}	-
51.8	-	89.76	-	0.3868
96.0	-	89.76	-	0.3868
100.0	0.0591	-	4.26×10^{-5}	-
106.0	0.0688	-	4.95×10^{-5}	-
109.0	-	88.29	-	0.3750
126.0	-	88.44	-	0.3551
146.0	-	87.47	-	0.3344

<p>COMPONENTS:</p> <p>(1) Phosphoric acid tributyl ester (tributyl phosphate); $C_{12}H_{27}O_4P$; [126-73-8]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988</p>
---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION:

Quantitative solubility data for the phosphoric acid tributyl ester (1) - water (2) system have been reported in the publications listed in Table 1.

TABLE 1: Quantitative Solubility Studies of the
Phosphoric acid tributyl ester (1) - Water (2) System

Reference	T/K	Solubility	Method
Alcock et al. (ref 1)	289-298	mutual	analytical
Burger and Forsman (ref 2)	293	mutual	unspecified
Higgins et al. (ref 3)	277-323	(1) in (2)	analytical
Hasegawa (ref 4)	298	(1) in (2)	titration
Mikhailov et al. (ref 5)	279-299	(2) in (1)	Karl Fischer, synthetic
Hardy et al. (ref 6)	273-333	(2) in (1)	Karl Fischer
Nikolaev et al. (ref 7)	273-460	mutual	Karl Fischer, synthetic

The original data in these publications are compiled in the Data Sheets immediately following this Critical Evaluation. For convenience, further discussion of this system will be divided into two parts.

1. SOLUBILITY OF PHOSPHORIC ACID TRIBUTYL ESTER (1) IN WATER (2)

All the available data for the solubility of phosphoric acid tributyl ester (1) in water (2) are summarized in Table 2 with the following exceptions. The data of Alcock et al. (ref 1) are much lower, and that of Hasegawa (ref 4) much higher, than all other studies and have therefore been rejected.

Agreement among the remaining data (Table 2) is in general good and a number of the average "Best" values have been Recommended. At higher temperatures ($T > 323$ K) only the data of Nikolaev (ref 7) are available and must therefore be considered as Tentative. Selected data are plotted in Figure 1.

(continued next page)

COMPONENTS: (1) Phosphoric acid tributyl ester (tributyl phosphate); $C_{12}H_{27}O_4P$; [126-73-8] (2) Water; H_2O ; [7732-18-5]	EVALUATOR: G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988
---------------------------------------------------------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

TABLE 2: Recommended (R) and Tentative Solubilities
of Phosphoric acid tributyl ester (1) in Water (2)

T/K	Solubilities		
	Reported values	"Best" values ($\pm\sigma_n$) ^a	
	10^2 g(1)/100g sln	10^2 g(1)/100g sln 10^5x_1	
283	7.5* (ref 3), 7.5* (ref 7)	7.5 (R)	5.1
293	4.8* (ref 3), 4.6* (ref 7)	4.7 \pm 0.1 (R)	3.2
298	3.9 (ref 2), 4.2* (ref 3), 3.95* (ref 7)	4.0 \pm 0.1 (R)	2.7
303	3.8* (ref 3), 3.5* (ref 7)	3.7 \pm 0.2 (R)	2.5
313	3.4* (ref 3), 2.9* (ref 7)	3.2 \pm 0.3	2.2
323	2.9* (ref 3), 2.6* (ref 7)	2.8 \pm 0.2	1.9
333	2.5* (ref 7)	2.5	1.7
343	2.6* (ref 7)	2.6	1.8
353	2.9* (ref 7)	2.9	2.0
363	3.6* (ref 7)	3.6	2.5
373	5.0* (ref 7)	5.0	3.4

^a Obtained by averaging where appropriate; σ_n has no statistical significance. Mole fraction solubilities (x_1) have the same status and (relative) percentage uncertainties as the mass % solubilities.

2. SOLUBILITY OF WATER (2) IN PHOSPHORIC ACID TRIBUTYL ESTER (1)

All the available data for the solubility of water (2) in phosphoric acid tributyl ester (1) are summarized in Table 3 with the exception of the datum of Alcock et al. (ref 1) which is lower than all other studies and has therefore been rejected.

The remaining data are in good agreement (Table 3) enabling a number of average "Best" values to be Recommended. At higher temperatures ($T > 343$ K) only the data of Nikolaev (ref 7) are available and must therefore be considered as Tentative. Selected data are plotted in Figure 2.

(continued next page)

COMPONENTS:

- (1) Phosphoric acid tributyl ester
(*tributyl phosphate*);
 $C_{12}H_{27}O_4P$; [126-73-8]
- (2) Water; H_2O ; [7732-18-5]

EVALUATOR:

G.T. Hefter, School of Mathematical
and Physical Sciences,
Murdoch University, Perth, W.A.,
Australia
December, 1988

CRITICAL EVALUATION: (continued)




FIGURE 1. Selected data for the solubility of phosphoric acid tributyl ester (1) in water (2): ref 3 (●); ref 7 (O). Solid line is a least square polynomial fitted to the "Best" values from Table 2.

(continued next page)

COMPONENTS: (1) Phosphoric acid tributyl ester (tributyl phosphate); $C_{12}H_{27}O_4P$; [126-73-8] (2) Water; H_2O ; [7732-18-5]	EVALUATOR: G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988
---------------------------------------------------------------------------------------------------------------------------------------------------------	-------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

TABLE 3: Recommended (R) and Tentative Solubilities
of Water (2) in Phosphoric acid tributyl ester (1)

T/K	Solubilities		
	Reported values g(2)/100g sln	"Best" values ($\pm\sigma_n$) ^a g(2)/100g sln	x_2
273	7.09 (ref 6), 7.52 (ref 7)	7.3 \pm 0.2 (R)	0.54
283	6.38* (ref 5), 6.92* (ref 6), 7.05* (ref 7)	6.8 \pm 0.3 (R)	0.52
293	6.24* (ref 5), 6.73* (ref 6), 6.70* (ref 7)	6.6 \pm 0.2 (R)	0.51
298	6.24 (ref 5), 6.64 (ref 6), 6.56 (ref 7)	6.5 \pm 0.2 (R)	0.51
303	6.53* (ref 6), 6.45* (ref 7)	6.49 \pm 0.04 (R)	0.507
313	6.31* (ref 6), 6.40* (ref 7)	6.36 \pm 0.05 (R)	0.501
323	6.08 (ref 6), 6.37* (ref 7)	6.2 \pm 0.1 (R)	0.49
333	5.82 (ref 6), 6.40* (ref 7)	6.1 \pm 0.3 (R)	0.49
343	6.60* (ref 7)	6.6	0.51
353	6.85* (ref 7)	6.9	0.52
363	7.17* (ref 7)	7.2	0.53
373	7.55* (ref 7)	7.6	0.55
393	8.40* (ref 7)	8.4	0.58
413	9.3* (ref 7)	9	0.59
433	10.1* (ref 7)	10	0.62
453	10.9* (ref 7)	11	0.65

^a Obtained by averaging where appropriate; σ_n has no statistical significance. Mole fraction solubilities (x_2) have the same status and (relative) percentage uncertainties as the mass % solubilities.

(continued next page)

<p>COMPONENTS:</p> <p>(1) Phosphoric acid tributyl ester (tributyl phosphate); $C_{12}H_{27}O_4P$; [126-73-8]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988</p>
---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)




FIGURE 2. Selected data for the solubility of water (2) in phosphoric acid tributyl ester (1): ref 5 (X); ref 6 (●); ref 7 (O). Solid line is a least square polynomial fitted to the "Best" values from Table 3.

REFERENCES

- Alcock, K.; Grimley, S. S.; Healy, T. V.; Kennedy, J.; McKay, H. A. C. *Trans. Faraday Soc.* 1956, *52*, 39-47.
- Burger, L. L.; Forsman, R. C. Report HW-20936, 1951; quoted in Bruce, F. R. et al. *Progress in Nuclear Energy, Series 3, Process Chemistry*, Vol. 2, Pergamon Press, 1958, 546-56.
- Higgins, C. E.; Baldwin, W. H.; Soldano, B. A. *J. Phys. Chem.* 1959, *63*, 113-8.
- Hasegawa, T. *Kogyo Kagaku Zasshi* 1961, *64*, 1239-41.
- Mikhailov, V. A.; Kharachenko, S. K.; Nazin, A. G. *Izv. Sib. Otd. Akad. Nauk SSSR* 1962, (7), 50-6.
- Hardy, C. J.; Fairhurst, D.; McKay, H. A. C.; Willson, A. M. *Trans. Faraday Soc.* 1964, *60*, 1626-36.
- Nikolaev, A. V.; Dyadin, Yu. A.; Yakovlev, I. I.; Durasov, V. B.; Mironova, Z. N. *Izv. Sib. Otd. Akad. Nauk SSSR, Ser. Khim. Nauk* 1965, (3), 27-31.

<p>COMPONENTS:</p> <p>(1) Phosphoric acid tributyl ester (tributyl phosphate); $C_{12}H_{27}O_4P$; [126-73-8]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Alcock, K.; Grimley, S.S.; Healy, T.V.; Kennedy, J.; McKay, H.A.C. <i>Trans. Faraday Soc.</i> <u>1956</u>, 52, 39-47.</p>																								
<p>VARIABLES:</p> <p>$T/K = 289 - 298$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>																								
<p>EXPERIMENTAL VALUES:</p> <p>Solubility of phosphoric acid tributyl ester in water</p> <table border="1" data-bbox="161 520 865 808"> <thead> <tr> <th>$t/^\circ C$</th> <th>$g(1)/dm^3$</th> <th>$mol(1)/dm^3$ sln (compiler)</th> </tr> </thead> <tbody> <tr> <td>16.</td> <td>0.420</td> <td>0.001577</td> </tr> <tr> <td>17.</td> <td>0.410</td> <td>0.001540</td> </tr> <tr> <td>19.</td> <td>0.397</td> <td>0.001491</td> </tr> <tr> <td>22.</td> <td>0.380</td> <td>0.001427</td> </tr> </tbody> </table> <p>Solubility of water in phosphoric acid tributyl ester</p> <table border="1" data-bbox="161 909 865 1130"> <thead> <tr> <th>$t/^\circ C$</th> <th>$mol(1)/dm^3$ sln</th> <th>$g(2)/dm^3$ sln (compiler)</th> </tr> </thead> <tbody> <tr> <td>25.</td> <td>3.22</td> <td>58.0</td> </tr> <tr> <td>25.</td> <td>3.43</td> <td>61.8</td> </tr> </tbody> </table>		$t/^\circ C$	$g(1)/dm^3$	$mol(1)/dm^3$ sln (compiler)	16.	0.420	0.001577	17.	0.410	0.001540	19.	0.397	0.001491	22.	0.380	0.001427	$t/^\circ C$	$mol(1)/dm^3$ sln	$g(2)/dm^3$ sln (compiler)	25.	3.22	58.0	25.	3.43	61.8
$t/^\circ C$	$g(1)/dm^3$	$mol(1)/dm^3$ sln (compiler)																							
16.	0.420	0.001577																							
17.	0.410	0.001540																							
19.	0.397	0.001491																							
22.	0.380	0.001427																							
$t/^\circ C$	$mol(1)/dm^3$ sln	$g(2)/dm^3$ sln (compiler)																							
25.	3.22	58.0																							
25.	3.43	61.8																							
<p>AUXILIARY INFORMATION</p>																									
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method was used. 0.5 cm³ of ³²P-labelled organic phase was shaken with 50 cm³ of aq. phase for 20 min. After a short standing period, the aqueous phase was separated and centrifuged. A sample was diluted 10-fold, and 10 cm³ were placed in a GM6 liquid counter for counting and comparison with a suitable standard. The method was reported together with the systems phosphoric acid tributyl ester-water-diluent-nitric acid.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified; purified by boiling with dilute NaOH solution to distill volatile impurities; the remaining portion washed repeatedly with water. ³²P-labelled (1) supplied by Radiochemical Center, Amersham; purified as described above.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <p>REFERENCES:</p>																								

<p>COMPONENTS:</p> <p>(1) Phosphoric acid tributyl ester (tributyl phosphate); $C_{12}H_{27}O_4P$; [126-73-8]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Burger, L.L.; Forsman, R.C. Report HW-20936, 1951.</p> <p>Data taken from: Bruce, F.R.; Fletcher, J.M.; Hyman, H.H. Progress in Nuclear Energy, ser.3, Process Chemistry vol.2, Pergamon Press, 1958, 546-56.</p>
<p>VARIABLES:</p> <p>$T/K = 298$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of phosphoric acid tributyl ester in water at 25°C was reported to be 0.39 g(1)/dm³ sln. The corresponding value on a mole/volume basis is 0.00146 mol(1)/dm³.</p> <p>The solubility of water in phosphoric acid tributyl ester at 25°C was reported to be 64 g(2)/dm³ sln. The corresponding value on a mole/volume basis is 3.55 mol(2)/dm³ sln.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The method was not specified.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Not specified. (2) Not specified.</p> <hr/> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <hr/> <p>REFERENCES:</p>

COMPONENTS: (1) Phosphoric acid tributyl ester (tributyl phosphate); $C_{12}H_{27}O_4P$; [126-73-8] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Higgins, C.E.; Baldwin, W.H.; Soldano, B.A. <i>J. Phys. Chem.</i> <u>1959</u> , 63, 113-8.
VARIABLES: $T/K = 277 - 323$	PREPARED BY: A. Skrzecz

EXPERIMENTAL VALUES:

Solubility of phosphoric acid tributyl ester in water

$t/^\circ C$	mg(1)/dm ³ sln	mol(1)/dm ³ sln
3.4	1075	0.0040
4.0	1012	0.0038 ^b
5.0	957	0.0036 ^b
13.0	640	0.0024 ^b
25.0	422 ^a	0.0016
50.0	285	0.0011

^a The average of 30 measurements.^b Calculated by the compiler.**AUXILIARY INFORMATION**

METHOD/APPARATUS/PROCEDURE: The analytical method was used. The mixture of water and ester labeled with ³² P were tumbled in a constant temperature bath for 1/2 to 1 h. and stored overnight to ensure complete separation. The aqueous phase was analysed by counting a dilution of an aliquot containing sufficient activity in the solution counter as described by Higgins and Baldwin (ref 1). The specific activity of the labeled ester was measured at the same time for comparison. The averages of at least two separated determinations were presented.	SOURCE AND PURITY OF MATERIALS: (1) Source not specified, C.P. grade; tributyl phosphate- ³² P prepared in the laboratory as described by Higgins and Baldwin (ref 2); specific activity 40-50(micro)c/mmole. (2) Not specified.
	ESTIMATED ERROR: Temp. $\pm 0.2^\circ C$. Soly. $\pm 1.8\%$ (the largest average deviation of a single determination at 25°C).
	REFERENCES: 1. Higgins, C.E.; Baldwin, W.H. <i>Anal. Chem.</i> <u>1955</u> , 27, 1780. 2. Higgins, C.E.; Baldwin, W.H. <i>J. Org. Chem.</i> <u>1956</u> , 21, 1156.

<p>COMPONENTS:</p> <p>(1) Phosphoric acid tributyl ester (tributyl phosphate); $C_{12}H_{27}O_4P$; [126-73-8]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Hasegawa T. <i>Kogyo Kagaku Zasshi</i> <u>1961</u>, 64(7), 1239-41.</p>
<p>VARIABLES:</p> <p>$T/K = 298$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of phosphoric acid tributyl ester in water 25°C was reported to be 6.8 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 0.0049.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/Apparatus/Procedure:</p> <p>The titration method was used. The cloudiness was observed visually. The data and method were reported together with the ternary systems phosphoric acid tributyl ester-water-2-furancarbal (tributyl phosphate-water-furfural). No further details were reported in the paper.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified; boiled with 5 volumes of 0.4% NaOH(aq) washed several times with water, distilled at reduced pressure; b.p. 132°C at 9 mm Hg, d_4^{25} 0.9538.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <p>REFERENCES:</p>

<p>COMPONENTS:</p> <p>(1) Phosphoric acid tributyl ester (tributyl phosphate); $C_{12}H_{27}O_4P$; [126-73-8]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Mikhailov, V.A.; Kharchenko, S.K.; Nazin, A.G.</p> <p>Izv. Sib. Otd. Akad. Nauk SSSR 1962, (7), 50-6.</p>																		
<p>VARIABLES:</p> <p>$T/K = 279 - 299$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>																		
<p>EXPERIMENTAL VALUES:</p> <p>Solubility of water in phosphoric acid tributyl ester</p> <table border="1" data-bbox="171 514 1029 796"> <thead> <tr> <th>$t/^\circ C$</th> <th>$g(2)/100g\ sln$</th> <th>x_2 (compiler)</th> </tr> </thead> <tbody> <tr> <td>5.5^b</td> <td>6.37</td> <td>0.501</td> </tr> <tr> <td>17.^b</td> <td>6.25</td> <td>0.496</td> </tr> <tr> <td>25.^a</td> <td>6.244(0.014)^d</td> <td>0.4961(0.0005) (authors)</td> </tr> <tr> <td>25.^c</td> <td>6.38</td> <td>0.502</td> </tr> <tr> <td>26.^b</td> <td>6.22</td> <td>0.495</td> </tr> </tbody> </table> <p>^a Analytical method. ^b Alekseev's synthetic method. ^c Isopiestic method. ^d The mean of 8 experiments.</p>		$t/^\circ C$	$g(2)/100g\ sln$	x_2 (compiler)	5.5 ^b	6.37	0.501	17. ^b	6.25	0.496	25. ^a	6.244(0.014) ^d	0.4961(0.0005) (authors)	25. ^c	6.38	0.502	26. ^b	6.22	0.495
$t/^\circ C$	$g(2)/100g\ sln$	x_2 (compiler)																	
5.5 ^b	6.37	0.501																	
17. ^b	6.25	0.496																	
25. ^a	6.244(0.014) ^d	0.4961(0.0005) (authors)																	
25. ^c	6.38	0.502																	
26. ^b	6.22	0.495																	
<p>AUXILIARY INFORMATION</p>																			
<p>METHOD/AppARATUS/PROCEDURE:</p> <p>The analytical, Alekseev's synthetic, and the isoplezitic methods were used. In the analytical method, the phases were separated, after thermostating 2-3 h., using a centrifuge for 20 sec. (for most accurate results), and the water was titrated with Karl Fischer reagent. For the Alekseev's method, sealed ampoules containing a known sample composition along with a magnetic stirrer were placed into a water bath and the temperature of cloudiness was observed. In the isopiestic procedure, the assumption that the ester phase was in equilibrium with pure water was made. No further details were provided in the paper.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified; distilled at vacuum over solid Na_2CO_3; d_4^{25} 0.9728, n_D^{25} 1.4244.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm 0.05^\circ C^a$ and $\pm (1-2)^\circ C^b$.</p> <p>REFERENCES:</p>																		

<p>COMPONENTS:</p> <p>(1) Phosphoric acid tributyl ester (tributyl phosphate); $C_{12}H_{27}O_4P$; [126-73-8]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Hardy, C.J.; Fairhurst, D.; McKay, H.A.C.; Willson, A.M. <i>Trans. Faraday Soc.</i> 1964, 60, 1626-36.</p>																														
<p>VARIABLES:</p> <p>$T/K = 273 - 333$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>																														
<p>EXPERIMENTAL VALUES:</p> <p style="text-align: center;">Solubility of water in phosphoric acid tributyl ester</p> <table border="1" style="width: 100%; border-collapse: collapse;"> <thead> <tr> <th style="text-align: center;">$t/^\circ C$</th> <th style="text-align: center;">$mol(2)/dm^3$ sln</th> <th style="text-align: center;">density g/cm^3</th> <th style="text-align: center;">$g(2)/100g$ sln (compiler)</th> <th style="text-align: center;">x_2 (compiler)</th> </tr> </thead> <tbody> <tr> <td style="text-align: center;">0.(0.2)</td> <td style="text-align: center;">3.92</td> <td style="text-align: center;">0.9982</td> <td style="text-align: center;">7.09</td> <td style="text-align: center;">0.5296</td> </tr> <tr> <td style="text-align: center;">25.(0.2)</td> <td style="text-align: center;">3.59(0.03)^a</td> <td style="text-align: center;">0.9760</td> <td style="text-align: center;">6.64</td> <td style="text-align: center;">0.5120</td> </tr> <tr> <td style="text-align: center;">40.(0.1)</td> <td style="text-align: center;">3.34</td> <td style="text-align: center;">0.9640</td> <td style="text-align: center;">6.26</td> <td style="text-align: center;">0.4960</td> </tr> <tr> <td style="text-align: center;">50.(0.1)</td> <td style="text-align: center;">3.23</td> <td style="text-align: center;">0.9573^b</td> <td style="text-align: center;">6.08</td> <td style="text-align: center;">0.4889</td> </tr> <tr> <td style="text-align: center;">60.(0.1)</td> <td style="text-align: center;">3.07</td> <td style="text-align: center;">0.9508^b</td> <td style="text-align: center;">5.82</td> <td style="text-align: center;">0.4773</td> </tr> </tbody> </table> <p>^a The average of 17 determinations (each in duplicate): 5 by shaking, 5 by "very fast" stirring, 3 by "fast" stirring and 4 by saturation with water vapour.</p> <p>^b Calculated from authors' equation: $d_t = 0.9982 (1.0 - 9.6 \times 10^{-4} (t/^\circ C) + 2.8 \times 10^{-6} (t/^\circ C)^2)$</p>		$t/^\circ C$	$mol(2)/dm^3$ sln	density g/cm^3	$g(2)/100g$ sln (compiler)	x_2 (compiler)	0.(0.2)	3.92	0.9982	7.09	0.5296	25.(0.2)	3.59(0.03) ^a	0.9760	6.64	0.5120	40.(0.1)	3.34	0.9640	6.26	0.4960	50.(0.1)	3.23	0.9573 ^b	6.08	0.4889	60.(0.1)	3.07	0.9508 ^b	5.82	0.4773
$t/^\circ C$	$mol(2)/dm^3$ sln	density g/cm^3	$g(2)/100g$ sln (compiler)	x_2 (compiler)																											
0.(0.2)	3.92	0.9982	7.09	0.5296																											
25.(0.2)	3.59(0.03) ^a	0.9760	6.64	0.5120																											
40.(0.1)	3.34	0.9640	6.26	0.4960																											
50.(0.1)	3.23	0.9573 ^b	6.08	0.4889																											
60.(0.1)	3.07	0.9508 ^b	5.82	0.4773																											
<p>AUXILIARY INFORMATION</p>																															
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method was used. The solutions were prepared by shaking together known weights of components for 15 min or by saturating ester with steam. The ester was equilibrated with water at 25°C by:</p> <p>(i) shaking in a mechanical shaker for 15 minutes and centrifuged,</p> <p>(ii) stirring 10 cm³ of each phase in a 40 cm³ centrifuge tube for 15-60 min at one of 3 different speeds (slow, fast, very fast)</p> <p>(iii) bubbling nitrogen saturated with steam through the tube containing the pure ester; water concentration becoming constant after 24 h.</p> <p>The phases at 0°C were separated by centrifugation and at 40-60°C they were settled for 1 h. in a stoppered tube at the required temperature. The water concentration was determined by the standard Karl Fischer reagent using the dead-stop end-point method.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, commercially available; washed and heated with 0.1 N alkali, volatile matter distilled in a stream of nitrogen, the remaining alkaline solution washed six times with water, dried under vacuum at 40-50°C with a nitrogen bleed, stored in a dark glass stoppered bottle.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>See above.</p> <p>REFERENCES:</p>																														

<p>COMPONENTS:</p> <p>(1) Phosphoric acid tributyl ester (tributyl phosphate); $C_{12}H_{27}O_4P$; [126-73-8]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Nikolaev, A.V.; Dyadin, Yu.A.; Yakovlev, I.I.; Durasov, V.B.; Mironova, Z.N.</p> <p><i>Izv. Sib. Otd. Akad. Nauk SSSR,</i> <i>Ser. Khim. Nauk</i> <u>1965</u>, (3), 27-31.</p>																																																																					
<p>VARIABLES:</p> <p>$T/K = 273 - 460$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>																																																																					
<p>EXPERIMENTAL VALUES:</p> <p>Mutual solubility of phosphoric acid tributyl ester and water</p> <table border="1" data-bbox="134 566 1136 1124"> <thead> <tr> <th rowspan="2">$t/^\circ C$</th> <th colspan="2">g(1)/100g sln</th> <th colspan="2">x_1 (compiler)</th> </tr> <tr> <th>(2)-rich phase</th> <th>(1)-rich phase</th> <th>(2)-rich phase</th> <th>(1)-rich phase</th> </tr> </thead> <tbody> <tr><td>0.^a</td><td>-</td><td>92.48</td><td>-</td><td>0.4541</td></tr> <tr><td>4.6</td><td>-</td><td>92.70</td><td>-</td><td>0.4621</td></tr> <tr><td>5.4</td><td>0.0994</td><td>-</td><td>6.73×10^{-5}</td><td>-</td></tr> <tr><td>9.1</td><td>-</td><td>92.89</td><td>-</td><td>0.4692</td></tr> <tr><td>13.9</td><td>0.0620</td><td>-</td><td>4.20×10^{-5}</td><td>-</td></tr> <tr><td>14.6</td><td>-</td><td>93.09</td><td>-</td><td>0.4756</td></tr> <tr><td>16.3</td><td>-</td><td>93.18</td><td>-</td><td>0.4803</td></tr> <tr><td>19.3</td><td>0.0474</td><td>-</td><td>3.21×10^{-5}</td><td>-</td></tr> <tr><td>21.6</td><td>-</td><td>93.33</td><td>-</td><td>0.4863</td></tr> <tr><td>24.2</td><td>-</td><td>93.38</td><td>-</td><td>0.4883</td></tr> <tr><td>24.7</td><td>0.0403</td><td>-</td><td>2.73×10^{-5}</td><td>-</td></tr> <tr><td>25.0^a</td><td>-</td><td>93.44</td><td>-</td><td>0.4907</td></tr> </tbody> </table> <p style="text-align: right;">(continued next page)</p>		$t/^\circ C$	g(1)/100g sln		x_1 (compiler)		(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase	0. ^a	-	92.48	-	0.4541	4.6	-	92.70	-	0.4621	5.4	0.0994	-	6.73×10^{-5}	-	9.1	-	92.89	-	0.4692	13.9	0.0620	-	4.20×10^{-5}	-	14.6	-	93.09	-	0.4756	16.3	-	93.18	-	0.4803	19.3	0.0474	-	3.21×10^{-5}	-	21.6	-	93.33	-	0.4863	24.2	-	93.38	-	0.4883	24.7	0.0403	-	2.73×10^{-5}	-	25.0 ^a	-	93.44	-	0.4907
$t/^\circ C$	g(1)/100g sln		x_1 (compiler)																																																																			
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase																																																																		
0. ^a	-	92.48	-	0.4541																																																																		
4.6	-	92.70	-	0.4621																																																																		
5.4	0.0994	-	6.73×10^{-5}	-																																																																		
9.1	-	92.89	-	0.4692																																																																		
13.9	0.0620	-	4.20×10^{-5}	-																																																																		
14.6	-	93.09	-	0.4756																																																																		
16.3	-	93.18	-	0.4803																																																																		
19.3	0.0474	-	3.21×10^{-5}	-																																																																		
21.6	-	93.33	-	0.4863																																																																		
24.2	-	93.38	-	0.4883																																																																		
24.7	0.0403	-	2.73×10^{-5}	-																																																																		
25.0 ^a	-	93.44	-	0.4907																																																																		
<p>AUXILIARY INFORMATION</p>																																																																						
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The synthetic method proposed by Aleksejev was used. Light filters and a high intensity light source were used to measure the cloudiness point with good accuracy. The Karl Fischer analytical method was used as a check. No further details were reported in the paper.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, pure grade; washed several times with NaOH(aq), diluted HCL, H_2O, three times distilled under vacuum; b.p. $132-3^\circ C$ at 3 mm Hg, d_4^{25} 0.9723, n_D^{25} 1.4224.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm(0.2-0.6)^\circ C$ below $30^\circ C$, $\pm(1-3)^\circ C$ at high temperature.</p> <p>REFERENCES:</p>																																																																					

<p>COMPONENTS:</p> <p>(1) Phosphoric acid tributyl ester (tributyl phosphate); $C_{12}H_{27}O_4P$; [126-73-8]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Nikolaev, A.V.; Dyadin, Yu.A.; Yakovlev, I.I.; Durasov, V.B.; Mironova, Z.N.</p> <p><i>Izv. Sib. Otd. Akad. Nauk SSSR,</i> <i>Ser. Khim. Nauk</i> <u>1965</u>, (3), 27-31.</p>
---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------

EXPERIMENTAL VALUES: (continued)

Mutual solubility of phosphoric acid tributyl ester and water

t/°C	g(1)/100g sln		x_1 (compiler)	
	(2)-rich phase	(1)-rich phase	(2)-rich phase	(1)-rich phase
25.1	-	93.41	-	0.4895
25.3	-	93.46	-	0.4915
29.5	-	93.52	-	0.4940
33.1	0.0329	-	2.23×10^{-5}	-
34.2	-	93.58	-	0.4965
39.2	0.0292	-	1.98×10^{-5}	-
60.2	-	93.58	-	0.4965
65.0	-	93.52	-	0.4940
69.5	-	93.41	-	0.4895
71.0	-	93.38	-	0.4883
78.0	-	93.18	-	0.4803
80.8	0.0292	-	1.98×10^{-5}	-
82.0	-	93.09	-	0.4756
86.0	0.0329	-	2.23×10^{-5}	-
94.5	0.0403	92.70	2.73×10^{-5}	0.4621
106.0	-	92.22	-	0.4450
117.0	-	91.72	-	0.4284
131.0	-	92.77	-	0.4646
134.0	-	89.02	-	0.3542
187.0	-	88.77	-	0.3484

^a Karl Fischer analytical method.

<p>COMPONENTS:</p> <p>(1) Phosphoric acid tris(2-methylpropyl) ester (<i>tri(2-methylpropyl) phosphate</i>); $C_{12}H_{27}O_4P$; [126-71-6]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Apelblat, A. <i>J. Chem. Soc. B</i> <u>1969</u>, 175-7.</p>
<p>VARIABLES:</p> <p>$T/K = 298$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of water in phosphoric acid tris(2-methylpropyl) ester at 25°C was reported to be 3.50 mol(2)/L sln. The corresponding value on a mole/volume basis is 63.1 g(2)/L sln (compiler).</p> <p>The density of the ester-phase was reported was to be d^{25} 0.9655.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method was used. The mixture was equilibrated by vigorous stirring for 30 min. and settling for 12 h. The organic phase was analyzed with the Karl Fischer reagent. Preliminary experiments showed that stirring and settling times were sufficient.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Albright and Wilson Co; purified by method described in (ref 1); d_4^{25} 0.9614, n_D^{25} 1.4168.</p> <p>(2) Not specified.</p> <hr/> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm 0.1^\circ C$.</p> <hr/> <p>REFERENCES:</p> <p>1. Alcock, K.; Grimley, S.S.; Healy, T.V.; Kennedy, J.; McKay, H.A.C. <i>Trans. Faraday Soc.</i> <u>1956</u>, 52, 39.</p>

COMPONENTS:	EVALUATOR:		
(1) Benzoic acid, 2-hydroxy-, phenyl ester (phenyl salicylate); $C_{13}H_{10}O_3$; [118-55-8] (2) Water; H_2O ; [7732-18-5]	G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia March, 1990		
CRITICAL EVALUATION:			
Quantitative solubility data for the 2-hydroxybenzoic acid phenyl ester (1) - water (2) system have been reported in the publications listed in Table 1.			
<u>TABLE 1: Quantitative Solubility Studies of the 2-Hydroxybenzoic acid phenyl ester (1) - Water (2) System</u>			
Reference	T/K	Solubility g(1)/100 g sln	Method
Seidell (ref 1)	298	0.015	gravimetric
Dehn (ref 2)	RT ^a	0.04	gravimetric
^a Room temperature (293-298 K).			
As can be seen from Table 1, the data of Seidell (ref 1) and Dehn (ref 2) for the solubility of phenyl salicylate (1) in water (2) are not in particularly good agreement. In the absence of further information it is not possible to prefer either value although it may be noted that Seidell's data in other systems are generally close to Recommended values.			
REFERENCES			
1. Seidell, A. <i>Hygienic Lab. Bull.</i> <u>1910</u> , No. 67, 98pp. (U.S. Govt. Printing Office, Washington, DC). 2. Dehn, W. M. <i>J. Am. Chem. Soc.</i> <u>1917</u> , 39, 1399-404.			

COMPONENTS: (1) Benzoic acid, 2-hydroxy-, phenyl ester (phenyl salicylate); $C_{13}H_{10}O_3$; [118-55-8] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Seidell, A. <i>Hygienic Lab. Bull.</i> 1910, No. 67, 98 pp (US Govt. Printing Office, Washington, DC).
VARIABLES: $T/K = 298$	PREPARED BY: G.T. Hefter
EXPERIMENTAL VALUES: The solubility of 2-hydroxybenzoic acid phenyl ester in water at 25°C was reported to be 0.015 g(1)/100g sln. The corresponding mole fraction, x_1 , value calculated by the compiler is 1.3×10^{-5} .	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The analytical method was used. Appropriate quantities of (1) and (2) were shaken together for two days. The amount of dissolved (1) was determined by evaporation of a known quantity of the saturated solution. The residue was dried to a constant weight in a vacuum desiccator.	SOURCE AND PURITY OF MATERIALS: (1) Source not stated; purity determined to be practically 100% by saponification. (2) Distilled (no details given).
	ESTIMATED ERROR: Temp. not stated. Soly. not stated.
	REFERENCES:

<p>COMPONENTS:</p> <p>(1) Benzoic acid, 2-hydroxy-, phenyl ester (phenyl salicylate); $C_{13}H_{10}O_3$; [118-55-8]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Dehn, W.M. <i>J. Am. Chem. Soc.</i> <u>1917</u>, <i>39</i>, 1399-1404.</p>
<p>VARIABLES:</p> <p>T/K = room temperature</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of 2-hydroxybenzoic acid phenyl ester in water at room temperature^a was reported to be 0.04 g(1)/100g(1). The corresponding mass percentage and mole fraction, x_1, values calculated by the compiler are 0.040 g(1)/100g sln and 3.4×10^{-5}.</p> <p>^a Room temperature was reported to be 20-25°C.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>Excess of the substance and 5 mL of water were enclosed in small vials which, when stoppered, were shaken or were left until equilibrium was established. The solutions were filtered into weighed crucibles and reweighed. After drying in a vacuum desiccator or on a water bath, the crucibles were again weighed and the loss of solvent calculated. The measurements were made at room temperatures from 20 to 25°C.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, commercial grade; used as received. (2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <p>REFERENCES:</p>

COMPONENTS: (1) Phosphoric acid methyl diphenyl ester (methyl diphenyl phosphate); $C_{13}H_{13}O_4P$; [115-89-9] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Apelblat, A. <i>J. Chem. Soc. B</i> <u>1969</u> , 175-7.
VARIABLES: $T/K = 298$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: The solubility of water in phosphoric acid methyl diphenyl ester at 25°C was reported to be 1.49 mol(2)/L sln. The corresponding value on a mass/volume basis is 26.8 g(2)/L sln (compiler). The density of the ester-phase was reported was to be d^{25} 1.2195.	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The analytical method was used. The mixture was equilibrated by vigorous stirring for 30 min. and settling for 12 h. The organic phase was analyzed with the Karl Fischer reagent. Preliminary experiments showed that stirring and settling times were sufficient.	SOURCE AND PURITY OF MATERIALS: (1) Aldrich Chemical Co.; purified by method described by Alcock, Grimley, Healy, Kennedy and McKay (ref 1); d_4^{25} 1.2258, n_D^{25} 1.5356. (2) Not specified. ESTIMATED ERROR: Temp. $\pm 0.1^\circ C$. REFERENCES: 1. Alcock, K.; Grimley, S.S.; Healy, T.V.; Kennedy, J.; McKay, H.A.C. <i>Trans. Faraday Soc.</i> <u>1956</u> , 52, 39.

<p>COMPONENTS:</p> <p>(1) Nonanedioic acid diethyl ester (diethyl nonanedioate); $C_{13}H_{24}O_4$; [624-17-9]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Sobotka, H.; Kahn, J. <i>J. Am. Chem. Soc.</i> <u>1931</u>, 53, 2935-8.</p>
<p>VARIABLES:</p> <p>$T/K = 293$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of nonanedioic acid diethyl ester in water at 20°C was reported to be 0.025 g(1)/100mL(2). The corresponding mass per cent and mole fraction, x_1, values calculated by the compiler are 0.025 g(1)/100g sln and 1.8×10^{-5}.</p> <p>Density of water $d_4^{20} = 0.9982$ (ref 1) was used in the calculation.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The titration method was used. The ester was added dropwise from a micro-burette with a capillary tip to 100, 250, or 500 mL of water in a narrow-mouthed stock bottle with a well-ground glass stopper. The bottle was shaken after each addition of ester. 1-5 mg of Sudan IV dye was put into the water to improve the end-point of the titration. At saturation, one additional drop of ester was sufficient to convert the floating rough indicator particles into dark transparent droplets.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified (Eastman Kodak Laboratories or synthesized); twice-distilled under reduced pressure; d_4^{20} 0.9729, n_D^{20} 1.4358.</p> <p>(2) Distilled.</p> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <p>REFERENCES:</p> <p>1. <i>Selected Values of Properties of Hydrocarbons and Related Compounds</i>, API Research Project 44, Thermodynamics Research Center, Texas A and M University, Texas, <u>1973</u>.</p>

<p>COMPONENTS:</p> <p>(1) 1,2-Benzenedicarboxylic acid di-2-propenyl ester (diallyl phthalate); C₁₄H₁₄O₄; [131-17-9]</p> <p>(2) Water; H₂O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Leyder, F.; Boulanger, P.</p> <p><i>Bull. Environ. Contam. Toxicol.</i> <u>1983</u>, 30(2), 152-7.</p>
<p>VARIABLES:</p> <p>T/K = 293</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of 1,2-benzenedicarboxylic acid di-2-propenyl ester in water at 20°C was reported to be 7.39 x 10⁻⁴ mol(1)/L and 0.182 g(1)/L.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method as described in the <i>OECD Guidelines for Testing of Chemicals</i> (ref 1) was used. Gas chromatography analyses were conducted on a Girdel 3000 FFLE apparatus with flame-ionization detection. A 4 mm x 180 m stainless steel column packed with 3% OV-1 on 80-100 mesh Chromosorb W-AW-DMCS and helium at a 20mL/min flow rate were used. Samples were fortified with an internal standard, extracted, and concentrated to 1 mL of hexane solution. "Solvent flush" and "hot needle" injection techniques were used. The two methods gave similar results. Peak heights were measured throughout. UV measurements were made by using 1-cm cells in a Perkin-Elmer model 552 uv-Visible spectrophotometer.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) CEPEA; purity not specified; used as received.</p> <p>(2) Deionized and distilled from KMnO₄.</p> <p>ESTIMATED ERROR:</p> <p>Temp. ±1°C.</p> <p>REFERENCES:</p> <p>1. <i>OECD Guidelines for Testing of Chemicals</i>, Paris, OECD, <u>1981</u>. (methods 101, 105).</p>

<p>COMPONENTS:</p> <p>(1) 1,2-Benzenedicarboxylic acid bis(1-methylethyl) ester (diisopropyl phthalate); $C_{14}H_{18}O_4$; [605-45-8]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Leyder, F.; Boulanger, P.</p> <p><i>Bull. Environ. Contam. Toxicol.</i> <u>1983</u>, 30(2), 152-7.</p>
<p>VARIABLES:</p> <p>$T/K = 293$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of 1,2-benzenedicarboxylic acid bis(1-methylethyl) ester in water at 20°C was reported to be 1.33×10^{-3} mol(1)/L and 0.332 g(1)/L.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method as described in the <i>OECD Guidelines for Testing of Chemicals</i> (ref 1) was used. Gas chromatography analyses were conducted on a Girdel 3000 FFLE apparatus with flame-ionization detection. A 4 mm x 180 m stainless steel column packed with 3% OV-1 on 80-100 mesh Chromosorb W-AW-DMCS and helium at a 20mL/min flow rate were used. Samples were fortified with an internal standard, extracted, and concentrated to 1 mL of hexane solution. "Solvent flush" and "hot needle" injection techniques were used. The two methods gave similar results. Peak heights were measured throughout. UV measurements were made by using 1-cm cells in a Perkin-Elmer model 552 uv-Visible spectrophotometer.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Chrompack, stationary phase for gas chromatography; used as received.</p> <p>(2) Deionized and distilled from $KMnO_4$.</p> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm 1^\circ C$.</p> <p>REFERENCES:</p> <p>1. <i>OECD Guidelines for Testing of Chemicals</i>, Paris, OECD, <u>1981</u>. (methods 101, 105).</p>

<p>COMPONENTS:</p> <p>(1) 1,2-Benzenedicarboxylic acid dipropyl ester (dipropyl phthalate); C₁₄H₁₈O₄; [131-16-8]</p> <p>(2) Water; H₂O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Leyder, F.; Boulanger, P. <i>Bull. Environ. Contam. Toxicol.</i> <u>1983</u>, 30(2), 152-7.</p>
<p>VARIABLES:</p> <p>T/K = 293</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of 1,2-benzenedicarboxylic acid dipropyl ester in water at 20°C was reported to be 4.32 x 10⁻⁴ mol(1)/L and 0.108 g(1)/L.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method as described in the <i>OECD Guidelines for Testing of Chemicals</i> (ref 1) was used. Gas chromatography analyses were conducted on a Girdel 3000 FFLE apparatus with flame-ionization detection. A 4 mm x 180 m stainless steel column packed with 3% OV-1 on 80-100 mesh Chromosorb W-AW-DMCS and helium at a 20mL/min flow rate were used. Samples were fortified with an internal standard, extracted, and concentrated to 1 mL of hexane solution. "Solvent flush" and "hot needle" injection techniques were used. The two methods gave similar results. Peak heights were measured throughout. UV measurements were made by using 1-cm cells in a Perkin-Elmer model 552 uv-Visible spectrophotometer.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Chrompack, stationary phase for gas chromatography; used as received.</p> <p>(2) Deionized and distilled from KMnO₄.</p> <p>ESTIMATED ERROR:</p> <p>Temp. ±1°C.</p> <p>REFERENCES:</p> <p>1. <i>OECD Guidelines for Testing of Chemicals</i>, Paris, OECD, <u>1981</u>. (methods 101, 105).</p>

<p>COMPONENTS:</p> <p>(1) Decanedioic acid diethyl ester (<i>diethyl decanedioate</i>); $C_{14}H_{26}O_4$; [110-40-7]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Sobotka, H.; Kahn, J. <i>J. Am. Chem. Soc.</i> <u>1931</u>, 53, 2935-8.</p>
<p>VARIABLES:</p> <p>$T/K = 293$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of decanedioic acid diethyl ester in water at 20°C was reported to be 0.008 g(1)/100mL(2). The corresponding mass per cent and mole fraction, x_1, values calculated by the compiler are 0.008 g(1)/100g sln and 6×10^{-6}.</p> <p>Density of water $d_4^{20} = 0.9982$ (ref 1) was used in the calculation.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The titration method was used. The ester was added dropwise from a micro-burette with a capillary tip to 100, 250, or 500 mL of water in a narrow-mouthed stock bottle with a well-ground glass stopper. The bottle was shaken after each addition of ester. 1-5 mg of Sudan IV dye was put into the water to improve the end-point of the titration. At saturation, one additional drop of ester was sufficient to convert the floating rough indicator particles into dark transparent droplets.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified (Eastman Kodak Laboratories or synthesized); twice-distilled under reduced pressure; d_4^{20} 0.9646, n_D^{20} 1.4368.</p> <p>(2) Distilled.</p> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <p>REFERENCES:</p> <p>1. <i>Selected Values of Properties of Hydrocarbons and Related Compounds</i>, API Research Project 44, Thermodynamics Research Center, Texas A and M University, Texas, <u>1973</u>.</p>

COMPONENTS: (1) Phosphoric acid dibutyl methylphenyl ester (dibutyl tolyl phosphate); $C_{15}H_{25}O_4P$; [26446-69-5] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Apelblat, A. <i>J. Chem. Soc. B</i> <u>1969</u> , 175-7.
VARIABLES: $T/K = 298$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: The solubility of water in phosphoric acid dibutyl methylphenyl ester at 25°C was reported to be 1.50 mol(2)/L sln. The corresponding value on a mass/volume basis is 27.0 g(2)/L sln (compiler).	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The analytical method was used. The mixture was equilibrated by vigorous stirring for 30 min. and settling for 12 h. The organic phase was analyzed with the Karl Fischer reagent. Preliminary experiments showed that stirring and settling times were sufficient.	SOURCE AND PURITY OF MATERIALS: (1) Synthesized at the Plastics Research Laboratory of the Weizmann Institute of Science; used as received; b.p. 132-5°C at 0.3 mm Hg, n_D^{25} 1.4722. (2) Not specified.
	ESTIMATED ERROR: Temp. $\pm 0.1^\circ C$.
	REFERENCES:

<p>COMPONENTS:</p> <p>(1) 1,2-Benzenedicarboxylic acid bis(2-methylpropyl) ester (diisobutyl phthalate); $C_{16}H_{22}O_4$; [84-69-5]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Leyder, F.; Boulanger, P. <i>Bull. Environ. Contam. Toxicol.</i> 1983, 30(2), 152-7.</p>
<p>VARIABLES:</p> <p>$T/K = 293$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of 1,2-benzenedicarboxylic acid bis(2-methylpropyl) ester in water at 20°C was reported to be 7.30×10^{-5} mol(1)/L and 0.0203 g(1)/L.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method as described in the <i>OECD Guidelines for Testing of Chemicals</i> (ref 1) was used. Gas chromatography analyses were conducted on a Girdel 3000 FFLE apparatus with flame-ionization detection. A 4 mm x 180 m stainless steel column packed with 3% OV-1 on 80-100 mesh Chromosorb W-AW-DMCS and helium at a 20mL/min flow rate were used. Samples were fortified with an internal standard, extracted, and concentrated to 1 mL of hexane solution. "Solvent flush" and "hot needle" injection techniques were used. The two methods gave similar results. Peak heights were measured throughout. UV measurements were made by using 1-cm cells in a Perkin-Elmer model 552 uv-Visible spectrophotometer.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) BASF "Palatinol IC"; used as received.</p> <p>(2) Deionized and distilled from $KMnO_4$.</p> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm 1^\circ C$.</p> <p>REFERENCES:</p> <p>1. <i>OECD Guidelines for Testing of Chemicals</i>, Paris, OECD, 1981. (methods 101, 105).</p>

<p>COMPONENTS:</p> <p>(1) 1,2-Benzenedicarboxylic acid dibutyl ester (dibutyl phthalate); $C_{16}H_{22}O_4$; [84-74-2]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988</p>
------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION:

Quantitative solubility data for the 1,2-benzenedicarboxylic acid dibutyl ester (1) - water (2) system have been reported in the publications listed in Table 1.

TABLE 1: Quantitative Solubility Studies of the 1,2-Benzenedicarboxylic acid dibutyl ester (1) - Water (2) System

Reference	T/K	Solubility	Method
Krupatkin and Glagoleva (ref 1)	298	mutual	titration
Schwarz (ref 2)	297 ^a	(1) in (2)	chromatographic
Leyder and Boulanger (ref 3)	293	(1) in (2)	GLC

^a 23.5 °C.

The original data in these publications are compiled in the Data Sheets immediately following this Critical Evaluation. For convenience, further discussion of this system will be divided into two parts.

1. SOLUBILITY OF 1,2-BENZENEDICARBOXYLIC ACID DIBUTYL ESTER (1)
IN WATER (2)

All the available data for the solubility of 1,2-benzenedicarboxylic acid dibutyl ester (1) in water (2) are summarized in Table 2. The value of Krupatkin and Glagoleva (ref 1) is much higher than the other studies and is rejected. Pending further independent studies the remaining data (ref 2,3) must be regarded as very Tentative.

(continued next page)

<p>COMPONENTS:</p> <p>(1) 1,2-Benzenedicarboxylic acid dibutyl ester (dibutyl phthalate); C₁₆H₂₂O₄; [84-74-2]</p> <p>(2) Water; H₂O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988</p>
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	-----------------------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION: (continued)

TABLE 2: Reported Solubilities
of 1,2-Benzenedicarboxylic acid dibutyl ester (1) in Water (2)

T/K	Reported Solubilities	
	10 ³ g(1)/100g sln	10 ⁷ x ₁
293	1.01 (ref 3)	6
297 ^a	1.83 (ref 2)	12
298	40 (ref 1) ^b	260 ^b

^a 23.5 °C.

^b Rejected value.

2. SOLUBILITY OF WATER (2) IN 1,2-BENZENEDICARBOXYLIC ACID
DIBUTYL ESTER (1)

As the only available datum for the solubility of water (2) in 1,2-benzenedicarboxylic acid dibutyl ester (1) is that of Krupatkin and Glagoleva (ref 1) no Critical Evaluation is possible. The interested user is referred to the relevant Data Sheet for the experimental solubility but it may be noted that solubilities reported by these authors often differ from reliable values.

REFERENCES

1. Krupatkin, I. L.; Glagoleva, M. F. *Zh. Prikl. Khim.* 1972, *45*, 1317-20.
2. Schwarz, F. P. *Anal. Chem.* 1980, *52*, 10-15.
3. Leyder, F.; Boulanger, P. *Bull. Environ. Contam. Toxicol.* 1983, *30*, 152-7.

<p>COMPONENTS:</p> <p>(1) 1,2-Benzenedicarboxylic acid dibutyl ester (dibutyl phthalate); $C_{16}H_{22}O_4$; [84-74-2]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Krupatkin, I.L.; Glagoleva, M.F. <i>Zh. Prikl. Khim.</i> <u>1972</u>, <i>45</i>, 1317-20.</p>
<p>VARIABLES:</p> <p>$T/K = 298$</p>	<p>PREPARED BY:</p> <p>Z. Maczynska</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of 1,2-benzenedicarboxylic acid dibutyl ester in water at 25°C was reported to be 0.04 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 3×10^{-5}.</p> <p>The solubility of water in 1,2-benzenedicarboxylic acid dibutyl ester at 25°C was reported to be 0.62 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.088.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The titration method was used. The measurements were carried out in a water thermostat. The data were reported together with the ternary system 1,2-benzenedicarboxylic acid dibutyl ester-water-2-furancarboxal (dibutyl phthalate-water-furfural). No further details were reported in the paper.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified; b.p. 340°C, d_4^{25} 1.0485, n_D^{25} 1.4882.</p> <p>(2) Twice distilled.</p> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm 0.1^\circ C$.</p> <p>REFERENCES:</p>

<p>COMPONENTS:</p> <p>(1) 1,2-Benzenedicarboxylic acid dibutyl ester (dibutyl phthalate); $C_{16}H_{22}O_4$; [84-74-2]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Schwarz, F.P. Anal. Chem. <u>1980</u>, 52, 10-5.</p>
<p>VARIABLES:</p> <p>$T/K = 297$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of 1,2-benzenedicarboxylic acid dibutyl ester in water at 23.5°C was reported to be 0.00183 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 1.18×10^{-6}.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The elution chromatographic method was used. The column was filled with Chromosorb P (80-120 mesh, 4-6 m²/g by BET) coated with the solute and prepared in a special way as described in the paper. The water passing through the solute column was collected in graduated vessels. The measurements were initiated after the appearance of the solute depleted light pink colored zone (solvent zone) on Chromosorb P. Each measurement consisted of turning off the flow, measuring the length of the zone and the total volume of the water passed through the column. Measurements were initiated after the solvent zone extended 0.3 cm down the length of the column and were continued until the solvent zone extended down at least 25% of the column length.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified, reagent grade; used as received; $d^{20} 1.047$.</p> <p>(2) Distilled.</p> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm 1.5^\circ C$. Soly. ± 0.00006 g(1)/100g sln (standard deviation).</p> <p>REFERENCES:</p>

<p>COMPONENTS:</p> <p>(1) 1,2-Benzenedicarboxylic acid dibutyl ester (dibutyl phthalate); $C_{16}H_{22}O_4$; [84-74-2]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Leyder, F.; Boulanger, P. <i>Bull. Environ. Contam. Toxicol.</i> <u>1983</u>, 30(2), 152-7.</p>
<p>VARIABLES:</p> <p>T/K = 293</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of 1,2-benzenedicarboxylic acid dibutyl ester in water at 20°C was reported to be 3.63×10^{-5} mol(1)/L and 0.0101 g(1)/L.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method as described in the <i>OECD Guidelines for Testing of Chemicals</i> (ref 1) was used. Gas chromatography analyses were conducted on a Girdel 3000 FFLE apparatus with flame-ionization detection. A 4 mm x 180 m stainless steel column packed with 3% OV-1 on 80-100 mesh Chromosorb W-AW-DMCS and helium at a 20mL/min flow rate were used. Samples were fortified with an internal standard, extracted, and concentrated to 1 mL of hexane solution. "Solvent flush" and "hot needle" injection techniques were used. The two methods gave similar results. Peak heights were measured throughout. UV measurements were made by using 1-cm cells in a Perkin-Elmer model 552 uv-Visible spectrophotometer.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Fluka; purity >98%; used as received.</p> <p>(2) Deionized and distilled from $KMnO_4$.</p> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm 1^\circ C$.</p> <p>REFERENCES:</p> <p>1. <i>OECD Guidelines for Testing of Chemicals</i>, Paris, OECD, <u>1981</u>. (methods 101, 105).</p>

<p>COMPONENTS:</p> <p>(1) Phosphoric acid butyl bis(methylphenyl) ester (butyl ditolyl phosphate); $C_{18}H_{23}O_4P$; [25657-10-7]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Apelblat, A. <i>J. Chem. Soc. B</i> <u>1969</u>, 175-7.</p>
<p>VARIABLES:</p> <p>$T/K = 298$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of water in phosphoric acid butyl bis(methylphenyl) ester at 25°C was reported to be 0.69 mol(2)/L sln. The corresponding value on a mass/volume basis is 12.4 g(2)/L sln (compiler).</p> <p>The density of the ester-phase was reported to be d^{25}_4 1.1097.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method was used. The mixture was equilibrated by vigorous stirring for 30 min. and settling for 12 h. The organic phase was analyzed with the Karl Fischer reagent. Preliminary experiments showed that stirring and settling times were sufficient.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Synthesized at the Plastics Research Laboratory of the Weizmann Institute of Science; used as received; b.p. 180°C at 0.25 mm Hg, d^{25}_4 1.1118, n^{25}_D 1.5160.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm 0.1^\circ C$.</p> <p>REFERENCES:</p>

<p>COMPONENTS:</p> <p>(1) 1,2-Benzenedicarboxylic acid dipentyl ester (dipentyl phthalate); $C_{18}H_{26}O_4$; [131-18-0]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Leyder, F.; Boulanger, P. <i>Bull. Environ. Contam. Toxicol.</i> <u>1983</u>, 30(2), 152-7.</p>
<p>VARIABLES:</p> <p>$T/K = 293$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of 1,2-benzenedicarboxylic acid dipentyl ester in water at 20°C was reported to be 0.3×10^{-6} ^a, 2.6×10^{-6} ^b mol(1)/L and 0.0001^a, 0.0008^b g(1)/L.</p> <p>^a Gas chromatography analysis. ^b UV spectrophotometry.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method as described in the <i>OECD Guidelines for Testing of Chemicals</i> (ref 1) was used. Gas chromatography analyses were conducted on a Girdel 3000 FFLE apparatus with flame-ionization detection. A 4 mm x 180 m stainless steel column packed with 3% OV-1 on 80-100 mesh Chromosorb W-AW-DMCS and helium at a 20mL/min flow rate were used. Samples were fortified with an internal standard, extracted, and concentrated to 1 mL of hexane solution. "Solvent flush" and "hot needle" injection techniques were used. The two methods gave similar results. Peak heights were measured throughout. UV measurements were made by using 1-cm cells in a Perkin-Elmer model 552 uv-Visible spectrophotometer.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Kodak, practical grade; purity >90% ; used as received.</p> <p>(2) Deionized and distilled from $KMnO_4$.</p> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm 1^\circ C$.</p> <p>REFERENCES:</p> <p>1. <i>OECD Guidelines for Testing of Chemicals</i>, Paris, OECD, <u>1981</u>. (methods 101, 105).</p>

<p>COMPONENTS:</p> <p>(1) Phosphoric acid tris(butoxyethyl) ester (tri(butoxyethyl) phosphate); $C_{18}H_{39}O_7P$; [78-51-3]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Apelblat, A. <i>J. Chem. Soc. B</i> <u>1969</u>, 175-7.</p>
<p>VARIABLES:</p> <p>$T/K = 298$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of water in phosphoric acid tris(butoxyethyl) ester at 25°C was reported to be 4.26 mol(2)/L sln. The corresponding value on a mass/volume basis is 76.7 g(2)/L sln (compiler).</p> <p>The density of the ester-phase was reported to be d^{25} 1.0204.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method was used. The mixture was equilibrated by vigorous stirring for 30 min. and settling for 12 h. The organic phase was analyzed with the Karl Fischer reagent. Preliminary experiments showed that stirring and settling times were sufficient.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Albright and Wilson Co; purified by method described by Alcock, Grimley, Healy, Kennedy and McKay (ref 1); d_4^{25} 1.0181, n_D^{25} 1.4358.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm 0.1^\circ C$.</p> <p>REFERENCES:</p> <p>1. Alcock, K.; Grimley, S.S.; Healy, T.V.; Kennedy, J.; McKay, H.A.C. <i>Trans. Faraday Soc.</i> <u>1956</u>, 52, 39.</p>

COMPONENTS: (1) 1,2-Benzenedicarboxylic acid butyl phenylmethyl ester (butyl benzyl phthalate); $C_{19}H_{20}O_4$; [85-68-7] (2) Water; H_2O ; [7732-18-5]	EVALUATOR: G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia December, 1988
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------------------------------

CRITICAL EVALUATION:

Quantitative solubility data for the 1,2-benzenedicarboxylic acid butyl phenylmethyl ester (1) - water (2) system have been reported in the publications listed in Table 1.

TABLE 1: Quantitative Solubility Studies of the 1,2-Benzenedicarboxylic acid butyl phenylmethyl ester (1) - Water (2) System

Reference	T/K	Solubility	Method
Krupatkin and Glagoleva (ref 1)	298	mutual	titration
Leyder and Boulanger (ref 2)	293	(1) in (2)	GLC

The original data in these publications are compiled in the Data Sheets immediately following this Critical Evaluation. For convenience, further discussion of this system will be divided into two parts.

1. SOLUBILITY OF 1,2-BENZENEDICARBOXYLIC ACID BUTYL PHENYLMETHYL ESTER (1) IN WATER (2)

All the available data for the solubility of 1,2-benzenedicarboxylic acid butyl phenylmethyl ester (1) in water (2) are summarized in Table 2. The data are in serious disagreement differing by three orders of magnitude! The more recent value of Leyder and Boulanger (ref 2) using a more appropriate technique is probably more reasonable but, in the absence of further studies, it is not possible to reject the datum of Krupatkin and Glagoleva (ref 1).

TABLE 2: Reported Solubilities of 1,2-Benzenedicarboxylic acid butyl phenylmethyl ester (1) in Water (2)

T/K	Reported Solubilities	
	10^4 g(1)/100g sln	$10^7 x_1$
293	2.82 ^a (ref 2)	1.63 ^a
298	2100 ^b (ref 1)	1200 ^b

^a Preferred value, see text.

^b Unlikely value, see text.

(continued next page)

<p>COMPONENTS:</p> <p>(1) 1,2-Benzenedicarboxylic acid butyl phenylmethyl ester (butyl benzyl phthalate); $C_{19}H_{20}O_4$; [85-68-7]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>EVALUATOR:</p> <p>G.T. Hefter, School of Mathematical and Physical Sciences, Murdoch University, Perth, W.A., Australia</p> <p>December, 1988</p>
<p>CRITICAL EVALUATION: (continued)</p> <p>2. SOLUBILITY OF WATER (2) IN 1,2-BENZENEDICARBOXYLIC ACID BUTYL PHENYLMETHYL ESTER (1)</p> <p>The only available datum for the solubility of water (2) in 1,2-benzenedicarboxylic acid butyl phenylmethyl ester (1) is that of Krupatkin and Glagoleva (ref 1) and thus no Critical Evaluation is possible. The interested user is referred to the relevant Data Sheet for the experimental solubility but it should be noted that solubilities reported by these authors often differ from reliable values in well characterized systems.</p> <p>REFERENCES</p> <ol style="list-style-type: none">1. Krupatkin, I. L.; Glagoleva, M. F. <i>Zh. Prikl. Khim.</i> <u>1972</u>, <i>45</i>, 1795-9.2. Leyder, F.; Boulanger, P. <i>Bull. Environ. Contam. Toxicol.</i> <u>1983</u>, <i>30</i>, 152-7.	

<p>COMPONENTS:</p> <p>(1) 1,2-Benzenedicarboxylic acid butyl phenylmethyl ester (butyl benzyl phthalate); $C_{19}H_{20}O_4$; [85-68-7]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Krupatkin, I.L.; Glagoleva, M.F. <i>Zh. Prikl. Khim.</i> <u>1972</u>, 45, 1795-9.</p>
<p>VARIABLES:</p> <p>$T/K = 298$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of 1,2-benzenedicarboxylic acid butyl phenylmethyl ester in water at 25°C was reported to be 0.21 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 1.21×10^{-4}.</p> <p>The solubility of water in 1,2-benzenedicarboxylic acid butyl phenylmethyl ester at 25°C was reported to be 0.07 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.012.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The titration method was used as given by Krupatkin and Glagoleva (ref 1). The samples were titrated up to turbidity. The data were reported together with the ternary system 1,2-benzenedicarboxylic acid butyl phenylmethyl ester-water-2-furancarboxal (butyl benzyl phthalate-water-furfural). No further details were reported in the paper.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified; b.p. 236°C, d_{25}^{25} 1.1065, n_D^{25} 1.5180.</p> <p>(2) Twice distilled.</p> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <p>REFERENCES:</p> <p>1. Krupatkin, I.L.; Glagoleva, M.F. <i>Zh. Prikl. Khim.</i> <u>1969</u>, 42, 880.</p>

<p>COMPONENTS:</p> <p>(1) 1,2-Benzenedicarboxylic acid butyl phenylmethyl ester (butyl benzyl phthalate); $C_{19}H_{20}O_4$; [85-68-7]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Leyder, F.; Boulanger, P. <i>Bull. Environ. Contam. Toxicol.</i> <u>1983</u>, 30(2), 152-7.</p>
<p>VARIABLES:</p> <p>$T/K = 293$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of 1,2-benzenedicarboxylic acid butyl phenylmethyl ester in water at 20°C was reported to be 9.02×10^{-6} mol(1)/L and 0.00282 g(1)/L.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method as described in the <i>OECD Guidelines for Testing of Chemicals</i> (ref 1) was used. Gas chromatography analyses were conducted on a Girdel 3000 FFLE apparatus with flame-ionization detection. A 4 mm x 180 m stainless steel column packed with 3% OV-1 on 80-100 mesh Chromosorb W-AW-DMCS and helium at a 20mL/min flow rate were used. Samples were fortified with an internal standard, extracted, and concentrated to 1 mL of hexane solution. "Solvent flush" and "hot needle" injection techniques were used. The two methods gave similar results. Peak heights were measured throughout. UV measurements were made by using 1-cm cells in a Perkin-Elmer model 552 uv-Visible spectrophotometer.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Bayer Unimoll BB; purity not specified; used as received.</p> <p>(2) Deionized and distilled from $KMnO_4$.</p> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm 1^\circ C$.</p> <p>REFERENCES:</p> <p>1. <i>OECD Guidelines for Testing of Chemicals</i>, Paris, OECD, <u>1981</u>. (methods 101, 105).</p>

<p>COMPONENTS:</p> <p>(1) 1,2-Benzenedicarboxylic acid butyl octyl ester (butyl octyl phthalate); $C_{20}H_{30}O_4$; [84-78-6]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Krupatkin, I.L.; Glagoleva, M.F. <i>Zh. Prikl. Khim.</i> <u>1972</u>, 45, 1795-9.</p>
<p>VARIABLES:</p> <p>$T/K = 298$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of 1,2-benzenedicarboxylic acid butyl octyl ester in water at 25°C was reported to be 0.04 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 2.2×10^{-5}.</p> <p>The solubility of water in 1,2-benzenedicarboxylic acid butyl octyl ester at 25°C was reported to be 0.02 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.0037.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The titration method was used as given by Krupatkin and Glagoleva (ref 1). The samples were titrated up to turbidity. The data were reported together with the ternary system 1,2-benzenedicarboxylic acid butyl octyl ester-water-2-furan-carbonal (butyl octyl phthalate-water-furfural). No further details were reported in the paper.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified; $d_{25}^{25} 1.0097$, $n_D^{25} 1.4868$.</p> <p>(2) Twice distilled.</p> <p>ESTIMATED ERROR:</p> <p>Not specified.</p> <p>REFERENCES:</p> <p>1. Krupatkin, I.L.; Glagoleva, M.F. <i>Zh. Prikl. Khim.</i> <u>1969</u>, 42, 880.</p>

COMPONENTS: (1) L-Ascorbic acid 6-hexadecanoate (L-ascorbyl palmitate); $C_{22}H_{38}O_7$; [137-66-6] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Swern, D. <i>J. Am. Chem. Soc.</i> <u>1949</u> , 71, 3256.
VARIABLES: $T/K = 298$	PREPARED BY: A. Skrzecz
EXPERIMENTAL VALUES: The solubility of L-ascorbic acid 6-hexadecanoate in water at 25°C was reported to be 0.56 g(1)/100g ^{a,b} and 0.31 g(1)/100g ^{a,c} . The corresponding mass per cent and mole fractions, x_1 , values calculated by the compiler are 0.56 g(1)/100g sln, 0.31 g(1)/100g sln and 2.4×10^{-4} , 1.4×10^{-4} . The units g(1)/100g were reported as above.	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The analytical methods were used. An excess of ester was shaken with water until an equilibrium was obtained. The dissolved ester was determined both by titration with 0.1 N NaOH and by evaporation of water. At least two determinations were made.	SOURCE AND PURITY OF MATERIALS: (1) Not specified. (2) Not specified. ESTIMATED ERROR: Temp. $\pm 0.10^\circ C$. Soly. $\pm 0.5\%$ (precision of duplicates). REFERENCES:

<p>COMPONENTS:</p> <p>(1) 1,2-Benzenedicarboxylic acid bis(2-ethylhexyl) ester (bis(2-ethylhexyl) phthalate); $C_{24}H_{38}O_4$; [117-81-7]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Leyder, F.; Boulanger, P. <i>Bull. Environ. Contam. Toxicol.</i> <u>1983</u>, 30(2), 152-7.</p>
<p>VARIABLES:</p> <p>T/K = 293</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of 1,2-benzenedicarboxylic acid bis(2-ethylhexyl) ester in water at 20°C was reported to be 1.05×10^{-7} mol(1)/L and 4.1×10^{-5} g(1)/L.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/Apparatus/PROCEDURE:</p> <p>The analytical method as described in the <i>OECD Guidelines for Testing of Chemicals</i> (ref 1) was used. Gas chromatography analyses were conducted on a Girdel 3000 FFLE apparatus with flame-ionization detection. A 4 mm x 180 m stainless steel column packed with 3% OV-1 on 80-100 mesh Chromosorb W-AW-DMCS and helium at a 20mL/min flow rate were used. Samples were fortified with an internal standard, extracted, and concentrated to 1 mL of hexane solution. "Solvent flush" and "hot needle" injection techniques were used. The two methods gave similar results. Peak heights were measured throughout. UV measurements were made by using 1-cm cells in a Perkin-Elmer model 552 uv-Visible spectrophotometer.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Essochem DOP; purity not specified; used as received.</p> <p>(2) Deionized and distilled from $KMnO_4$.</p> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm 1^\circ C$.</p> <p>REFERENCES:</p> <p>1. <i>OECD Guidelines for Testing of Chemicals</i>, Paris, OECD, <u>1981</u>. (methods 101, 105).</p>

<p>COMPONENTS:</p> <p>(1) Phosphoric acid tris(2-ethylhexyl) ester (tri(2-ethylhexyl) phosphate); $C_{24}H_{51}O_4P$; [78-42-2]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Apelblat, A. <i>J. Chem. Soc. B</i> <u>1969</u>, 175-7.</p>
<p>VARIABLES:</p> <p>$T/K = 298$</p>	<p>PREPARED BY:</p> <p>A. Skrzecz</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of water in phosphoric acid tris(2-ethylhexyl) ester at 25°C was reported to be 0.81 mol(2)/L sln. The corresponding value on a mass/volume basis is 14.6 g(2)/L sln (compiler).</p> <p>The density of the ester-phase was reported to be d^{25} 0.9235.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The analytical method was used. The mixture was equilibrated by vigorous stirring for 30 min. and settling for 12 h. The organic phase was analyzed with the Karl Fischer reagent. Preliminary experiments showed that stirring and settling times were sufficient.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Albright and Wilson Co; purified by method described by Alcock, Grimley, Healy, Kennedy and McKay (ref 1); d_4^{25} 0.9220, n_D^{25} 1.4417.</p> <p>(2) Not specified.</p> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm 0.1^\circ C$.</p> <p>REFERENCES:</p> <p>1. Alcock, K.; Grimley, S.S.; Healy, T.V.; Kennedy, J.; McKay, H.A.C. <i>Trans. Faraday Soc.</i> <u>1956</u>, 52, 39.</p>

<p>COMPONENTS:</p> <p>(1) 1,2-Benzenedicarboxylic acid dinonyl ester (dinonyl phthalate); $C_{26}H_{42}O_4$; [84-76-4]</p> <p>(2) Water; H_2O; [7732-18-5]</p>	<p>ORIGINAL MEASUREMENTS:</p> <p>Krupatkin, I.L.; Glagoleva, M.F. <i>Zh. Prikl. Khim.</i> 1972, 45, 1317-20.</p>
<p>VARIABLES:</p> <p>$T/K = 298$</p>	<p>PREPARED BY:</p> <p>Z. Maczynska</p>
<p>EXPERIMENTAL VALUES:</p> <p>The solubility of 1,2-benzenedicarboxylic acid dinonyl ester in water at 25°C was reported to be 0.02 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 9×10^{-6}.</p> <p>The solubility of water in 1,2-benzenedicarboxylic acid dinonyl ester at 25°C was reported to be 0.09 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.02.</p>	
<p>AUXILIARY INFORMATION</p>	
<p>METHOD/APPARATUS/PROCEDURE:</p> <p>The titration method was used. The measurements were carried out in a water thermostat. The data were reported together with the ternary system 1,2-benzenedicarboxylic acid dinonyl ester-water-2-furancarboxal (dinonyl phthalate-water-furfural). No further details were reported in the paper.</p>	<p>SOURCE AND PURITY OF MATERIALS:</p> <p>(1) Source not specified; b.p. 240°C, d_4^{25} 0.9819, n_D^{25} 1.4852.</p> <p>(2) Twice distilled.</p> <hr/> <p>ESTIMATED ERROR:</p> <p>Temp. $\pm 0.1^\circ C$.</p> <hr/> <p>REFERENCES:</p>

COMPONENTS: (1) Benzoic acid, 2-hydroxy-, cinchonan-9-ol, 6'-methoxy, (8 α , 9R) ester (quinine salicylate); $C_{27}H_{30}N_2O_5$; [750-90-3] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Seidell, A. <i>Hygienic Lab. Bull.</i> <u>1910</u> , No. 67, 98 pp (US Govt. Printing Office, Washington, DC).
VARIABLES: $T/K = 298$	PREPARED BY: G.T. Hefter
EXPERIMENTAL VALUES: The solubility of 2-hydroxybenzoic acid 6'-methoxycinchonan-9-ol' (8 α , 9R) ester in water at 25°C was reported to be 0.065 g(1)/100g sln. The corresponding mole fraction, x_1 , value calculated by the compiler is 2.5×10^{-5} .	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: The analytical method was used. Appropriate quantities of (1) and (2) were shaken together. The amount of dissolved (1) was determined by evaporation of a known quantity of the saturated solution. The residue was dried to a constant weight in a vacuum desiccator.	SOURCE AND PURITY OF MATERIALS: (1) Merck, m.p. ca. 195°C. (2) Distilled (no details given). ESTIMATED ERROR: Temp. not stated. Soly. not stated. REFERENCES:

COMPONENTS: (1) 1,2-Benzenedicarboxylic acid didodecyl ester (<i>didodecyl phthalate</i>); $C_{32}H_{54}O_4$; [2432-90-8] (2) Water; H_2O ; [7732-18-5]	ORIGINAL MEASUREMENTS: Krupatkin, I.L.; Glagoleva, M.F. <i>Zh. Prikl. Khim.</i> <u>1972</u> , 45, 1317-20.
VARIABLES: $T/K = 298$	PREPARED BY: Z. Maczynska
EXPERIMENTAL VALUES: <p>The solubility of 1,2-benzenedicarboxylic acid didodecyl ester in water at 25°C was reported to be 0.07 g(1)/100g sln. The corresponding mole fraction, x_1, value calculated by the compiler is 3×10^{-5}.</p> <p>The solubility of water in 1,2-benzenedicarboxylic acid didodecyl ester at 25°C was reported to be 0.02 g(2)/100g sln. The corresponding mole fraction, x_2, value calculated by the compiler is 0.006.</p>	
AUXILIARY INFORMATION	
METHOD/APPARATUS/PROCEDURE: <p>The titration method was used. The measurements were carried out in a water thermostat. The data were reported together with the ternary system 1,2-benzenedicarboxylic acid didodecyl ester-water-2-furan-carbonyl (<i>didodecyl phthalate-water-furfural</i>). No further details were reported in the paper.</p>	SOURCE AND PURITY OF MATERIALS: (1) Source not specified; d_4^{25} 0.9548, n_D^{25} 1.4839. (2) Twice distilled. ESTIMATED ERROR: Temp. $\pm 0.1^\circ C$. REFERENCES:

SYSTEM INDEX

Pages preceded by E refer to evaluation texts whereas those not preceded by E refer to compiled tables.

Acetic acid 2-(2-butoxyethoxy)ethyl ester	E169, E170, 171, 172
Acetic acid 2-butoxyethyl ester	123
see ethanol, 2-butoxy-, acetate	
Acetic acid cyclohexyl ester	E85-E87, 88-90
Acetic acid 1,3-dimethylbutyl ester	97
Acetic acid 2-ethyl-1-butyl ester	98
Acetic acid 2-ethylhexyl ester	161
Acetic acid heptyl ester	140
Acetic acid hexyl ester	E99-E101, 102-104
Acetic acid 3-methoxy-1-butyl ester	72
Acetic acid 1-methyl-1-butyl ester	E6, 7, 8
Acetic acid 3-methyl-1-butyl ester	E9-E14, 15-24
Acetic acid 1-methylpentyl ester	E105, 106, 107
Acetic acid isononyl ester	179
Acetic acid octyl ester	162
Acetic acid pentyl ester	E25-E30, 31-40
Acetic acid phenyl ester	74
Acetic acid phenylmethyl ester	E124, E125, 126, 127
L-Ascorbic acid 6-hexadecanoate	232
see L-ascorbic acid 6-hexadecanoate	
1,2-Benzenedicarboxylic acid bis(2-ethylhexyl) ester	233
1,2-Benzenedicarboxylic acid bis(1-methylethyl) ester	214
1,2-Benzenedicarboxylic acid bis(2-methylpropyl) ester	218
1,2-Benzenedicarboxylic acid butyl octyl ester	231
1,2-Benzenedicarboxylic acid butyl phenylmethyl ester	E227, E228, 229, 230
1,2-Benzenedicarboxylic acid dibutyl ester	E219, E220, 221-223
1,2-Benzenedicarboxylic acid didodecyl ester	237
1,2-Benzenedicarboxylic acid diethyl ester	185, 186
1,2-Benzenedicarboxylic acid dimethyl ester	E153, 154, 155
1,2-Benzenedicarboxylic acid dinonyl ester	235
1,2-Benzenedicarboxylic acid dipentyl ester	225
1,2-Benzenedicarboxylic acid di-2-propenyl ester	213
1,2-Benzenedicarboxylic acid dipropyl ester	215
Benzoic acid ethyl ester	E128-E130, 131, 132
Benzoic acid, 2-hydroxy-, cinchonan-9-ol, 6'-methoxy, (8 α ,9R) ester	236
Benzoic acid, 2-hydroxy-, ethyl ester	E133
Benzoic acid, 2-hydroxy-, methyl ester	E81, E82, 83, 84
Benzoic acid, 2-hydroxy-, phenyl ester	E208, 209, 210
Benzoic acid methyl ester	E75-E77, 78-80
Benzyl acetate	
see acetic acid phenylmethyl ester	
Bis(2-ethylhexyl) phthalate	
see 1,2-benzenedicarboxylic acid bis(2-ethylhexyl) ester	
Butanedioic acid diethyl ester	E91, E92, 93, 94
2,3-Butanediol diacetate	95
Butanoic acid butyl ester	E108-E110, 111-114
Butanoic acid 1-methylethyl ester	E41-E44, 45, 46
Butanoic acid 3-methyl-1-butyl ester	141
Butanoic acid, 3-methyl-, ethyl ester	E47-E49, 50-52
Butanoic acid pentyl ester	E142, E143, 144, 145
Butanoic acid propyl ester	E53, E54, 55, 56
2-(2-Butoxyethoxy)ethyl acetate	
see ethanol, 2-(2-butoxyethoxy)-, acetate	
2-Butoxyethyl acetate	
see ethanol, 2-butoxy-, acetate	

- Butyl acrylate
 see 2-propenoic acid butyl ester
 Butyl benzyl phthalate
 see 1,2-benzenedicarboxylic acid butyl phenylmethyl ester
 Butyl butyrate
 see butanoic acid butyl ester
 Butyl isobutyrate
 see propanoic acid, 2-methyl-, butyl ester
 Butyl dibutylphosphinate
 see phosphinic acid, dibutyl-, butyl ester
 Butyl dipropylphosphinate
 see phosphinic acid, dipropyl-, butyl ester
 2,3-Butylene diacetate
 see 2,3-butanediol diacetate
 Butyl lactate
 see propanoic acid, 2-hydroxy-, 1-butyl ester
 Butyloctyl phthalate
 see 1,2-benzene dicarboxylic acid butyl octyl ester
 Butyl propionate
 see propanoic acid butyl ester
 Butyl valerate
 see pentanoic acid 1-butyl ester

 2-Cyanoethyl methacrylate
 see 2-propenoic acid, 2-methyl-, 2-cyanoethyl ester
 Cyclohexyl acetate
 see acetic acid cyclohexyl ester
 Cyclohexyl formate
 see formic acid cyclohexyl ester
 Cyclohexyl propionate
 see propanoic acid cyclohexyl ester

 Decanedioic acid diethyl ester 216
 Decanoic acid ethyl ester 188
 Decanoic acid methyl ester 180
 Diallyl phthalate
 see 1,2-benzenedicarboxylic acid di-2-propenyl ester
 Dibutyl butylphosphinate
 see phosphinic acid, butyl-, dibutyl ester
 Dibutyl ethylphosphonate
 see phosphonic acid, ethyl-, dibutyl ester
 Dibutyl methylphosphonate
 see phosphonic acid, methyl-, dibutyl ester
 Dibutyl propylphosphonate
 see phosphonic acid, propyl-, dibutyl ester
 Dibutyl tolyl phosphate
 see phosphoric acid dibutyl methylphenyl ester
 Didodecyl phthalate
 see 1,2-benzenedicarboxylic acid didodecyl ester
 Diethyl adipate
 see hexanedioic acid diethyl ester
 Diethyl decanedioate
 see decanedioic acid diethyl ester
 Diethyl glutarate
 see pentanedioic acid diethyl ester
 Diethyl heptanedioate
 see heptanedioic acid diethyl ester
 Diethyl malonate
 see malonic acid diethyl ester
 Diethylnonanedioate
 see nonanedioic acid diethyl ester
 Diethyl octanedioate
 see octanedioic acid diethyl ester
 Diethyl phthalate
 see 1,2-benzenedicarboxylic acid diethyl ester
 Diethyl succinate
 see butanedioic acid diethyl ester

Diisopropyl phthalate	
see 1,2-benzenedicarboxylic acid bis(1-methylethyl) ester	
Dimethyl adipate	
see hexanedioic acid dimethyl ester	
1,3-Dimethylbutyl acetate	
see acetic acid 1,3-dimethylbutyl ester	
Dimethyl glutarate	
see pentanedioic acid dimethyl ester	
Dimethyl phthalate	
see 1,2-benzenedicarboxylic acid dimethyl ester	
Dinonyl phthalate	
see 1,2-benzenedicarboxylic acid dinonyl ester	
Dipropyl phthalate	
see 1,2-benzenedicarboxylic acid dipropyl ester	
Ethanol, 2-butoxy-, acetate	
see acetic acid 2-butoxyethyl ester	
Ethanol, 2(2-butoxyethyl)-, acetate	
see acetic acid 2-(2-butoxyethoxy)ethyl ester	
Ethanol, 2-butoxy-, phosphate (3:1)	
see Phosphoric acid tris(butoxyethyl) ester	
Ethyl benzoate	
see benzoic acid ethyl ester	
2-Ethylbutyl acetate	
see acetic acid 2-ethylbutyl ester	
Ethyl caproate	
see hexanoic acid ethyl ester	
Ethyl caprylate	
see octanoic acid ethyl ester	
Ethyl decanoate	
see decanoic acid ethyl ester	
Ethyl dibutylphosphinate	
see phosphinic acid, dibutyl-, ethyl ester	
Ethyl enanthate	
see heptanoic acid ethyl ester	
Ethyl heptanoate	
see heptanoic acid ethyl ester	
2-Ethylhexyl acetate	
see acetic acid 2-ethylhexyl ester	
Ethyl isovalerate	
see butanoic acid, 3-methyl-, ethyl ester	
Ethyl nonanoate	
see nonanoic acid ethyl ester	
Ethyl salicylate	
see benzoic acid, 2-hydroxy-, ethyl ester	
Ethyl trimethylacetate	
see propanoic acid, 2,2-dimethyl-, ethyl ester	
Ethyl valerate	
see pentanoic acid ethyl ester	
Formic acid cyclohexyl ester	2
Formic acid heptyl ester	115
Formic acid hexyl ester	57
Formic acid octyl ester	146
Glycerol triacetate	
see 1,2,3-propanetriol triacetate	
Heptanedioic acid diethyl ester	178
Heptanoic acid ethyl ester	147
Heptanoic acid methyl ester	116
Heptyl acetate	
see acetic acid heptyl ester	
Heptyl formate	
see formic acid heptyl ester	
Hexanedioic acid diethyl ester	E156, E157, 158-160
Hexanedioic acid dimethyl ester	96
Hexanoic acid ethyl ester	117, 118

Hexanoic acid methyl ester	58
Hexyl acetate	
see acetic acid hexyl ester	
Hexyl formate	
see formic acid hexyl ester	
Hexyl isobutyrate	
see propanoic acid, 2-methyl-, hexyl ester	
Hexyl propionate	
see propanoic acid hexyl ester	
Isobutyl isobutyrate	
see propanoic acid, 2-methyl-, 2-methylpropyl ester	
Isononyl acetate	
see acetic acid isononyl ester	
Isopentyl acetate	
see acetic acid 3-methylbutyl ester	
Isopentyl butyrate	
see butanoic acid 3-methylbutyl ester	
Isopentyl propionate	
see propanoic acid 3-methylbutyl ester	
Isopropyl butyrate	
see butanoic acid 1-methylethyl ester	
Malonic acid diethyl ester	
see propanedioic acid diethyl ester	
3-Methoxybutyl acetate	
see acetic acid 3-methoxybutyl ester	
Methyl benzoate	
see benzoic acid methyl ester	
1-Methylbutyl acetate	
see acetic acid 1-methyl-1-butyl ester	
Methyl caproate	
see hexanoic acid methyl ester	
Methyl decanoate	
see decanoic acid methyl ester	
Methyl dibutylphosphinate	
see phosphinic acid dibutylmethyl ester	
Methyl diphenyl phosphate	
see phosphoric acid, methyl-, diphenyl ester	
Methyl enanthate	
see heptanoic acid methyl ester	
Methyl hexanoate	
see hexanoic acid methyl ester	
Methyl nonanoate	
see nonanoic acid methyl ester	
1-Methylpentyl acetate	
see acetic acid 1-methylpentyl ester	
Methyl salicylate	
see benzoic acid, 2-hydroxy-, methyl ester	
Nonanedioic acid diethyl ester	212
Nonanoic acid ethyl ester	181
Nonanoic acid methyl ester	163
Octanedioic acid diethyl ester	187
Octanoic acid ethyl ester	E164, E165, 166, 167
Octyl acetate	
see acetic acid octyl ester	
Octyl formate	
see formic acid octyl ester	
Pentanedioic acid diethyl ester	139
Pentanedioic acid dimethyl ester	5
Pentanoic acid 1-butyl ester	148
Pentanoic acid ethyl ester	E59, E60, 61, 62
Pentyl acetate	
see acetic acid pentyl ester	

Pentyl butyrate	
see butanoic acid pentyl ester	
Pentyl propionate	
see propanoic acid pentyl ester	
Phenyl acetate	
see acetic acid phenyl ester	
Phenyl salicylate	
see benzoic acid,2-hydroxy-, phenyl ester	
Phosphinic acid, dibutyl-, butyl ester	189-191
Phosphinic acid, dibutyl-, ethyl ester	173, 174
Phosphinic acid, dibutyl-, methyl ester	150, 151
Phosphinic acid, dibutyl-, propyl ester	182, 183
Phosphinic acid, dipropyl-, butyl ester	175, 176
Phosphonic acid, butyl-, dibutyl ester	192, 193
Phosphonic acid, ethyl-, dibutyl ester	177
Phosphonic acid, propyl-, dibutyl ester	184
Phosphonic acid, methyl-, dibutyl ester	152
Phosphoric acid butyl bis(methylphenyl) ester	224
Phosphoric acid dibutyl methylphenyl ester	217
Phosphoric acid methyl diphenyl ester	211
Phosphoric acid tributyl ester	E194-E198, 199-206
Phosphoric acid tris(butoxyethyl) ester	226
Phosphoric acid tris(2-ethylhexyl) ester	234
Phosphoric acid tris(2-methylpropyl) ester	207
Propanedioic acid diethyl ester	4
Propanoic acid butyl ester	E63-E65, 66-70
Propanoic acid cyclohexyl ester	138
Propanoic acid, 2,2-dimethyl-, ethyl ester	71
Propanoic acid hexyl ester	149
Propanoic acid, 2-hydroxy-, 1-butyl ester	73
Propanoic acid, 2-methyl-, butyl ester	119
Propanoic acid 3-methyl-1-butyl ester	120
Propanoic acid, 2-methyl-, hexyl ester	168
Propanoic acid, 2-methyl-, 2-methylpropyl ester	121
Propanoic acid pentyl ester	122
1,2,3-Propanetriol triacetate	E134, E135, 136, 137
2-Propenoic acid butyl ester	3
2-Propenoic acid, 2-methyl-, 2-cyanoethyl ester	1
Propyl butyrate	
see butanoic acid propyl ester	
Propyl dibutylphosphinate	
see phosphinic acid dibutylpropyl ester	
Quinine monosalicylate	
see Benzoic acid, 2-hydroxy-, cinchonan-9-ol, 6'-methoxy, (8 α ,9R) ester	
Tributyl phosphate	
see phosphoric acid tributyl ester	
Tri(2-ethylhexyl) phosphate	
see phosphoric acid tris(2-ethylhexyl) ester	
Triisobutyl phosphate	
see phosphoric acid tris(2-methylpropyl) ester	

REGISTRY NUMBER INDEX

Pages preceded by E refer to evaluation texts whereas those not preceded by E refer to compiled tables.

78-42-2	234
78-46-6	192, 193
78-51-3	226
84-66-2	185, 186
84-69-5	218
84-74-2	E219, E220, 221-223
84-76-4	235
84-78-6	231
85-68-7	E227, E228, 229, 230
93-58-3	E75-E77, 78-80
93-89-0	E128-130, 131, 132
97-85-8	121
97-87-0	119
102-76-1	E134, E135, 136, 137
103-09-3	161
105-53-3	4
105-66-8	E53, E54, 55, 56
105-68-0	120
106-27-4	141
106-30-9	147
106-32-1	E164, E165, 166, 167
106-70-7	58
106-73-0	116
108-64-5	E47-E49, 50-52
108-84-9	97
109-21-7	E108-E110, 111-114
110-38-3	188
110-40-7	216
110-42-9	180
112-06-1	140
112-07-2	123
112-14-1	162
112-23-2	115
112-32-3	146
115-89-9	211
117-81-7	233
118-55-8	E208, 209, 210
118-61-6	133
119-36-8	E81, E82, 83, 84
122-79-2	74
123-25-1	E91, E92, 93, 94

123-29-5	181
123-66-0	117, 118
123-92-2	E9-E14, 15-24
124-17-4	E169, E170, 171, 172
126-71-6	207
126-73-8	E194-E198, 199-206
131-11-3	E153, 154, 155
131-16-8	215
131-17-9	213
131-18-0	225
137-66-6	232
138-22-7	73
140-11-4	E124, E125, 126, 127
141-28-6	E156, E157, 158-160
141-32-2	3
142-92-7	E99-E101, 102-104
539-82-2	E59, E60, 61, 62
540-18-1	E142, E143, 144, 145
590-01-2	E63-E65, 66-70
591-68-4	148
605-45-8	214
622-45-7	E85-E87, 88-90
624-17-9	212
624-54-4	122
626-38-0	E6, 7, 8
627-93-0	96
628-63-7	E25-E30, 31-40
629-33-4	57
638-11-9	E41-E44, 45, 46
750-90-3	236
818-38-2	139
1114-92-7	95
1119-40-0	5
1731-84-6	163
2050-20-6	178
2050-23-9	187
2349-07-7	168
2404-58-2	177
2404-73-1	152
2432-90-8	237
2445-76-3	149
2950-47-2	189-191
3938-95-2	71
4351-54-6	2
4435-53-4	72
4513-53-5	1
4628-12-0	184
5953-49-1	E105, 106, 107
6222-35-1	138

7100-92-7	173, 174
7100-93-8	182, 183
7163-67-9	150, 151
7732-18-5	1-5, E6, 7, 8, E9-E14, 15-24, E25-E30, 31-40, E41-E44, 45, 46, E47-E49, 50-52, E53, E54, 55-58, E59, E60, 61, 62, E63-E65, 66-74, E75-E77, 78-80, E81, E82, 83, 84, E85-E87, 88-90, E91, E92, 93- 98, E99-E101, 102-104, E105, 106, 107, E108-E110, 111-123, E124, E125, 126, 127, E128-E130, 131- 133, E134, E135, 136-141, E142, E143, 144-152, E153, 154, 155, E156, E157, 158-163, E164, E165, 166-168, E169, E170, 171-193, 154, 155, E156, E157, 158-163, E164, E165, 166-168, E170, 171-193, E194-E198, 199-207, E208, 209-218, E219, E220, 221-226, E227, E228, 229-237
10031-87-5	98
16984-10-4	175, 176
25657-10-7	224
26446-69-5	217
40379-24-6	179

AUTHOR INDEX

Pages preceded by E refer to evaluation texts whereas those not preceded by E refer to compiled tables.

Alcock, K.	E194-E198, 199
Alvarez, J.R.	E99-E101, 103
Andreeva, N.G.	E9-E13, 20
Apelblat, A.	207, 211, 217, 224, 226, 234
Balashov, M.I.	E108-E110, 113
Baldwin, W.H.	E134, E135, 137, E194-E198, 201
Bancroft, W.D.	E9-E13, 15, E47-E49, 50
Bergen, W.S.	95
Bomshtein, A.L.	E41-E44, 45, E59, E60, 62, E63-E65, 69, 148
Boulangier, P.	E153, 155, 186, 213-215, 218, E219, E220, 223, 225, E227, E228, 230, 233
Bridgman, J.A.	E63-E65, 66, E108-E110, 111
Bruce, F.R.	E194-E198, 200
Bruins, P.F.	95
Burger, L.L.	E194-E198, 200
Chang, Y.C.	E47-E49, 51
Chari, K.S.	E9-E13, 17, 19
Chebotaev, V.F.	E108-E110, 113
Chiranjivi, C.	E9-E13, 21
Chubarov, G.A.	3
Danov, S.M.	3
Dehn, W.M.	E208, 210
Doolittle, A.K.	E6, 8, E25-E29, 33, E63-E65, 68, 72, 73, E85-E87, 97, 98, E105, 107, E108-E110, 112, 123, 161, E169, E170, 171
Durasov, V.B.	150-152, 173-177, 182-184, 189-193, E194-E198, 205, 206
Dyadin, Yu.A.	150-152, 173-177, 182-184, 189-193, E194-E198, 205, 206
Erykov, V.G.	1
Fairhurst, D.	E194-E198, 204
Forsman, R.C.	E194-E198, 200
Frolov, A.F.	1
Fuehner, H.	E53, E54, 55
Fuse, K.	E9-E13, 18
Garber, Yu.N.	E9-E13, 20
Gilbert, E.C.	E75-E77, 78
Glagoleva, M.F.	E128-E130, 131, E219, E220, 221, E227, E228, 229, 231, 235, 237
Gorman, M.A.	E156, E157, 159
Goto, S.	E124, E125, 126
Greenspan, F.P.	E153, 154, 185
Grimley, S.S.	E194-E198, 199
Gross, P.M.	E156, E157, 159
Hardy, C.J.	E194-E198, 204
Hasegawa, T.	E194-E198, 202
Healy, T.V.	E194-E198, 199
Hemptinne, A.	E25-E29, 31, E59, E60, 61, 122, E142, E143, 144
Higgins, C.E.	E194-E198, 201
Hopkins, M.B.	E6, 7, E25-E29, 32, E63-E65, 67, E105, 106
Husain, A.	E9-E13, 18

Iguchi, A.	E9-E13, 18
Kahn, J.	4, E91, E92, 93, 117, 139, 147, E156, E157, 158, E164, E165, 166, 178, 181, 187, 188, 212, 216
Kendall, J.	E194-E198, 199
Kharachencho, S.K.	E194-E198, 203
King, C.J.	E99-E101, 102
Komarova, L.F.	E9-E13, 20
Korshukov, M.A.	1
Kraus, K.A.	E134, E135, 137
Krupatkin, I.L.	E25-E29, 37, E128-E130, 131, E219, E220, 221, E227, E228, 229, 231, 235, 237
Lauer, B.E.	E75-77, 78
Leyder, F.	E153, 155, 186, 213-215, 218, E219, E220, 223, 225, E227, E228, 230, 233
Logutov, V.I.	3
Maiorova, N.M.	1
Martire, D.E.	163, 180
Massaldi, H.A.	E99-E101, 102
Matsubara, M.	E124, E125, 126
McKay, H.A.C.	E194-E198, 199, 204
Miller, M.M.	163, 180
Mikhailov, V.A.	E194-E198, 203
Mironova, Z.N.	189, 190, 192, 193
Moulton, R.W.	E47-E49, 51
Narasimhan, K.S.	E9-E13, 17
Nazin, A.G.	E194-E198, 203
Neila, J.J.	E99-E101, 103
Nikolaev, A.V.	150-152, 173-177, 182-184, 189-193, E194-E198, 205, 206
Othmer, D.F	E9-E13, 16, E85-E87, 89, 95, E134, E135, 136, 162, E169, E170, 172
Park, J.G.	E6, 7, E25-E29, 32, E63-E65, 67, E105, 106
Ramanarao, M.V.	E9-E13, 19
Rao, C.V.	E25-E29, 34-36
Rao, D.S.	E9-E13, 21
Rao, J.R.	E25-E29, 34-36
Rao, K.V.	E9-E13, 19
Rardon, R.J.	E134, E135, 137
Reddy, C.C.	E9-E13, 17
Richon, D.	E25-E30, 39, 74, E75-E77, 79
Saylor, J.H.	E156, E157, 159
Schlechter, N.	95
Schlesinger, N.	95
Schwarz, F.P.	E219, E220, 222
Seidell, A.	E81, E82, 83, E208, 209, 236
Serafimov, L.A.	E41-E44, 45, E59, E60, 62, E63-E65, 69, E108-E110, 113, 148
Shanley, E.S.	E153, 154, 185
Shcherbakova, T.A.	E25-E29, 37
Skrzecz, A.	E9-E14, 22, 23, E25-E30, 38
Sobotka, H.	4, E91, E92, 93, 117, 139, 147, E156, 157, 158, E164, E165, 166, 178, 181, 187, 188, 212, 216
Soldano, B.A.	E194-E198, 201
Stephenson, R.	2, 5, E9-E14, 24, E25-E30, 40, E41-E44, 46, E47-E49, 52, E53, E54, 56-58, E63-E65, 70, 71, E75-E77, 80, E81, E82, 84, E85-E87, 90, E91, E92, 94, 96, E99-E101, 104, E108-E110, 114-116, 118-121, E124, E125, 127, E128-E130, 132, 133, 138, 140,

Stephenson, R. 141, E142, E143, 145, 146, 149, E156, E157, 160, E164, E165, 168, 179

Stuart, J. 2, 5, E9-E14, 24, E25-E30, 40, E41-E44, 46, E47-E49, 52, E53, E54, 56-58, E63-E65, 70, 71, E75-E77, 80, E81, E82, 84, E85-E87, 90, E91, E92, 94, 96, E99-E101, 104, E108-E110, 114-116, 118-121, E124, E125, 127, E128-E130, 132, 133, 138, 140, 141, E142, E143, 145, 146, 149, E156, E157, 160, E164, E165, 168, 179

Swern, D. 232

Tewari, Y.B. 163, 180

Tomashahuk, V.I. 3

Trofimov, A.N. E41-E44, 45, E59, E60, 62, E63-E65, 69, 148

Trueger, E. E9-E13, 16, E85-E87, 89, E134, E135, 162, E169, E170, 172

Venkataratnam, A. E25-E29, 35

Viallard, A. E25-E30, 39, 74, E75-E77, 79

Washino, K. E124, E125, 126

Wasik, S.P. 163, 180

White, R.E. E9-E13, 16, E85-E87, 89, E134, E135, 162, E169, E170, 172

Willson, A.M. E194-E198, 204

Yakovlev, I.I. 150-152, 173-177, 182-184, 189-193, E194-E198, 205, 206

Yakovleva, N.I. 150-152, 173-177, 182-184, 191

SOLUBILITY DATA SERIES

- Volume 1 H. L. Clever, *Helium and Neon*
- Volume 2 H. L. Clever, *Krypton, Xenon and Radon*
- Volume 3 M. Salomon, *Silver Azide, Cyanide, Cyanamides, Cyanate, Selenocyanate and Thiocyanate*
- Volume 4 H. L. Clever, *Argon*
- Volume 5/6 C. L. Young, *Hydrogen and Deuterium*
- Volume 7 R. Battino, *Oxygen and Ozone*
- Volume 8 C. L. Young, *Oxides of Nitrogen*
- Volume 9 W. Hayduk, *Ethane*
- Volume 10 R. Battino, *Nitrogen and Air*
- Volume 11 B. Scrosati and C. A. Vincent, *Alkali Metal, Alkaline Earth Metal and Ammonium Halides, Amide Solvents*
- Volume 12 C. L. Young, *Sulfur Dioxide, Chlorine, Fluorine and Chlorine Oxides*
- Volume 13 S. Siekierski, T. Mioduski and M. Salomon, *Scandium, Yttrium, Lanthanum and Lanthanide Nitrates*
- Volume 14 H. Miyamoto, M. Salomon and H. L. Clever, *Alkaline Earth Metal Halates*
- Volume 15 A. F. M. Barton, *Alcohols with Water*
- Volume 16/17 E. Tomlinson and A. Regosz, *Antibiotics: I, β -Lactam Antibiotics*
- Volume 18 O. Popovych, *Tetraphenylborates*
- Volume 19 C. L. Young, *Cumulative Index: Volumes 1-18*
- Volume 20 A. L. Horvath and F. W. Getzen, *Halogenated Benzenes, Toluenes and Phenols with Water*
- Volume 21 C. L. Young and P. G. T. Fogg, *Ammonia, Amines, Phosphine, Arsine, Stibine, Silane, Germane and Stannane in Organic Solvents*
- Volume 22 T. Mioduski and M. Salomon, *Scandium, Yttrium, Lanthanum and Lanthanide Halides in Nonaqueous Solvents*
- Volume 23 T. P. Dirkse, *Copper, Silver, Gold and Zinc, Cadmium, Mercury Oxides and Hydroxides*
- Volume 24 W. Hayduk, *Propane, Butane and 2-Methylpropane*
- Volume 25 C. Hirayama, Z. Galus and C. Guminski, *Metals in Mercury*
- Volume 26 M. R. Masson, H. D. Lutz and B. Engelen, *Sulfites, Selenites and Tellurites*
- Volume 27/28 H. L. Clever and C. L. Young, *Methane*
- Volume 29 H. L. Clever, *Mercury in Liquids, Compressed Gases, Molten Salts and Other Elements*
- Volume 30 H. Miyamoto and M. Salomon, *Alkali Metal Halates, Ammonium Iodate and Iodic Acid*
- Volume 31 J. Eysseltová and T. P. Dirkse, *Alkali Metal Orthophosphates*

- Volume 32 P. G. T. Fogg and C. L. Young, *Hydrogen Sulfide, Deuterium Sulfide and Hydrogen Selenide*
- Volume 33 P. Franzosini, *Molten Alkali Metal Alkanoates*
- Volume 34 A. N. Paruta and R. Piekos, *4-Aminobenzenesulfonamides. Part I: Non-cyclic Substituents*
- Volume 35 A. N. Paruta and R. Piekos, *4-Aminobenzenesulfonamides. Part II: 5-Membered Heterocyclic Substituents*
- Volume 36 A. N. Paruta and R. Piekos, *4-Aminobenzenesulfonamides. Part III: 6-Membered Heterocyclic Substituents and Miscellaneous Systems*
- Volume 37 D. G. Shaw, *Hydrocarbons with Water and Seawater. Part I: Hydrocarbons C₅ to C₇*
- Volume 38 D. G. Shaw, *Hydrocarbons with Water and Seawater. Part II: Hydrocarbons C₈ to C₃₆*
- Volume 39 C. L. Young, *Cumulative Index: Volumes 20-38*
- Volume 40 J. Hala, *Halides, Oxyhalides and Salts of Halogen Complexes of Titanium, Zirconium, Hafnium, Vanadium, Niobium and Tantalum*
- Volume 41 C.-Y. Chan, I. N. Lepeshkov and K. H. Khoo, *Alkaline Earth Metal Perchlorates*
- Volume 42 P. G. T. Fogg and W. Gerrard, *Hydrogen Halides in Non-aqueous Solvents*
- Volume 43 R. W. Cargill, *Carbon Monoxide*
- Volume 44 H. Miyamoto, E. M. Woolley and M. Salomon, *Copper and Silver Halates*
- Volume 45/46 R. P. T. Tomkins and N. P. Bansal, *Gases in Molten Salts*
- Volume 47 R. Cohen-Adad and J. W. Lorimer, *Alkali Metal and Ammonium Chlorides in Water and Heavy Water (Binary Systems)*
- Volume 48 F. Getzen, G. Hefter and A. Maczynski, *Esters with Water. Part I: Esters 2-C to 6-C*
- Volume 49 F. Getzen, G. Hefter and A. Maczynski, *Esters with Water. Part II: Esters 7-C to 32-C*